

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C12N 15/12, C12Q 1/68, C07K 14/47, 16/18, G01N 33/68, A61K 38/17		A2	(11) International Publication Number: WO 00/44900 (43) International Publication Date: 3 August 2000 (03.08.00)
(21) International Application Number: PCT/US00/02237 (22) International Filing Date: 28 January 2000 (28.01.00) (30) Priority Data: 60/117,905 29 January 1999 (29.01.99) US 60/117,904 29 January 1999 (29.01.99) US (63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Applications US 60/117,904 (CIP) Filed on 29 January 1999 (29.01.99) US 60/117,905 (CIP) Filed on 29 January 1999 (29.01.99) (71) Applicant (for all designated States except US): INCYTE PHARMACEUTICALS, INC. [US/US]; 3174 Porter Drive, Palo Alto, CA 94304 (US). (72) Inventors; and (75) Inventors/Applicants (for US only): TANG, Y., Tom [CN/US]; 4230 Ranwick Court, San Jose, CA 95118 (US). LAL, Pretti [IN/US]; 2382 Lass Drive, Santa Clara, CA 95054 (US). HILLMAN, Jennifer, L. [US/US]; 230 Monrow Drive #12, Mountain View, CA 94040 (US). YUE, Henry [US/US]; 826		Lois Avenue, Sunnyvale, CA 94087 (US). AZIMZAI, Yalda [US/US]; 2045 Rock Springs Drive, Hayward, CA 94545 (US). LU, Aina, M., D. [US/US]; 55 Park Belmont Place, San Jose, CA 95136 (US). BAUGHN, Mariah, R. [US/US]; 14244 Santiago Road, San Leandro, CA 94577 (US). TRAN, Bao [US/US]; 744 Kiely Boulevard, Santa Clara, CA 95051 (US). SHIH, Leo, L. [US/US]; Apartment B., 1081 Tanland Drive, Palo Alto, CA 94303 (US). AU-YOUNG, Janice, L. [US/US]; 1419 Kains Avenue, Berkeley, CA 94702 (US). (74) Agents: HAMLET-COX, Diana et al.; Incyte Pharmaceuticals, Inc., 3174 Porter Drive, Palo Alto, CA 94304 (US). (81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>	
(54) Title: NUCLEIC-ACID BINDING PROTEINS (57) Abstract <p>The invention provides human nucleic-acid binding proteins (NuABP) and polynucleotides which identify and encode NuABP. The invention also provides expression vectors, host cells, antibodies, agonists, and antagonists. The invention also provides methods for diagnosing, treating, or preventing disorders associated with expression of NuABP.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

NUCLEIC-ACID BINDING PROTEINS

TECHNICAL FIELD

This invention relates to nucleic acid and amino acid sequences of nucleic-acid binding proteins and to the use of these sequences in the diagnosis, treatment, and prevention of reproductive, immune, and neurological disorders, and cell proliferative disorders including cancer.

5

BACKGROUND OF THE INVENTION

Multicellular organisms are comprised of diverse cell types that differ dramatically both in structure and function. The identity of a cell is determined by its characteristic pattern of gene expression, and different cell types express overlapping but distinct sets of genes throughout development. Spatial and temporal regulation of gene expression is critical for the control of cell proliferation, cell differentiation, apoptosis, and other processes that contribute to organismal development. Furthermore, gene expression is regulated in response to extracellular signals that mediate cell-cell communication and coordinate the activities of different cell types. Appropriate gene regulation also ensures that cells function efficiently by expressing only those genes whose functions are required at a given time.

Transcriptional regulatory proteins are essential for the control of gene expression. Some of these proteins function as transcription factors that initiate, activate, repress, or terminate gene transcription. Transcription factors generally bind to promoter, enhancer, or upstream regulatory regions of a gene in a sequence-specific manner, although some factors bind regulatory elements within or downstream of the coding region. Transcription factors may bind to a specific region of DNA singly or as a complex with other accessory factors. (Reviewed in Lewin, B. (1990) Genes IV, Oxford University Press, New York, NY, pp. 554-570.)

The double helix structure and repeated sequences of DNA create topological and chemical features which can be recognized by transcription factors. These features include hydrogen bond donor and acceptor groups, hydrophobic patches, major and minor grooves, and regular repeated stretches of sequence which induce distinct bends in the helix. Typically, transcription factors recognize specific DNA sequence motifs of about 20 nucleotides in length. Multiple adjacent transcription factor-binding motifs may be required for gene regulation.

Many transcription factors incorporate DNA-binding structural motifs which comprise either α helices or β sheets that bind to the major groove of DNA. Four well-characterized structural motifs are helix-turn-helix, zinc finger, leucine zipper, and helix-loop-helix. Proteins containing these motifs may act alone as monomers or form homo- or heterodimers that interact with DNA.

The helix-turn-helix motif consists of two α helices connected at a fixed angle by a short

chain of amino acids. One of the helices binds to the major groove. Helix-turn-helix motifs are exemplified by the homeobox motif which is present in homeodomain proteins. These proteins are critical for specifying the anterior-posterior body axis during development and are conserved throughout the animal kingdom. The Antennapedia and Ultrabithorax proteins of Drosophila melanogaster are prototypical homeodomain proteins. (Pabo, C.O. and R.T. Sauer (1992) Ann. Rev. Biochem. 61:1053-1095.)

The zinc finger motif, which binds zinc ions, generally contains tandem repeats of about 30 amino acids consisting of periodically spaced cysteine and histidine residues. Examples of this sequence pattern include the C2H2-type and the C3HC4-type zinc fingers, and the PHD domain. (Lewin, supra ; Aasland, R., et al. (1995) Trends Biochem. Sci 20:56 - 59.) Zinc finger proteins each contain an α helix and an antiparallel β sheet whose proximity and conformation are maintained by the zinc ion. Contact with DNA is made by the arginine preceding the α helix and by the second, third, and sixth residues of the α helix. Variants of the zinc finger motif include poorly defined cysteine-rich motifs which bind zinc or other metal ions. These motifs may not contain histidine residues and are generally nonrepetitive.

The leucine zipper motif comprises a stretch of amino acids rich in leucine which can form an amphipathic α helix. This structure provides the basis for dimerization of two leucine zipper proteins. The region adjacent to the leucine zipper is usually basic, and upon protein dimerization, is optimally positioned for binding to the major groove. Proteins containing such motifs are generally referred to as bZIP transcription factors.

The helix-loop-helix motif (HLH) consists of a short α helix connected by a loop to a longer α helix. The loop is flexible and allows the two helices to fold back against each other and to bind to DNA. The transcription factor Myc contains a prototypical HLH motif.

Most transcription factors contain characteristic DNA binding motifs, and variations on the above motifs and new motifs have been and are currently being characterized. (Faisst, S. and S. Meyer (1992) Nucl. Acids Res. 20:3-26.)

Mutations in transcription factors contribute to oncogenesis. This is likely due to the role of transcription factors in the expression of genes involved in cell proliferation. For example, mutations in transcription factors encoded by proto-oncogenes, such as Fos, Jun, Myc, Rel, and Sp1, may be oncogenic due to increased stimulation of cell proliferation. Conversely, mutations in transcription factors encoded by tumor suppressor genes, such as p53, RB1, and WT1, may be oncogenic due to decreased inhibition of cell proliferation. (Latchman, D. (1995) Gene Regulation: A Eukaryotic Perspective, Chapman and Hall, London, UK, pp 242-255.)

Gene expression is also affected by chromatin-associated proteins. In the nucleus, DNA is

packaged into chromatin, the compact organization of which limits the accessibility of DNA to transcription factors and plays a key role in gene regulation. (Lewin, supra, pp. 409-410.) The compact structure of chromatin is determined and influenced by chromatin-associated proteins such as histones, high mobility group (HMG) proteins, helicases, and chromodomain proteins. There are five
5 classes of histones, H1, H2A, H2B, H3, and H4, all of which are highly basic, low molecular weight proteins. The fundamental unit of chromatin, the nucleosome, consists of 200 base pairs of DNA associated with two copies each of H2A, H2B, H3, and H4. H1 links adjacent nucleosomes. HMG proteins are low molecular weight, non-histone proteins that may play a role in unwinding DNA and stabilizing single-stranded DNA. Helicases, which are DNA-dependent ATPases, unwind DNA,
10 allowing access for transcription factors. Chromodomain proteins play a key role in the formation of highly-compacted, transcriptionally silent heterochromatin.

Much of the regulation of gene expression in eucaryotic cells occurs at the posttranscriptional level. Messenger RNAs (mRNA), which are produced in the cell nucleus from primary transcripts of protein-encoding genes, are processed and transported to the cytoplasm where the protein synthesis
15 machinery is located. RNA-binding proteins are a group of proteins that participate in the processing, editing, transport, localization, and posttranscriptional regulation of mRNAs, and comprise the protein component of ribosomes as well. The RNA-binding activity of many of these proteins is mediated by a series of RNA-binding motifs identified within them. These domains include the RNP motif, the arginine-rich motif, the RGG box, and the KH motif. (Reviewed in Burd, C. G. and Dreyfuss, G.
20 (1994) *Science* 265:615 - 621.) The RNP motif is the most widely found and best characterized of these motifs. The RNP motif is composed of 90-100 amino acids which form an RNA-binding domain and is found in one or more copies in proteins that bind pre-mRNA, mRNA, pre-ribosomal RNA, and small nuclear RNAs. The RNP motif is composed of two short sequences (RNP-1 and RNP-2) and a number of other mostly hydrophobic, conserved amino acids interspersed throughout
25 the motif. (Burd, supra; ExPASy PROSITE document PDOC0030.)

Many neoplastic disorders in humans can be attributed to inappropriate gene expression. Malignant cell growth may result from either excessive expression of tumor promoting genes or insufficient expression of tumor suppressor genes. (Cleary, M.L. (1992) *Cancer Surv.* 15:89-104.) Chromosomal translocations may also produce chimeric loci which fuse the coding sequence of one
30 gene with the regulatory regions of a second unrelated gene. Such an arrangement often results in inappropriate gene transcription. The Wilms tumor suppressor gene product, WT1, is a protein containing a DNA-binding domain consisting of four zinc fingers and a proline-glutamine rich region capable of regulating transcription. (ExPASy PROSITE document PR00049.) Deletions of the WT1 gene, or point mutations which destroy the DNA-binding activity of the protein are associated with
35 development of the pediatric nephroblastoma, Wilms tumor, and Denys-Drash syndrome. (Rauscher,

F.J. (1993) FASEB J. 7:896-903.)

Certain proteins enriched in glutamine are associated with various neurological disorders including spinocerebellar ataxia, bipolar effective disorder, schizophrenia, and autism. (Margolis, R.L. et al. (1997) Human Genetics 100:114-122.) These proteins contain regions with as many as 15 or more consecutive glutamine residues and may function as transcription factors with a potential role in regulation of neurodevelopment or neuroplasticity.

The immune system responds to infection or trauma by activating a cascade of events that coordinate the progressive selection, amplification, and mobilization of cellular defense mechanisms. A complex and balanced program of gene activation and repression is involved in this process.

However, hyperactivity of the immune system as a result of improper or insufficient regulation of gene expression may result in considerable tissue or organ damage. This damage is well documented in immunological responses associated with arthritis, allergens, heart attack, stroke, and infections. (Harrison's Principles of Internal Medicine, 13/e, McGraw Hill, Inc. and Teton Data Systems Software, 1996.) In particular, a zinc finger protein termed Staf50 (for Stimulated trans-acting factor of 50 kDa) is a transcriptional regulator and is induced in various cell lines by interferon-I and -II. Staf50 appears to mediate the antiviral activity of interferon by down-regulating the viral transcription directed by the long terminal repeat promoter region of human immunodeficiency virus type-1 in transfected cells. (Tissot, C. (1995) J. Biol. Chem. 270:14891-14898.)

Furthermore, the generation of multicellular organisms is based upon the induction and coordination of cell differentiation at the appropriate stages of development. Central to this process is differential gene expression, which confers the distinct identities of cells and tissues throughout the body. Failure to regulate gene expression during development could result in developmental disorders.

The discovery of new nucleic-acid binding proteins and the polynucleotides encoding them satisfies a need in the art by providing new compositions which are useful in the diagnosis, prevention, and treatment of reproductive, immune, and neurological disorders, and cell proliferative disorders including cancer.

SUMMARY OF THE INVENTION

The invention features purified polypeptides, protnames, referred to collectively as "ABBR" and individually as "NuABP-1," "NuABP-2," "NuABP-3," "NuABP-4," "NuABP-5," "NuABP-6," "NuABP-7," "NuABP-8," "NuABP-9," "NuABP-10," "NuABP-11," "NuABP-12," "NuABP-13," "NuABP-14," "NuABP-15," "NuABP-16," "NuABP-17," "NuABP-18," "NuABP-19," "NuABP-20," "NuABP-21," "NuABP-22," "NuABP-23," "NuABP-24," "NuABP-25," "NuABP-26," "NuABP-27," "NuABP-28," "NuABP-29," "NuABP-30," "NuABP-31," "NuABP-32," "NuABP-33,"

“NuABP-34,” “NuABP-35,” “NuABP-36,” “NuABP-37,” “NuABP-38,” “NuABP-39,” “NuABP-40” “NuABP-41,” “NuABP-42,” “NuABP-43,” “NuABP-44,” “NuABP-45,” “NuABP-46,” “NuABP-47,” “NuABP-48,” “NuABP-49,” “NuABP-50” “NuABP-51,” “NuABP-52,” “NuABP-53,” “NuABP-54,” and “NuABP-55.” In one aspect, the invention provides an isolated polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. In one alternative, the invention provides an isolated polypeptide comprising the amino acid sequence of SEQ ID NO:1-55.

The invention further provides an isolated polynucleotide encoding a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. In one alternative, the polynucleotide is selected from the group consisting of SEQ ID NO:56-110.

Additionally, the invention provides a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. In one alternative, the invention provides a cell transformed with the recombinant polynucleotide. In another alternative, the invention provides a transgenic organism comprising the recombinant polynucleotide.

The invention also provides a method for producing a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. The method comprises a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said cell is transformed with a recombinant polynucleotide comprising a promoter sequence operably linked to a polynucleotide encoding the polypeptide, and b) recovering the polypeptide so expressed.

Additionally, the invention provides an isolated antibody which specifically binds to a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active
5 fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55.

The invention further provides an isolated polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110, b) a naturally occurring
10 polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110, c) a polynucleotide sequence complementary to a), or d) a polynucleotide sequence complementary to b). In one alternative, the polynucleotide comprises at least 60 contiguous nucleotides.

Additionally, the invention provides a method for detecting a target polynucleotide in a
15 sample, said target polynucleotide having a sequence of a polynucleotide comprising a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110, b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110, c) a polynucleotide sequence complementary to a), or d) a polynucleotide sequence complementary to b). The method comprises a)
20 hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and b) detecting the presence or absence of said hybridization complex, and optionally, if present, the amount thereof. In one alternative, the probe
25 comprises at least 30 contiguous nucleotides. In another alternative, the probe comprises at least 60 contiguous nucleotides.

The invention further provides a pharmaceutical composition comprising an effective amount of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an
30 amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, and a pharmaceutically acceptable excipient. The invention additionally provides a method of treating a disease or condition associated with decreased expression of functional NuABP,
35 comprising administering to a patient in need of such treatment the pharmaceutical composition.

The invention also provides a method for screening a compound for effectiveness as an agonist of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting agonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an agonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with decreased expression of functional NuABP, comprising administering to a patient in need of such treatment the pharmaceutical composition.

Additionally, the invention provides a method for screening a compound for effectiveness as an antagonist of a polypeptide comprising a) an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, b) a naturally occurring amino acid sequence having at least 90% sequence identity to an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, c) a biologically active fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55, or d) an immunogenic fragment of an amino acid sequence selected from the group consisting of SEQ ID NO:1-55. The method comprises a) exposing a sample comprising the polypeptide to a compound, and b) detecting antagonist activity in the sample. In one alternative, the invention provides a pharmaceutical composition comprising an antagonist compound identified by the method and a pharmaceutically acceptable excipient. In another alternative, the invention provides a method of treating a disease or condition associated with overexpression of functional NuABP, comprising administering to a patient in need of such treatment the pharmaceutical composition.

The invention further provides a method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence selected from the group consisting of SEQ ID NO:56-110, the method comprising a) exposing a sample comprising the target polynucleotide to a compound, and b) detecting altered expression of the target polynucleotide.

BRIEF DESCRIPTION OF THE TABLES

Table 1 shows polypeptide and nucleotide sequence identification numbers (SEQ ID NOs), clone identification numbers (clone IDs), cDNA libraries, and cDNA fragments used to assemble full-length sequences encoding NuABP.

Table 2 shows features of each polypeptide sequence, including potential motifs, homologous sequences, and methods, algorithms, and searchable databases used for analysis of NuABP.

Table 3 shows selected fragments of each nucleic acid sequence; the tissue-specific expression patterns of each nucleic acid sequence as determined by northern analysis; diseases,
5 disorders, or conditions associated with these tissues; and the vector into which each cDNA was cloned.

Table 4 describes the tissues used to construct the cDNA libraries from which cDNA clones encoding NuABP were isolated.

Table 5 shows the tools, programs, and algorithms used to analyze NuABP, along with
10 applicable descriptions, references, and threshold parameters.

DESCRIPTION OF THE INVENTION

Before the present proteins, nucleotide sequences, and methods are described, it is understood that this invention is not limited to the particular machines, materials and methods described, as these
15 may vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention which will be limited only by the appended claims.

It must be noted that as used herein and in the appended claims, the singular forms "a," "an," and "the" include plural reference unless the context clearly dictates otherwise. Thus, for example, a
20 reference to "a host cell" includes a plurality of such host cells, and a reference to "an antibody" is a reference to one or more antibodies and equivalents thereof known to those skilled in the art, and so forth.

Unless defined otherwise, all technical and scientific terms used herein have the same meanings as commonly understood by one of ordinary skill in the art to which this invention belongs.
25 Although any machines, materials, and methods similar or equivalent to those described herein can be used to practice or test the present invention, the preferred machines, materials and methods are now described. All publications mentioned herein are cited for the purpose of describing and disclosing the cell lines, protocols, reagents and vectors which are reported in the publications and which might be used in connection with the invention. Nothing herein is to be construed as an admission that the
30 invention is not entitled to antedate such disclosure by virtue of prior invention.

DEFINITIONS

"NuABP" refers to the amino acid sequences of substantially purified NuABP obtained from any species, particularly a mammalian species, including bovine, ovine, porcine, murine, equine, and human, and from any source, whether natural, synthetic, semi-synthetic, or recombinant.

35 The term "agonist" refers to a molecule which intensifies or mimics the biological activity of

NuABP. Agonists may include proteins, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of NuABP either by directly interacting with NuABP or by acting on components of the biological pathway in which NuABP participates.

An "allelic variant" is an alternative form of the gene encoding NuABP. Allelic variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. A gene may have none, one, or many allelic variants of its naturally occurring form. Common mutational changes which give rise to allelic variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

"Altered" nucleic acid sequences encoding NuABP include those sequences with deletions, insertions, or substitutions of different nucleotides, resulting in a polypeptide the same as NuABP or a polypeptide with at least one functional characteristic of NuABP. Included within this definition are polymorphisms which may or may not be readily detectable using a particular oligonucleotide probe of the polynucleotide encoding NuABP, and improper or unexpected hybridization to allelic variants, with a locus other than the normal chromosomal locus for the polynucleotide sequence encoding NuABP. The encoded protein may also be "altered," and may contain deletions, insertions, or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent NuABP. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity, and/or the amphipathic nature of the residues, as long as the biological or immunological activity of NuABP is retained. For example, negatively charged amino acids may include aspartic acid and glutamic acid, and positively charged amino acids may include lysine and arginine. Amino acids with uncharged polar side chains having similar hydrophilicity values may include: asparagine and glutamine; and serine and threonine. Amino acids with uncharged side chains having similar hydrophilicity values may include: leucine, isoleucine, and valine; glycine and alanine; and phenylalanine and tyrosine.

The terms "amino acid" and "amino acid sequence" refer to an oligopeptide, peptide, polypeptide, or protein sequence, or a fragment of any of these, and to naturally occurring or synthetic molecules. Where "amino acid sequence" is recited to refer to an amino acid sequence of a naturally occurring protein molecule, "amino acid sequence" and like terms are not meant to limit the amino acid sequence to the complete native amino acid sequence associated with the recited protein molecule.

"Amplification" relates to the production of additional copies of a nucleic acid sequence. Amplification is generally carried out using polymerase chain reaction (PCR) technologies well known in the art.

The term "antagonist" refers to a molecule which inhibits or attenuates the biological activity of NuABP. Antagonists may include proteins such as antibodies, nucleic acids, carbohydrates, small molecules, or any other compound or composition which modulates the activity of NuABP either by directly interacting with NuABP or by acting on components of the biological pathway in which NuABP participates.

The term "antibody" refers to intact immunoglobulin molecules as well as to fragments thereof, such as Fab, F(ab')₂, and Fv fragments, which are capable of binding an epitopic determinant. Antibodies that bind NuABP polypeptides can be prepared using intact polypeptides or using fragments containing small peptides of interest as the immunizing antigen. The polypeptide or oligopeptide used to immunize an animal (e.g., a mouse, a rat, or a rabbit) can be derived from the translation of RNA, or synthesized chemically, and can be conjugated to a carrier protein if desired. Commonly used carriers that are chemically coupled to peptides include bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin (KLH). The coupled peptide is then used to immunize the animal.

The term "antigenic determinant" refers to that region of a molecule (i.e., an epitope) that makes contact with a particular antibody. When a protein or a fragment of a protein is used to immunize a host animal, numerous regions of the protein may induce the production of antibodies which bind specifically to antigenic determinants (particular regions or three-dimensional structures on the protein). An antigenic determinant may compete with the intact antigen (i.e., the immunogen used to elicit the immune response) for binding to an antibody.

The term "antisense" refers to any composition containing a nucleic acid sequence which is complementary to the "sense" strand of a specific nucleic acid sequence. Antisense molecules may be produced by any method including synthesis or transcription. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form duplexes and to block either transcription or translation. The designation "negative" or "minus" can refer to the antisense strand, and the designation "positive" or "plus" can refer to the sense strand.

The term "biologically active" refers to a protein having structural, regulatory, or biochemical functions of a naturally occurring molecule. Likewise, "immunologically active" refers to the capability of the natural, recombinant, or synthetic NuABP, or of any oligopeptide thereof, to induce a specific immune response in appropriate animals or cells and to bind with specific antibodies.

The terms "complementary" and "complementarity" refer to the natural binding of polynucleotides by base pairing. For example, the sequence "5' A-G-T 3'" bonds to the complementary sequence "3' T-C-A 5'." Complementarity between two single-stranded molecules may be "partial," such that only some of the nucleic acids bind, or it may be "complete," such that total complementarity exists between the single stranded molecules. The degree of complementarity

between nucleic acid strands has significant effects on the efficiency and strength of the hybridization between the nucleic acid strands. This is of particular importance in amplification reactions, which depend upon binding between nucleic acid strands, and in the design and use of peptide nucleic acid (PNA) molecules.

A "composition comprising a given polynucleotide sequence" and a "composition comprising a given amino acid sequence" refer broadly to any composition containing the given polynucleotide or amino acid sequence. The composition may comprise a dry formulation or an aqueous solution. Compositions comprising polynucleotide sequences encoding NuABP or fragments of NuABP may be employed as hybridization probes. The probes may be stored in freeze-dried form and may be associated with a stabilizing agent such as a carbohydrate. In hybridizations, the probe may be deployed in an aqueous solution containing salts (e.g., NaCl), detergents (e.g., sodium dodecyl sulfate; SDS), and other components (e.g., Denhardt's solution, dry milk, salmon sperm DNA, etc.).

"Consensus sequence" refers to a nucleic acid sequence which has been resequenced to resolve uncalled bases, extended using the XL-PCR kit (Perkin-Elmer, Norwalk CT) in the 5' and/or the 3' direction, and resequenced, or which has been assembled from the overlapping sequences of one or more Incyte Clones and, in some cases, one or more public domain ESTs, using a computer program for fragment assembly, such as the GELVIEW fragment assembly system (GCG, Madison WI). Some sequences have been both extended and assembled to produce the consensus sequence.

"Conservative amino acid substitutions" are those substitutions that, when made, least interfere with the properties of the original protein, i.e., the structure and especially the function of the protein is conserved and not significantly changed by such substitutions. The table below shows amino acids which may be substituted for an original amino acid in a protein and which are regarded as conservative amino acid substitutions.

	Original Residue	Conservative Substitution
25	Ala	Gly, Ser
	Arg	His, Lys
	Asn	Asp, Gln, His
	Asp	Asn, Glu
	Cys	Ala, Ser
30	Gln	Asn, Glu, His
	Glu	Asp, Gln, His
	Gly	Ala
	His	Asn, Arg, Gln, Glu
	Ile	Leu, Val
35	Leu	Ile, Val
	Lys	Arg, Gln, Glu
	Met	Leu, Ile
	Phe	His, Met, Leu, Trp, Tyr
	Ser	Cys, Thr
40	Thr	Ser, Val

Trp	Phe, Tyr
Tyr	His, Phe, Trp
Val	Ile, Leu, Thr

5 Conservative amino acid substitutions generally maintain (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a beta sheet or alpha helical conformation, (b) the charge or hydrophobicity of the molecule at the site of the substitution, and/or (c) the bulk of the side chain.

10 A "deletion" refers to a change in the amino acid or nucleotide sequence that results in the absence of one or more amino acid residues or nucleotides.

 The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, hydroxyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function
15 of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

 A "fragment" is a unique portion of NuABP or the polynucleotide encoding NuABP which is identical in sequence to but shorter in length than the parent sequence. A fragment may comprise up
20 to the entire length of the defined sequence, minus one nucleotide/amino acid residue. For example, a fragment may comprise from 5 to 1000 contiguous nucleotides or amino acid residues. A fragment used as a probe, primer, antigen, therapeutic molecule, or for other purposes, may be at least 5, 10, 15, 20, 25, 30, 40, 50, 60, 75, 100, 150, 250 or at least 500 contiguous nucleotides or amino acid residues in length. Fragments may be preferentially selected from certain regions of a molecule. For example,
25 a polypeptide fragment may comprise a certain length of contiguous amino acids selected from the first 250 or 500 amino acids (or first 25% or 50% of a polypeptide) as shown in a certain defined sequence. Clearly these lengths are exemplary, and any length that is supported by the specification, including the Sequence Listing, tables, and figures, may be encompassed by the present embodiments.

 A fragment of SEQ ID NO:56-110 comprises a region of unique polynucleotide sequence that
30 specifically identifies SEQ ID NO:56-110, for example, as distinct from any other sequence in the same genome. A fragment of SEQ ID NO:56-110 is useful, for example, in hybridization and amplification technologies and in analogous methods that distinguish SEQ ID NO:56-110 from related polynucleotide sequences. The precise length of a fragment of SEQ ID NO:56-110 and the region of SEQ ID NO:56-110 to which the fragment corresponds are routinely determinable by one of
35 ordinary skill in the art based on the intended purpose for the fragment.

 A fragment of SEQ ID NO:1-55 is encoded by a fragment of SEQ ID NO:56-110. A

fragment of SEQ ID NO:1-55 comprises a region of unique amino acid sequence that specifically identifies SEQ ID NO:1-55. For example, a fragment of SEQ ID NO:1-55 is useful as an immunogenic peptide for the development of antibodies that specifically recognize SEQ ID NO:1-55. The precise length of a fragment of SEQ ID NO:1-55 and the region of SEQ ID NO:1-55 to which the
5 fragment corresponds are routinely determinable by one of ordinary skill in the art based on the intended purpose for the fragment.

The term "similarity" refers to a degree of complementarity. There may be partial similarity or complete similarity. The word "identity" may substitute for the word "similarity." A partially complementary sequence that at least partially inhibits an identical sequence from hybridizing to a
10 target nucleic acid is referred to as "substantially similar." The inhibition of hybridization of the completely complementary sequence to the target sequence may be examined using a hybridization assay (Southern or northern blot, solution hybridization, and the like) under conditions of reduced stringency. A substantially similar sequence or hybridization probe will compete for and inhibit the binding of a completely similar (identical) sequence to the target sequence under conditions of
15 reduced stringency. This is not to say that conditions of reduced stringency are such that non-specific binding is permitted, as reduced stringency conditions require that the binding of two sequences to one another be a specific (i.e., a selective) interaction. The absence of non-specific binding may be tested by the use of a second target sequence which lacks even a partial degree of complementarity (e.g., less than about 30% similarity or identity). In the absence of non-specific binding, the
20 substantially similar sequence or probe will not hybridize to the second non-complementary target sequence.

The phrases "percent identity" and "% identity," as applied to polynucleotide sequences, refer to the percentage of residue matches between at least two polynucleotide sequences aligned using a standardized algorithm. Such an algorithm may insert, in a standardized and reproducible way, gaps
25 in the sequences being compared in order to optimize alignment between two sequences, and therefore achieve a more meaningful comparison of the two sequences.

Percent identity between polynucleotide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program. This program is part of the LASERGENE software package, a suite of
30 molecular biological analysis programs (DNASTAR, Madison WI). CLUSTAL V is described in Higgins, D.G. and P.M. Sharp (1989) CABIOS 5:151-153 and in Higgins, D.G. et al. (1992) CABIOS 8:189-191. For pairwise alignments of polynucleotide sequences, the default parameters are set as follows: Ktuple=2, gap penalty=5, window=4, and "diagonals saved"=4. The "weighted" residue weight table is selected as the default. Percent identity is reported by CLUSTAL V as the "percent
35 similarity" between aligned polynucleotide sequence pairs.

Alternatively, a suite of commonly used and freely available sequence comparison algorithms is provided by the National Center for Biotechnology Information (NCBI) Basic Local Alignment Search Tool (BLAST) (Altschul, S.F. et al. (1990) J. Mol. Biol. 215:403-410), which is available from several sources, including the NCBI, Bethesda, MD. and on the Internet at

5 <http://www.ncbi.nlm.nih.gov/BLAST/>. The BLAST software suite includes various sequence analysis programs including "blastn," that is used to align a known polynucleotide sequence with other polynucleotide sequences from a variety of databases. Also available is a tool called "BLAST 2 Sequences" that is used for direct pairwise comparison of two nucleotide sequences. "BLAST 2 Sequences" can be accessed and used interactively at <http://www.ncbi.nlm.nih.gov/gorf/bl2.html>. The

10 "BLAST 2 Sequences" tool can be used for both blastn and blastp (discussed below). BLAST programs are commonly used with gap and other parameters set to default settings. For example, to compare two nucleotide sequences, one may use blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

15 *Reward for match: 1*

Penalty for mismatch: -2

Open Gap: 5 and Extension Gap: 2 penalties

Gap x drop-off: 50

Expect: 10

20 *Word Size: 11*

Filter: on

Percent identity may be measured over the length of an entire defined sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined sequence, for instance, a fragment of at

25 least 20, at least 30, at least 40, at least 50, at least 70, at least 100, or at least 200 contiguous nucleotides. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures, or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

Nucleic acid sequences that do not show a high degree of identity may nevertheless encode

30 similar amino acid sequences due to the degeneracy of the genetic code. It is understood that changes in a nucleic acid sequence can be made using this degeneracy to produce multiple nucleic acid sequences that all encode substantially the same protein.

The phrases "percent identity" and "% identity," as applied to polypeptide sequences, refer to the percentage of residue matches between at least two polypeptide sequences aligned using a

35 standardized algorithm. Methods of polypeptide sequence alignment are well-known. Some

alignment methods take into account conservative amino acid substitutions. Such conservative substitutions, explained in more detail above, generally preserve the hydrophobicity and acidity at the site of substitution, thus preserving the structure (and therefore function) of the polypeptide.

Percent identity between polypeptide sequences may be determined using the default parameters of the CLUSTAL V algorithm as incorporated into the MEGALIGN version 3.12e sequence alignment program (described and referenced above). For pairwise alignments of polypeptide sequences using CLUSTAL V, the default parameters are set as follows: Ktuple=1, gap penalty=3, window=5, and "diagonals saved"=5. The PAM250 matrix is selected as the default residue weight table. As with polynucleotide alignments, the percent identity is reported by CLUSTAL V as the "percent similarity" between aligned polypeptide sequence pairs.

Alternatively the NCBI BLAST software suite may be used. For example, for a pairwise comparison of two polypeptide sequences, one may use the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) with blastp set at default parameters. Such default parameters may be, for example:

Matrix: BLOSUM62

Open Gap: 11 and Extension Gap: 1 penalties

Gap x drop-off: 50

Expect: 10

Word Size: 3

Filter: on

Percent identity may be measured over the length of an entire defined polypeptide sequence, for example, as defined by a particular SEQ ID number, or may be measured over a shorter length, for example, over the length of a fragment taken from a larger, defined polypeptide sequence, for instance, a fragment of at least 15, at least 20, at least 30, at least 40, at least 50, at least 70 or at least 150 contiguous residues. Such lengths are exemplary only, and it is understood that any fragment length supported by the sequences shown herein, in the tables, figures or Sequence Listing, may be used to describe a length over which percentage identity may be measured.

"Human artificial chromosomes" (HACs) are linear microchromosomes which may contain DNA sequences of about 6 kb to 10 Mb in size, and which contain all of the elements required for stable mitotic chromosome segregation and maintenance.

The term "humanized antibody" refers to antibody molecules in which the amino acid sequence in the non-antigen binding regions has been altered so that the antibody more closely resembles a human antibody, and still retains its original binding ability.

"Hybridization" refers to the process by which a polynucleotide strand anneals with a complementary strand through base pairing under defined hybridization conditions. Specific hybridization is an indication that two nucleic acid sequences share a high degree of identity. Specific

hybridization complexes form under permissive annealing conditions and remain hybridized after the "washing" step(s). The washing step(s) is particularly important in determining the stringency of the hybridization process, with more stringent conditions allowing less non-specific binding, i.e., binding between pairs of nucleic acid strands that are not perfectly matched. Permissive conditions for annealing of nucleic acid sequences are routinely determinable by one of ordinary skill in the art and may be consistent among hybridization experiments, whereas wash conditions may be varied among experiments to achieve the desired stringency, and therefore hybridization specificity. Permissive annealing conditions occur, for example, at 68°C in the presence of about 6 x SSC, about 1% (w/v) SDS, and about 100 µg/ml denatured salmon sperm DNA.

Generally, stringency of hybridization is expressed, in part, with reference to the temperature under which the wash step is carried out. Generally, such wash temperatures are selected to be about 5°C to 20°C lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength and pH) at which 50% of the target sequence hybridizes to a perfectly matched probe. An equation for calculating T_m and conditions for nucleic acid hybridization are well known and can be found in Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; specifically see volume 2, chapter 9.

High stringency conditions for hybridization between polynucleotides of the present invention include wash conditions of 68°C in the presence of about 0.2 x SSC and about 0.1% SDS, for 1 hour.

Alternatively, temperatures of about 65°C, 60°C, 55°C, or 42°C may be used. SSC concentration may be varied from about 0.1 to 2 x SSC, with SDS being present at about 0.1%. Typically, blocking reagents are used to block non-specific hybridization. Such blocking reagents include, for instance, denatured salmon sperm DNA at about 100-200 µg/ml. Organic solvent, such as formamide at a concentration of about 35-50% v/v, may also be used under particular circumstances, such as for RNA:DNA hybridizations. Useful variations on these wash conditions will be readily apparent to those of ordinary skill in the art. Hybridization, particularly under high stringency conditions, may be suggestive of evolutionary similarity between the nucleotides. Such similarity is strongly indicative of a similar role for the nucleotides and their encoded polypeptides.

The term "hybridization complex" refers to a complex formed between two nucleic acid sequences by virtue of the formation of hydrogen bonds between complementary bases. A hybridization complex may be formed in solution (e.g., C_0t or R_0t analysis) or formed between one nucleic acid sequence present in solution and another nucleic acid sequence immobilized on a solid support (e.g., paper, membranes, filters, chips, pins or glass slides, or any other appropriate substrate to which cells or their nucleic acids have been fixed).

The words "insertion" and "addition" refer to changes in an amino acid or nucleotide

sequence resulting in the addition of one or more amino acid residues or nucleotides, respectively.

"Immune response" can refer to conditions associated with inflammation, trauma, immune disorders, or infectious or genetic disease, etc. These conditions can be characterized by expression of various factors, e.g., cytokines, chemokines, and other signaling molecules, which may affect

5 cellular and systemic defense systems.

The term "microarray" refers to an arrangement of distinct polynucleotides on a substrate.

The terms "element" and "array element" in a microarray context, refer to hybridizable polynucleotides arranged on the surface of a substrate.

The term "modulate" refers to a change in the activity of NuABP. For example, modulation
10 may cause an increase or a decrease in protein activity, binding characteristics, or any other biological, functional, or immunological properties of NuABP.

The phrases "nucleic acid" and "nucleic acid sequence" refer to a nucleotide, oligonucleotide, polynucleotide, or any fragment thereof. These phrases also refer to DNA or RNA of genomic or synthetic origin which may be single-stranded or double-stranded and may represent the sense or the
15 antisense strand, to peptide nucleic acid (PNA), or to any DNA-like or RNA-like material.

"Operably linked" refers to the situation in which a first nucleic acid sequence is placed in a functional relationship with the second nucleic acid sequence. For instance, a promoter is operably linked to a coding sequence if the promoter affects the transcription or expression of the coding sequence. Generally, operably linked DNA sequences may be in close proximity or contiguous and,
20 where necessary to join two protein coding regions, in the same reading frame.

"Peptide nucleic acid" (PNA) refers to an antisense molecule or anti-gene agent which comprises an oligonucleotide of at least about 5 nucleotides in length linked to a peptide backbone of amino acid residues ending in lysine. The terminal lysine confers solubility to the composition. PNAs preferentially bind complementary single stranded DNA or RNA and stop transcript elongation,
25 and may be pegylated to extend their lifespan in the cell.

"Probe" refers to nucleic acid sequences encoding NuABP, their complements, or fragments thereof, which are used to detect identical, allelic or related nucleic acid sequences. Probes are isolated oligonucleotides or polynucleotides attached to a detectable label or reporter molecule. Typical labels include radioactive isotopes, ligands, chemiluminescent agents, and enzymes.

30 "Primers" are short nucleic acids, usually DNA oligonucleotides, which may be annealed to a target polynucleotide by complementary base-pairing. The primer may then be extended along the target DNA strand by a DNA polymerase enzyme. Primer pairs can be used for amplification (and identification) of a nucleic acid sequence, e.g., by the polymerase chain reaction (PCR).

Probes and primers as used in the present invention typically comprise at least 15 contiguous
35 nucleotides of a known sequence. In order to enhance specificity, longer probes and primers may also

be employed, such as probes and primers that comprise at least 20, 25, 30, 40, 50, 60, 70, 80, 90, 100, or at least 150 consecutive nucleotides of the disclosed nucleic acid sequences. Probes and primers may be considerably longer than these examples, and it is understood that any length supported by the specification, including the tables, figures, and Sequence Listing, may be used.

5 Methods for preparing and using probes and primers are described in the references, for example Sambrook et al., 1989, Molecular Cloning: A Laboratory Manual, 2nd ed., vol. 1-3, Cold Spring Harbor Press, Plainview NY; Ausubel et al., 1987, Current Protocols in Molecular Biology, Greene Publ. Assoc. & Wiley-Intersciences, New York NY; Innis et al., 1990, PCR Protocols, A Guide to Methods and Applications, Academic Press, San Diego CA. PCR primer pairs can be
10 derived from a known sequence, for example, by using computer programs intended for that purpose such as Primer (Version 0.5, 1991, Whitehead Institute for Biomedical Research, Cambridge MA).

 Oligonucleotides for use as primers are selected using software known in the art for such purpose. For example, OLIGO 4.06 software is useful for the selection of PCR primer pairs of up to 100 nucleotides each, and for the analysis of oligonucleotides and larger polynucleotides of up to
15 5,000 nucleotides from an input polynucleotide sequence of up to 32 kilobases. Similar primer selection programs have incorporated additional features for expanded capabilities. For example, the PrimOU primer selection program (available to the public from the Genome Center at University of Texas South West Medical Center, Dallas TX) is capable of choosing specific primers from megabase sequences and is thus useful for designing primers on a genome-wide scope. The Primer3 primer
20 selection program (available to the public from the Whitehead Institute/MIT Center for Genome Research, Cambridge MA) allows the user to input a "mispriming library," in which sequences to avoid as primer binding sites are user-specified. Primer3 is useful, in particular, for the selection of oligonucleotides for microarrays. (The source code for the latter two primer selection programs may also be obtained from their respective sources and modified to meet the user's specific needs.) The
25 PrimeGen program (available to the public from the UK Human Genome Mapping Project Resource Centre, Cambridge UK) designs primers based on multiple sequence alignments, thereby allowing selection of primers that hybridize to either the most conserved or least conserved regions of aligned nucleic acid sequences. Hence, this program is useful for identification of both unique and conserved oligonucleotides and polynucleotide fragments. The oligonucleotides and polynucleotide fragments
30 identified by any of the above selection methods are useful in hybridization technologies, for example, as PCR or sequencing primers, microarray elements, or specific probes to identify fully or partially complementary polynucleotides in a sample of nucleic acids. Methods of oligonucleotide selection are not limited to those described above.

 A "recombinant nucleic acid" is a sequence that is not naturally occurring or has a sequence
35 that is made by an artificial combination of two or more otherwise separated segments of sequence.

This artificial combination is often accomplished by chemical synthesis or, more commonly, by the artificial manipulation of isolated segments of nucleic acids, e.g., by genetic engineering techniques such as those described in Sambrook, supra. The term recombinant includes nucleic acids that have been altered solely by addition, substitution, or deletion of a portion of the nucleic acid. Frequently, a recombinant nucleic acid may include a nucleic acid sequence operably linked to a promoter sequence. Such a recombinant nucleic acid may be part of a vector that is used, for example, to transform a cell.

Alternatively, such recombinant nucleic acids may be part of a viral vector, e.g., based on a vaccinia virus, that could be used to vaccinate a mammal wherein the recombinant nucleic acid is expressed, inducing a protective immunological response in the mammal.

The term "sample" is used in its broadest sense. A sample suspected of containing nucleic acids encoding NuABP, or fragments thereof, or NuABP itself, may comprise a bodily fluid; an extract from a cell, chromosome, organelle, or membrane isolated from a cell; a cell; genomic DNA, RNA, or cDNA, in solution or bound to a substrate; a tissue; a tissue print; etc.

The terms "specific binding" and "specifically binding" refer to that interaction between a protein or peptide and an agonist, an antibody, an antagonist, a small molecule, or any natural or synthetic binding composition. The interaction is dependent upon the presence of a particular structure of the protein, e.g., the antigenic determinant or epitope, recognized by the binding molecule. For example, if an antibody is specific for epitope "A," the presence of a polypeptide containing the epitope A, or the presence of free unlabeled A, in a reaction containing free labeled A and the antibody will reduce the amount of labeled A that binds to the antibody.

The term "substantially purified" refers to nucleic acid or amino acid sequences that are removed from their natural environment and are isolated or separated, and are at least 60% free, preferably at least 75% free, and most preferably at least 90% free from other components with which they are naturally associated.

A "substitution" refers to the replacement of one or more amino acids or nucleotides by different amino acids or nucleotides, respectively.

"Substrate" refers to any suitable rigid or semi-rigid support including membranes, filters, chips, slides, wafers, fibers, magnetic or nonmagnetic beads, gels, tubing, plates, polymers, microparticles and capillaries. The substrate can have a variety of surface forms, such as wells, trenches, pins, channels and pores, to which polynucleotides or polypeptides are bound.

"Transformation" describes a process by which exogenous DNA enters and changes a recipient cell. Transformation may occur under natural or artificial conditions according to various methods well known in the art, and may rely on any known method for the insertion of foreign nucleic acid sequences into a prokaryotic or eukaryotic host cell. The method for transformation is selected

based on the type of host cell being transformed and may include, but is not limited to, viral infection, electroporation, heat shock, lipofection, and particle bombardment. The term "transformed" cells includes stably transformed cells in which the inserted DNA is capable of replication either as an autonomously replicating plasmid or as part of the host chromosome, as well as transiently

5 transformed cells which express the inserted DNA or RNA for limited periods of time.

A "variant" of a particular nucleic acid sequence is defined as a nucleic acid sequence having at least 40% sequence identity to the particular nucleic acid sequence over a certain length of one of the nucleic acid sequences using blastn with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of nucleic acids may show, for example, at least 50%, at
10 least 60%, at least 70%, at least 80%, at least 85%, at least 90%, at least 95% or at least 98% or greater sequence identity over a certain defined length. A variant may be described as, for example, an "allelic" (as defined above), "splice," "species," or "polymorphic" variant. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding
15 polypeptide may possess additional functional domains or lack domains that are present in the reference molecule. Species variants are polynucleotide sequences that vary from one species to another. The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide
20 polymorphisms" (SNPs) in which the polynucleotide sequence varies by one nucleotide base. The presence of SNPs may be indicative of, for example, a certain population, a disease state, or a propensity for a disease state.

A "variant" of a particular polypeptide sequence is defined as a polypeptide sequence having at least 40% sequence identity to the particular polypeptide sequence over a certain length of one of
25 the polypeptide sequences using blastp with the "BLAST 2 Sequences" tool Version 2.0.9 (May-07-1999) set at default parameters. Such a pair of polypeptides may show, for example, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or at least 98% or greater sequence identity over a certain defined length of one of the polypeptides.

30 THE INVENTION

The invention is based on the discovery of new human nucleic-acid binding proteins (NuABP), the polynucleotides encoding NuABP, and the use of these compositions for the diagnosis, treatment, or prevention of reproductive, immune, and neurological disorders, and cell proliferative disorders including cancer.

35 Table 1 lists the Incyte clones used to assemble full length nucleotide sequences encoding

NuABP. Columns 1 and 2 show the sequence identification numbers (SEQ ID NOs) of the polypeptide and nucleotide sequences, respectively. Column 3 shows the clone IDs of the Incyte clones in which nucleic acids encoding each NuABP were identified, and column 4 shows the cDNA libraries from which these clones were isolated. Column 5 shows Incyte clones and their
 5 corresponding cDNA libraries. Clones for which cDNA libraries are not indicated were derived from pooled cDNA libraries. The Incyte clones in column 5 were used to assemble the consensus nucleotide sequence of each NuABP and are useful as fragments in hybridization technologies.

The columns of Table 2 show various properties of each of the polypeptides of the invention: column 1 references the SEQ ID NO; column 2 shows the number of amino acid residues in each
 10 polypeptide; column 3 shows potential phosphorylation sites; column 4 shows potential glycosylation sites; column 5 shows the amino acid residues comprising signature sequences and motifs; column 6 shows identification or homologous sequences as identified by BLAST analysis; and column 7 shows analytical methods and in some cases, searchable databases to which the analytical methods were applied. The methods of column 7 were used to characterize each polypeptide through sequence
 15 homology and protein motifs.

The columns of Table 3 show the tissue-specificity and diseases, disorders, or conditions associated with nucleotide sequences encoding NuABP. The first column of Table 3 lists the nucleotide SEQ ID NOs. Column 2 lists fragments of the nucleotide sequences of column 1. These fragments are useful, for example, in hybridization or amplification technologies to identify SEQ ID
 20 NO:56-110 and to distinguish between SEQ ID NO:56-110 and related polynucleotide sequences. The polypeptides encoded by these fragments are useful, for example, as immunogenic peptides. Column 3 lists tissue categories which express NuABP as a fraction of total tissues expressing NuABP. Column 4 lists diseases, disorders, or conditions associated with those tissues expressing NuABP as a fraction of total tissues expressing NuABP. Of particular note is the expression of SEQ
 25 ID NO:83 and SEQ ID NO:110 in neurological tissue. About 53% of the cDNA libraries expressing SEQ ID NO:83 are derived from neurological tissue. Furthermore, SEQ ID NO:110 expression is detected exclusively in a cDNA library derived from brain tissue afflicted with Huntington's disease. Column 5 lists the vectors used to subclone each cDNA library.

The columns of Table 4 show descriptions of the tissues used to construct the cDNA libraries
 30 from which cDNA clones encoding NuABP were isolated. Column 1 references the nucleotide SEQ ID NOs, column 2 shows the cDNA libraries from which these clones were isolated, and column 3 shows the tissue origins and other descriptive information relevant to the cDNA libraries in column 2.

Fragments of the nucleotide sequences encoding NuABP are useful, for example, in hybridization or amplification technologies to identify SEQ ID NOS:56-110 and to distinguish
 35 between SEQ ID NOS:56-110 and related polynucleotide sequences. The polypeptides encoded by

these fragments are useful, for example, as immunogenic peptides.

The invention also encompasses NuABP variants. A preferred NuABP variant is one which has at least about 80%, or alternatively at least about 90%, or even at least about 95% amino acid sequence identity to the NuABP amino acid sequence, and which contains at least one functional or structural characteristic of NuABP.

The invention also encompasses polynucleotides which encode NuABP. In a particular embodiment, the invention encompasses a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:56-110, which encodes NuABP.

The invention also encompasses a variant of a polynucleotide sequence encoding NuABP. In particular, such a variant polynucleotide sequence will have at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to the polynucleotide sequence encoding NuABP. A particular aspect of the invention encompasses a variant of a polynucleotide sequence comprising a sequence selected from the group consisting of SEQ ID NO:56-110 which has at least about 70%, or alternatively at least about 85%, or even at least about 95% polynucleotide sequence identity to a nucleic acid sequence selected from the group consisting of SEQ ID NO:56-110. Any one of the polynucleotide variants described above can encode an amino acid sequence which contains at least one functional or structural characteristic of NuABP.

It will be appreciated by those skilled in the art that as a result of the degeneracy of the genetic code, a multitude of polynucleotide sequences encoding NuABP, some bearing minimal similarity to the polynucleotide sequences of any known and naturally occurring gene, may be produced. Thus, the invention contemplates each and every possible variation of polynucleotide sequence that could be made by selecting combinations based on possible codon choices. These combinations are made in accordance with the standard triplet genetic code as applied to the polynucleotide sequence of naturally occurring NuABP, and all such variations are to be considered as being specifically disclosed.

Although nucleotide sequences which encode NuABP and its variants are generally capable of hybridizing to the nucleotide sequence of the naturally occurring NuABP under appropriately selected conditions of stringency, it may be advantageous to produce nucleotide sequences encoding NuABP or its derivatives possessing a substantially different codon usage, e.g., inclusion of non-naturally occurring codons. Codons may be selected to increase the rate at which expression of the peptide occurs in a particular prokaryotic or eukaryotic host in accordance with the frequency with which particular codons are utilized by the host. Other reasons for substantially altering the nucleotide sequence encoding NuABP and its derivatives without altering the encoded amino acid sequences include the production of RNA transcripts having more desirable properties, such as a greater half-life, than transcripts produced from the naturally occurring sequence.

The invention also encompasses production of DNA sequences which encode NuABP and NuABP derivatives, or fragments thereof, entirely by synthetic chemistry. After production, the synthetic sequence may be inserted into any of the many available expression vectors and cell systems using reagents well known in the art. Moreover, synthetic chemistry may be used to introduce mutations into a sequence encoding NuABP or any fragment thereof.

Also encompassed by the invention are polynucleotide sequences that are capable of hybridizing to the claimed polynucleotide sequences, and, in particular, to those shown in SEQ ID NO:56-110 and fragments thereof under various conditions of stringency. (See, e.g., Wahl, G.M. and S.L. Berger (1987) *Methods Enzymol.* 152:399-407; Kimmel, A.R. (1987) *Methods Enzymol.* 152:507-511.) Hybridization conditions, including annealing and wash conditions, are described in "Definitions."

Methods for DNA sequencing are well known in the art and may be used to practice any of the embodiments of the invention. The methods may employ such enzymes as the Klenow fragment of DNA polymerase I, SEQUENASE (US Biochemical, Cleveland OH), Taq polymerase (Perkin-Elmer), thermostable T7 polymerase (Amersham Pharmacia Biotech, Piscataway NJ), or combinations of polymerases and proofreading exonucleases such as those found in the ELONGASE amplification system (Life Technologies, Gaithersburg MD). Preferably, sequence preparation is automated with machines such as the MICROLAB 2200 liquid transfer system (Hamilton, Reno NV), PTC200 thermal cycler (MJ Research, Watertown MA) and ABI CATALYST 800 thermal cycler (Perkin-Elmer). Sequencing is then carried out using either the ABI 373 or 377 DNA sequencing system (Perkin-Elmer), the MEGABACE 1000 DNA sequencing system (Molecular Dynamics, Sunnyvale CA), or other systems known in the art. The resulting sequences are analyzed using a variety of algorithms which are well known in the art. (See, e.g., Ausubel, F.M. (1997) Short Protocols in Molecular Biology, John Wiley & Sons, New York NY, unit 7.7; Meyers, R.A. (1995) Molecular Biology and Biotechnology, Wiley VCH, New York NY, pp. 856-853.)

The nucleic acid sequences encoding NuABP may be extended utilizing a partial nucleotide sequence and employing various PCR-based methods known in the art to detect upstream sequences, such as promoters and regulatory elements. For example, one method which may be employed, restriction-site PCR, uses universal and nested primers to amplify unknown sequence from genomic DNA within a cloning vector. (See, e.g., Sarkar, G. (1993) *PCR Methods Applic.* 2:318-322.) Another method, inverse PCR, uses primers that extend in divergent directions to amplify unknown sequence from a circularized template. The template is derived from restriction fragments comprising a known genomic locus and surrounding sequences. (See, e.g., Triglia, T. et al. (1988) *Nucleic Acids Res.* 16:8186.) A third method, capture PCR, involves PCR amplification of DNA fragments adjacent to known sequences in human and yeast artificial chromosome DNA. (See, e.g., Lagerstrom, M. et al.

(1991) PCR Methods Applic. 1:111-119.) In this method, multiple restriction enzyme digestions and ligations may be used to insert an engineered double-stranded sequence into a region of unknown sequence before performing PCR. Other methods which may be used to retrieve unknown sequences are known in the art. (See, e.g., Parker, J.D. et al. (1991) Nucleic Acids Res. 19:3055-3060).

5 Additionally, one may use PCR, nested primers, and PROMOTERFINDER libraries (Clontech, Palo Alto CA) to walk genomic DNA. This procedure avoids the need to screen libraries and is useful in finding intron/exon junctions. For all PCR-based methods, primers may be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences, Plymouth MN) or another appropriate program, to be about 22 to 30 nucleotides in
10 length, to have a GC content of about 50% or more, and to anneal to the template at temperatures of about 68°C to 72°C.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. In addition, random-primed libraries, which often include sequences containing the 5' regions of genes, are preferable for situations in which an oligo d(T)
15 library does not yield a full-length cDNA. Genomic libraries may be useful for extension of sequence into 5' non-transcribed regulatory regions.

Capillary electrophoresis systems which are commercially available may be used to analyze the size or confirm the nucleotide sequence of sequencing or PCR products. In particular, capillary sequencing may employ flowable polymers for electrophoretic separation, four different nucleotide-
20 specific, laser-stimulated fluorescent dyes, and a charge coupled device camera for detection of the emitted wavelengths. Output/light intensity may be converted to electrical signal using appropriate software (e.g., GENOTYPER and SEQUENCE NAVIGATOR, Perkin-Elmer), and the entire process from loading of samples to computer analysis and electronic data display may be computer controlled. Capillary electrophoresis is especially preferable for sequencing small DNA fragments which may be
25 present in limited amounts in a particular sample.

In another embodiment of the invention, polynucleotide sequences or fragments thereof which encode NuABP may be cloned in recombinant DNA molecules that direct expression of NuABP, or fragments or functional equivalents thereof, in appropriate host cells. Due to the inherent degeneracy of the genetic code, other DNA sequences which encode substantially the same or a
30 functionally equivalent amino acid sequence may be produced and used to express NuABP.

The nucleotide sequences of the present invention can be engineered using methods generally known in the art in order to alter NuABP-encoding sequences for a variety of purposes including, but not limited to, modification of the cloning, processing, and/or expression of the gene product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic
35 oligonucleotides may be used to engineer the nucleotide sequences. For example, oligonucleotide-

mediated site-directed mutagenesis may be used to introduce mutations that create new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, and so forth.

In another embodiment, sequences encoding NuABP may be synthesized, in whole or in part, using chemical methods well known in the art. (See, e.g., Caruthers, M.H. et al. (1980) *Nucleic Acids Symp. Ser.* 7:215-223; and Horn, T. et al. (1980) *Nucleic Acids Symp. Ser.* 7:225-232.)

Alternatively, NuABP itself or a fragment thereof may be synthesized using chemical methods. For example, peptide synthesis can be performed using various solid-phase techniques. (See, e.g., Roberge, J.Y. et al. (1995) *Science* 269:202-204.) Automated synthesis may be achieved using the ABI 431A peptide synthesizer (Perkin-Elmer). Additionally, the amino acid sequence of NuABP, or any part thereof, may be altered during direct synthesis and/or combined with sequences from other proteins, or any part thereof, to produce a variant polypeptide.

The peptide may be substantially purified by preparative high performance liquid chromatography. (See, e.g., Chiez, R.M. and F.Z. Regnier (1990) *Methods Enzymol.* 182:392-421.) The composition of the synthetic peptides may be confirmed by amino acid analysis or by sequencing. (See, e.g., Creighton, T. (1984) Proteins, Structures and Molecular Properties, WH Freeman, New York NY.)

In order to express a biologically active NuABP, the nucleotide sequences encoding NuABP or derivatives thereof may be inserted into an appropriate expression vector, i.e., a vector which contains the necessary elements for transcriptional and translational control of the inserted coding sequence in a suitable host. These elements include regulatory sequences, such as enhancers, constitutive and inducible promoters, and 5' and 3' untranslated regions in the vector and in polynucleotide sequences encoding NuABP. Such elements may vary in their strength and specificity. Specific initiation signals may also be used to achieve more efficient translation of sequences encoding NuABP. Such signals include the ATG initiation codon and adjacent sequences, e.g. the Kozak sequence. In cases where sequences encoding NuABP and its initiation codon and upstream regulatory sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals including an in-frame ATG initiation codon should be provided by the vector. Exogenous translational elements and initiation codons may be of various origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of enhancers appropriate for the particular host cell system used. (See, e.g., Scharf, D. et al. (1994) *Results Probl. Cell Differ.* 20:125-162.)

Methods which are well known to those skilled in the art may be used to construct expression vectors containing sequences encoding NuABP and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques,

and in vivo genetic recombination. (See, e.g., Sambrook, J. et al. (1989) Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Plainview NY, ch. 4, 8, and 16-17; Ausubel, F.M. et al. (1995) Current Protocols in Molecular Biology, John Wiley & Sons, New York NY, ch. 9, 13, and 16.)

5 A variety of expression vector/host systems may be utilized to contain and express sequences encoding NuABP. These include, but are not limited to, microorganisms such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors; insect cell systems infected with viral expression vectors (e.g., baculovirus); plant cell systems transformed with viral expression vectors (e.g., cauliflower mosaic virus, CaMV, or tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids); or
10 animal cell systems. The invention is not limited by the host cell employed.

In bacterial systems, a number of cloning and expression vectors may be selected depending upon the use intended for polynucleotide sequences encoding NuABP. For example, routine cloning, subcloning, and propagation of polynucleotide sequences encoding NuABP can be achieved using a
15 multifunctional E. coli vector such as PBLUESCRIPT (Stratagene, La Jolla CA) or PSPORT1 plasmid (Life Technologies). Ligation of sequences encoding NuABP into the vector's multiple cloning site disrupts the *lacZ* gene, allowing a colorimetric screening procedure for identification of transformed bacteria containing recombinant molecules. In addition, these vectors may be useful for
20 in vitro transcription, dideoxy sequencing, single strand rescue with helper phage, and creation of nested deletions in the cloned sequence. (See, e.g., Van Heeke, G. and S.M. Schuster (1989) J. Biol. Chem. 264:5503-5509.) When large quantities of NuABP are needed, e.g. for the production of antibodies, vectors which direct high level expression of NuABP may be used. For example, vectors containing the strong, inducible T5 or T7 bacteriophage promoter may be used.

Yeast expression systems may be used for production of NuABP. A number of vectors
25 containing constitutive or inducible promoters, such as alpha factor, alcohol oxidase, and PGH promoters, may be used in the yeast Saccharomyces cerevisiae or Pichia pastoris. In addition, such vectors direct either the secretion or intracellular retention of expressed proteins and enable integration of foreign sequences into the host genome for stable propagation. (See, e.g., Ausubel, 1995, supra; Bitter, G.A. et al. (1987) Methods Enzymol. 153:516-544; and Scorer, C.A. et al. (1994)
30 Bio/Technology 12:181-184.)

Plant systems may also be used for expression of NuABP. Transcription of sequences encoding NuABP may be driven viral promoters, e.g., the 35S and 19S promoters of CaMV used alone or in combination with the omega leader sequence from TMV (Takamatsu, N. (1987) EMBO J. 6:307-311). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock
35 promoters may be used. (See, e.g., Coruzzi, G. et al. (1984) EMBO J. 3:1671-1680; Broglie, R. et al.

(1984) Science 224:838-843; and Winter, J. et al. (1991) Results Probl. Cell Differ. 17:85-105.)

These constructs can be introduced into plant cells by direct DNA transformation or pathogen-mediated transfection. (See, e.g., The McGraw Hill Yearbook of Science and Technology (1992) McGraw Hill, New York NY, pp. 191-196.)

5 In mammalian cells, a number of viral-based expression systems may be utilized. In cases where an adenovirus is used as an expression vector, sequences encoding NuABP may be ligated into an adenovirus transcription/translation complex consisting of the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome may be used to obtain infective virus which expresses NuABP in host cells. (See, e.g., Logan, J. and T. Shenk (1984) Proc. Natl. Acad. Sci. USA 81:3655-3659.) In addition, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, may be used to increase expression in mammalian host cells. SV40 or EBV-based vectors may also be used for high-level protein expression.

Human artificial chromosomes (HACs) may also be employed to deliver larger fragments of DNA than can be contained in and expressed from a plasmid. HACs of about 6 kb to 10 Mb are constructed and delivered via conventional delivery methods (liposomes, polycationic amino polymers, or vesicles) for therapeutic purposes. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355.)

For long term production of recombinant proteins in mammalian systems, stable expression of NuABP in cell lines is preferred. For example, sequences encoding NuABP can be transformed into cell lines using expression vectors which may contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells may be allowed to grow for about 1 to 2 days in enriched media before being switched to selective media. The purpose of the selectable marker is to confer resistance to a selective agent, and its presence allows growth and recovery of cells which successfully express the introduced sequences. Resistant clones of stably transformed cells may be propagated using tissue culture techniques appropriate to the cell type.

Any number of selection systems may be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase and adenine phosphoribosyltransferase genes, for use in *tk* and *apr* cells, respectively. (See, e.g., Wigler, M. et al. (1977) Cell 11:223-232; Lowy, I. et al. (1980) Cell 22:817-823.) Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, *dhfr* confers resistance to methotrexate; *neo* confers resistance to the aminoglycosides neomycin and G-418; and *als* and *pat* confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. (See, e.g., Wigler, M. et al. (1980) Proc. Natl. Acad. Sci. USA 77:3567-3570; Colbere-Garapin, F. et al. (1981) J. Mol. Biol. 150:1-14.) Additional selectable genes have been described, e.g., *trpB* and *hisD*, which

alter cellular requirements for metabolites. (See, e.g., Hartman, S.C. and R.C. Mulligan (1988) Proc. Natl. Acad. Sci. USA 85:8047-8051.) Visible markers, e.g., anthocyanins, green fluorescent proteins (GFP; Clontech), β glucuronidase and its substrate β -glucuronide, or luciferase and its substrate luciferin may be used. These markers can be used not only to identify transformants, but also to
5 quantify the amount of transient or stable protein expression attributable to a specific vector system. (See, e.g., Rhodes, C.A. (1995) Methods Mol. Biol. 55:121-131.)

Although the presence/absence of marker gene expression suggests that the gene of interest is also present, the presence and expression of the gene may need to be confirmed. For example, if the sequence encoding NuABP is inserted within a marker gene sequence, transformed cells containing
10 sequences encoding NuABP can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding NuABP under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the tandem gene as well.

In general, host cells that contain the nucleic acid sequence encoding NuABP and that express
15 NuABP may be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations, PCR amplification, and protein bioassay or immunoassay techniques which include membrane, solution, or chip based technologies for the detection and/or quantification of nucleic acid or protein sequences.

Immunological methods for detecting and measuring the expression of NuABP using either
20 specific polyclonal or monoclonal antibodies are known in the art. Examples of such techniques include enzyme-linked immunosorbent assays (ELISAs), radioimmunoassays (RIAs), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering epitopes on NuABP is preferred, but a competitive binding assay may be employed. These and other assays are well known in the art. (See,
25 e.g., Hampton, R. et al. (1990) Serological Methods, a Laboratory Manual, APS Press, St. Paul MN, Sect. IV; Coligan, J.E. et al. (1997) Current Protocols in Immunology, Greene Pub. Associates and Wiley-Interscience, New York NY; and Pound, J.D. (1998) Immunochemical Protocols, Humana Press, Totowa NJ.)

A wide variety of labels and conjugation techniques are known by those skilled in the art and
30 may be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding NuABP include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, the sequences encoding NuABP, or any fragments thereof, may be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available,
35 and may be used to synthesize RNA probes in vitro by addition of an appropriate RNA polymerase

such as T7, T3, or SP6 and labeled nucleotides. These procedures may be conducted using a variety of commercially available kits, such as those provided by Amersham Pharmacia Biotech, Promega (Madison WI), and US Biochemical. Suitable reporter molecules or labels which may be used for ease of detection include radionuclides, enzymes, fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Host cells transformed with nucleotide sequences encoding NuABP may be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The protein produced by a transformed cell may be secreted or retained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode NuABP may be designed to contain signal sequences which direct secretion of NuABP through a prokaryotic or eukaryotic cell membrane.

In addition, a host cell strain may be chosen for its ability to modulate expression of the inserted sequences or to process the expressed protein in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a "prepro" or "pro" form of the protein may also be used to specify protein targeting, folding, and/or activity. Different host cells which have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38) are available from the American Type Culture Collection (ATCC, Manassas VA) and may be chosen to ensure the correct modification and processing of the foreign protein.

In another embodiment of the invention, natural, modified, or recombinant nucleic acid sequences encoding NuABP may be ligated to a heterologous sequence resulting in translation of a fusion protein in any of the aforementioned host systems. For example, a chimeric NuABP protein containing a heterologous moiety that can be recognized by a commercially available antibody may facilitate the screening of peptide libraries for inhibitors of NuABP activity. Heterologous protein and peptide moieties may also facilitate purification of fusion proteins using commercially available affinity matrices. Such moieties include, but are not limited to, glutathione S-transferase (GST), maltose binding protein (MBP), thioredoxin (Trx), calmodulin binding peptide (CBP), 6-His, FLAG, *c-myc*, and hemagglutinin (HA). GST, MBP, Trx, CBP, and 6-His enable purification of their cognate fusion proteins on immobilized glutathione, maltose, phenylarsine oxide, calmodulin, and metal-chelate resins, respectively. FLAG, *c-myc*, and hemagglutinin (HA) enable immunoaffinity purification of fusion proteins using commercially available monoclonal and polyclonal antibodies that specifically recognize these epitope tags. A fusion protein may also be engineered to contain a proteolytic cleavage site located between the NuABP encoding sequence and the heterologous protein sequence, so that NuABP may be cleaved away from the heterologous moiety following purification.

Methods for fusion protein expression and purification are discussed in Ausubel (1995, supra, ch. 10). A variety of commercially available kits may also be used to facilitate expression and purification of fusion proteins.

In a further embodiment of the invention, synthesis of radiolabeled NuABP may be achieved
5 in vitro using the TNT rabbit reticulocyte lysate or wheat germ extract system (Promega). These systems couple transcription and translation of protein-coding sequences operably associated with the T7, T3, or SP6 promoters. Translation takes place in the presence of a radiolabeled amino acid precursor, for example, ³⁵S-methionine.

Fragments of NuABP may be produced not only by recombinant means, but also by direct
10 peptide synthesis using solid-phase techniques. (See, e.g., Creighton, supra, pp. 55-60.) Protein synthesis may be performed by manual techniques or by automation. Automated synthesis may be achieved, for example, using the ABI 431A peptide synthesizer (Perkin-Elmer). Various fragments of NuABP may be synthesized separately and then combined to produce the full length molecule.

THERAPEUTICS

15 Chemical and structural similarity, e.g., in the context of sequences and motifs, exists between regions of NuABP and nucleic-acid binding proteins. In addition, the expression of NuABP is closely associated with proliferative, neuronal, inflamed, and cancerous tissues and tissues of the reproductive system. Therefore, NuABP appears to play a role in reproductive, immune, and neurological disorders, and cell proliferative disorders including cancer. In the treatment of disorders associated
20 with increased NuABP expression or activity, it is desirable to decrease the expression or activity of NuABP. In the treatment of disorders associated with decreased NuABP expression or activity, it is desirable to increase the expression or activity of NuABP.

Therefore, in one embodiment, NuABP or a fragment or derivative thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity
25 of NuABP. Examples of such disorders include, but are not limited to, a reproductive disorder such as disorders of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, disruptions of the estrous cycle, disruptions of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, endometrial and ovarian tumors, uterine fibroids, autoimmune disorders, ectopic pregnancies, and teratogenesis; cancer of the breast, fibrocystic breast
30 disease, and galactorrhea; disruptions of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; an immune disorder such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, arteriosclerosis, asthma,
35 atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, bursitis,

cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis. Goodpasture's syndrome, gout, Graves' disease. Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel syndrome,

- 5 episodic lymphopenia with lymphocytotoxins, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, ulcerative colitis, uveitis,
- 10 Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy,
- 15 retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann-Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis,
- 20 tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases. muscular dystrophy and other neuromuscular disorders, peripheral nervous system disorders. dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies,
- 25 myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders. akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses. postherpetic neuralgia, and Tourette's disorder; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal hemoglobinuria, polycythemia vera, psoriasis,
- 30 primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus.

- 35 In another embodiment, a vector capable of expressing NuABP or a fragment or derivative

thereof may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of NuABP including, but not limited to, those described above.

In a further embodiment, a pharmaceutical composition comprising a substantially purified NuABP in conjunction with a suitable pharmaceutical carrier may be administered to a subject to treat
5 or prevent a disorder associated with decreased expression or activity of NuABP including, but not limited to, those provided above.

In still another embodiment, an agonist which modulates the activity of NuABP may be administered to a subject to treat or prevent a disorder associated with decreased expression or activity of NuABP including, but not limited to, those listed above.

10 In a further embodiment, an antagonist of NuABP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of NuABP. Examples of such disorders include, but are not limited to, those reproductive, immune, and neurological disorders, and cell proliferative disorders including cancer, described above. In one aspect, an antibody which specifically binds NuABP may be used directly as an antagonist or indirectly as a targeting or delivery
15 mechanism for bringing a pharmaceutical agent to cells or tissues which express NuABP.

In an additional embodiment, a vector expressing the complement of the polynucleotide encoding NuABP may be administered to a subject to treat or prevent a disorder associated with increased expression or activity of NuABP including, but not limited to, those described above.

In other embodiments, any of the proteins, antagonists, antibodies, agonists, complementary
20 sequences, or vectors of the invention may be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy may be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents may act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic
25 efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

An antagonist of NuABP may be produced using methods which are generally known in the art. In particular, purified NuABP may be used to produce antibodies or to screen libraries of pharmaceutical agents to identify those which specifically bind NuABP. Antibodies to NuABP may also be generated using methods that are well known in the art. Such antibodies may include, but are
30 not limited to, polyclonal, monoclonal, chimeric, and single chain antibodies, Fab fragments, and fragments produced by a Fab expression library. Neutralizing antibodies (i.e., those which inhibit dimer formation) are generally preferred for therapeutic use.

For the production of antibodies, various hosts including goats, rabbits, rats, mice, humans, and others may be immunized by injection with NuABP or with any fragment or oligopeptide thereof
35 which has immunogenic properties. Depending on the host species, various adjuvants may be used to

increase immunological response. Such adjuvants include, but are not limited to, Freund's, mineral gels such as aluminum hydroxide, and surface active substances such as lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, KLH, and dinitrophenol. Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and Corynebacterium parvum are especially preferable.

5 It is preferred that the oligopeptides, peptides, or fragments used to induce antibodies to NuABP have an amino acid sequence consisting of at least about 5 amino acids, and generally will consist of at least about 10 amino acids. It is also preferable that these oligopeptides, peptides, or fragments are identical to a portion of the amino acid sequence of the natural protein and contain the entire amino acid sequence of a small, naturally occurring molecule. Short stretches of NuABP amino
10 acids may be fused with those of another protein, such as KLH, and antibodies to the chimeric molecule may be produced.

Monoclonal antibodies to NuABP may be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma
15 technique. (See, e.g., Kohler, G. et al. (1975) Nature 256:495-497; Kozbor, D. et al. (1985) J. Immunol. Methods 81:31-42; Cote, R.J. et al. (1983) Proc. Natl. Acad. Sci. USA 80:2026-2030; and Cole, S.P. et al. (1984) Mol. Cell Biol. 62:109-120.)

In addition, techniques developed for the production of "chimeric antibodies," such as the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate
20 antigen specificity and biological activity, can be used. (See, e.g., Morrison, S.L. et al. (1984) Proc. Natl. Acad. Sci. USA 81:6851-6855; Neuberger, M.S. et al. (1984) Nature 312:604-608; and Takeda, S. et al. (1985) Nature 314:452-454.) Alternatively, techniques described for the production of single chain antibodies may be adapted, using methods known in the art, to produce NuABP-specific single chain antibodies. Antibodies with related specificity, but of distinct idiotypic composition, may be
25 generated by chain shuffling from random combinatorial immunoglobulin libraries. (See, e.g., Burton, D.R. (1991) Proc. Natl. Acad. Sci. USA 88:10134-10137.)

Antibodies may also be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature. (See, e.g., Orlandi, R. et al. (1989) Proc. Natl. Acad. Sci. USA
30 86:3833-3837; Winter, G. et al. (1991) Nature 349:293-299.)

Antibody fragments which contain specific binding sites for NuABP may also be generated. For example, such fragments include, but are not limited to, F(ab')₂ fragments produced by pepsin digestion of the antibody molecule and Fab fragments generated by reducing the disulfide bridges of the F(ab')₂ fragments. Alternatively, Fab expression libraries may be constructed to allow rapid and
35 easy identification of monoclonal Fab fragments with the desired specificity. (See, e.g., Huse, W.D.

et al. (1989) Science 246:1275-1281.)

Various immunoassays may be used for screening to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays using either polyclonal or monoclonal antibodies with established specificities are well known in the art. Such immunoassays typically involve the measurement of complex formation between NuABP and its specific antibody. A two-site, monoclonal-based immunoassay utilizing monoclonal antibodies reactive to two non-interfering NuABP epitopes is generally used, but a competitive binding assay may also be employed (Pound, supra).

Various methods such as Scatchard analysis in conjunction with radioimmunoassay techniques may be used to assess the affinity of antibodies for NuABP. Affinity is expressed as an association constant, K_a , which is defined as the molar concentration of NuABP-antibody complex divided by the molar concentrations of free antigen and free antibody under equilibrium conditions. The K_a determined for a preparation of polyclonal antibodies, which are heterogeneous in their affinities for multiple NuABP epitopes, represents the average affinity, or avidity, of the antibodies for NuABP. The K_a determined for a preparation of monoclonal antibodies, which are monospecific for a particular NuABP epitope, represents a true measure of affinity. High-affinity antibody preparations with K_a ranging from about 10^9 to 10^{12} L/mole are preferred for use in immunoassays in which the NuABP-antibody complex must withstand rigorous manipulations. Low-affinity antibody preparations with K_a ranging from about 10^6 to 10^7 L/mole are preferred for use in immunopurification and similar procedures which ultimately require dissociation of NuABP, preferably in active form, from the antibody (Catty, D. (1988) Antibodies, Volume I: A Practical Approach, IRL Press, Washington, DC; Liddell, J.E. and Cryer, A. (1991) A Practical Guide to Monoclonal Antibodies, John Wiley & Sons, New York NY).

The titer and avidity of polyclonal antibody preparations may be further evaluated to determine the quality and suitability of such preparations for certain downstream applications. For example, a polyclonal antibody preparation containing at least 1-2 mg specific antibody/ml, preferably 5-10 mg specific antibody/ml, is generally employed in procedures requiring precipitation of NuABP-antibody complexes. Procedures for evaluating antibody specificity, titer, and avidity, and guidelines for antibody quality and usage in various applications, are generally available. (See, e.g., Catty, supra, and Coligan et al. supra.)

In another embodiment of the invention, the polynucleotides encoding NuABP, or any fragment or complement thereof, may be used for therapeutic purposes. In one aspect, the complement of the polynucleotide encoding NuABP may be used in situations in which it would be desirable to block the transcription of the mRNA. In particular, cells may be transformed with sequences complementary to polynucleotides encoding NuABP. Thus, complementary molecules or

fragments may be used to modulate NuABP activity, or to achieve regulation of gene function. Such technology is now well known in the art, and sense or antisense oligonucleotides or larger fragments can be designed from various locations along the coding or control regions of sequences encoding NuABP.

5 Expression vectors derived from retroviruses, adenoviruses, or herpes or vaccinia viruses, or from various bacterial plasmids, may be used for delivery of nucleotide sequences to the targeted organ, tissue, or cell population. Methods which are well known to those skilled in the art can be used to construct vectors to express nucleic acid sequences complementary to the polynucleotides encoding NuABP. (See, e.g., Sambrook, supra; Ausubel, 1995, supra.)

10 Genes encoding NuABP can be turned off by transforming a cell or tissue with expression vectors which express high levels of a polynucleotide, or fragment thereof, encoding NuABP. Such constructs may be used to introduce untranslatable sense or antisense sequences into a cell. Even in the absence of integration into the DNA, such vectors may continue to transcribe RNA molecules until they are disabled by endogenous nucleases. Transient expression may last for a month or more
15 with a non-replicating vector, and may last even longer if appropriate replication elements are part of the vector system.

As mentioned above, modifications of gene expression can be obtained by designing complementary sequences or antisense molecules (DNA, RNA, or PNA) to the control, 5', or regulatory regions of the gene encoding NuABP. Oligonucleotides derived from the transcription
20 initiation site, e.g., between about positions -10 and +10 from the start site, may be employed. Similarly, inhibition can be achieved using triple helix base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or regulatory molecules. Recent therapeutic advances using triplex DNA have been described in the literature. (See, e.g., Gee, J.E. et al. (1994) in Huber,
25 B.E. and B.I. Carr, Molecular and Immunologic Approaches, Futura Publishing, Mt. Kisco NY, pp. 163-177.) A complementary sequence or antisense molecule may also be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Ribozymes, enzymatic RNA molecules, may also be used to catalyze the specific cleavage of RNA. The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme
30 molecule to complementary target RNA, followed by endonucleolytic cleavage. For example, engineered hammerhead motif ribozyme molecules may specifically and efficiently catalyze endonucleolytic cleavage of sequences encoding NuABP.

Specific ribozyme cleavage sites within any potential RNA target are initially identified by scanning the target molecule for ribozyme cleavage sites, including the following sequences: GUA,
35 GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides,

corresponding to the region of the target gene containing the cleavage site, may be evaluated for secondary structural features which may render the oligonucleotide inoperable. The suitability of candidate targets may also be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays.

5 Complementary ribonucleic acid molecules and ribozymes of the invention may be prepared by any method known in the art for the synthesis of nucleic acid molecules. These include techniques for chemically synthesizing oligonucleotides such as solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by in vitro and in vivo transcription of DNA sequences encoding NuABP. Such DNA sequences may be incorporated into a wide variety of
10 vectors with suitable RNA polymerase promoters such as T7 or SP6. Alternatively, these cDNA constructs that synthesize complementary RNA, constitutively or inducibly, can be introduced into cell lines, cells, or tissues.

RNA molecules may be modified to increase intracellular stability and half-life. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends
15 of the molecule, or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the backbone of the molecule. This concept is inherent in the production of PNAs and can be extended in all of these molecules by the inclusion of nontraditional bases such as inosine, queosine, and wybutosine, as well as acetyl-, methyl-, thio-, and similarly modified forms of adenine, cytidine, guanine, thymine, and uridine which are not as easily recognized by endogenous endonucleases.

20 Many methods for introducing vectors into cells or tissues are available and equally suitable for use in vivo, in vitro, and ex vivo. For ex vivo therapy, vectors may be introduced into stem cells taken from the patient and clonally propagated for autologous transplant back into that same patient. Delivery by transfection, by liposome injections, or by polycationic amino polymers may be achieved using methods which are well known in the art. (See, e.g., Goldman, C.K. et al. (1997) Nat.
25 Biotechnol. 15:462-466.)

Any of the therapeutic methods described above may be applied to any subject in need of such therapy, including, for example, mammals such as humans, dogs, cats, cows, horses, rabbits, and monkeys.

An additional embodiment of the invention relates to the administration of a pharmaceutical
30 or sterile composition, in conjunction with a pharmaceutically acceptable carrier, for any of the therapeutic effects discussed above. Such pharmaceutical compositions may consist of NuABP, antibodies to NuABP, and mimetics, agonists, antagonists, or inhibitors of NuABP. The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier including,
35 but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered

to a patient alone, or in combination with other agents, drugs, or hormones.

The pharmaceutical compositions utilized in this invention may be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, enteral, topical, sublingual, or rectal means.

In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be found in the latest edition of Remington's
10 Pharmaceutical Sciences (Maack Publishing, Easton PA).

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

15 Pharmaceutical preparations for oral use can be obtained through combining active compounds with solid excipient and processing the resultant mixture of granules (optionally, after grinding) to obtain tablets or dragee cores. Suitable auxiliaries can be added, if desired. Suitable excipients include carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, and sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose,
20 hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums, including arabic and tragacanth; and proteins, such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, and alginic acid or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar
25 solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of
30 gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with fillers or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

35 Pharmaceutical formulations suitable for parenteral administration may be formulated in

aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds may be prepared as appropriate oily
5 injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils, such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate, triglycerides, or liposomes. Non-lipid polycationic amino polymers may also be used for delivery. Optionally, the suspension may also contain suitable stabilizers or agents to increase the solubility of the compounds and allow for the preparation of highly concentrated solutions.

10 For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

15 The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, and succinic acids. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preparation may be a lyophilized powder which may contain any or all of the following: 1 mM to 50 mM histidine, 0.1% to 2% sucrose, and 2% to 7% mannitol, at a
20 pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. For administration of NuABP, such labeling would include amount, frequency, and method of administration.

25 Pharmaceutical compositions suitable for use in the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended purpose. The determination of an effective dose is well within the capability of those skilled in the art.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models such as mice, rats, rabbits, dogs, or pigs. An animal model may also be used to determine the appropriate concentration range and route of
30 administration. Such information can then be used to determine useful doses and routes for administration in humans.

A therapeutically effective dose refers to that amount of active ingredient, for example NuABP or fragments thereof, antibodies of NuABP, and agonists, antagonists or inhibitors of NuABP, which ameliorates the symptoms or condition. Therapeutic efficacy and toxicity may be
35 determined by standard pharmaceutical procedures in cell cultures or with experimental animals, such

as by calculating the ED_{50} (the dose therapeutically effective in 50% of the population) or LD_{50} (the dose lethal to 50% of the population) statistics. The dose ratio of toxic to therapeutic effects is the therapeutic index, which can be expressed as the LD_{50}/ED_{50} ratio. Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies are used to formulate a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that includes the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, the sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject requiring treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, the general health of the subject, the age, weight, and gender of the subject, time and frequency of administration, drug combination(s), reaction sensitivities, and response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or biweekly depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts may vary from about 0.1 μg to 100,000 μg , up to a total dose of about 1 gram, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

DIAGNOSTICS

In another embodiment, antibodies which specifically bind NuABP may be used for the diagnosis of disorders characterized by expression of NuABP, or in assays to monitor patients being treated with NuABP or agonists, antagonists, or inhibitors of NuABP. Antibodies useful for diagnostic purposes may be prepared in the same manner as described above for therapeutics. Diagnostic assays for NuABP include methods which utilize the antibody and a label to detect NuABP in human body fluids or in extracts of cells or tissues. The antibodies may be used with or without modification, and may be labeled by covalent or non-covalent attachment of a reporter molecule. A wide variety of reporter molecules, several of which are described above, are known in the art and may be used.

A variety of protocols for measuring NuABP, including ELISAs, RIAs, and FACS, are known in the art and provide a basis for diagnosing altered or abnormal levels of NuABP expression.

Normal or standard values for NuABP expression are established by combining body fluids or cell

extracts taken from normal mammalian subjects, for example, human subjects, with antibody to NuABP under conditions suitable for complex formation. The amount of standard complex formation may be quantitated by various methods, such as photometric means. Quantities of NuABP expressed in subject, control, and disease samples from biopsied tissues are compared with the standard values.

5 Deviation between standard and subject values establishes the parameters for diagnosing disease.

In another embodiment of the invention, the polynucleotides encoding NuABP may be used for diagnostic purposes. The polynucleotides which may be used include oligonucleotide sequences, complementary RNA and DNA molecules, and PNAs. The polynucleotides may be used to detect and quantify gene expression in biopsied tissues in which expression of NuABP may be correlated
10 with disease. The diagnostic assay may be used to determine absence, presence, and excess expression of NuABP, and to monitor regulation of NuABP levels during therapeutic intervention.

In one aspect, hybridization with PCR probes which are capable of detecting polynucleotide sequences, including genomic sequences, encoding NuABP or closely related molecules may be used to identify nucleic acid sequences which encode NuABP. The specificity of the probe, whether it is
15 made from a highly specific region, e.g., the 5' regulatory region, or from a less specific region, e.g., a conserved motif, and the stringency of the hybridization or amplification will determine whether the probe identifies only naturally occurring sequences encoding NuABP, allelic variants, or related sequences.

Probes may also be used for the detection of related sequences, and may have at least 50%
20 sequence identity to any of the NuABP encoding sequences. The hybridization probes of the subject invention may be DNA or RNA and may be derived from the sequence of SEQ ID NO:56-110 or from genomic sequences including promoters, enhancers, and introns of the NuABP gene.

Means for producing specific hybridization probes for DNAs encoding NuABP include the cloning of polynucleotide sequences encoding NuABP or NuABP derivatives into vectors for the
25 production of mRNA probes. Such vectors are known in the art, are commercially available, and may be used to synthesize RNA probes in vitro by means of the addition of the appropriate RNA polymerases and the appropriate labeled nucleotides. Hybridization probes may be labeled by a variety of reporter groups, for example, by radionuclides such as ^{32}P or ^{35}S , or by enzymatic labels, such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems, and the like.

30 Polynucleotide sequences encoding NuABP may be used for the diagnosis of disorders associated with expression of NuABP. Examples of such disorders include, but are not limited to, a reproductive disorder such as disorders of prolactin production, infertility, including tubal disease, ovulatory defects, and endometriosis, disruptions of the estrous cycle, disruptions of the menstrual cycle, polycystic ovary syndrome, ovarian hyperstimulation syndrome, endometrial and ovarian
35 tumors, uterine fibroids, autoimmune disorders, ectopic pregnancies, and teratogenesis; cancer of the

breast, fibrocystic breast disease, and galactorrhea; disruptions of spermatogenesis, abnormal sperm physiology, cancer of the testis, cancer of the prostate, benign prostatic hyperplasia, prostatitis, Peyronie's disease, impotence, carcinoma of the male breast, and gynecomastia; an immune disorder such as inflammation, actinic keratosis, acquired immunodeficiency syndrome (AIDS), Addison's disease, adult respiratory distress syndrome, allergies, ankylosing spondylitis, amyloidosis, anemia, 5 arteriosclerosis, asthma, atherosclerosis, autoimmune hemolytic anemia, autoimmune thyroiditis, bronchitis, bursitis, cholecystitis, cirrhosis, contact dermatitis, Crohn's disease, atopic dermatitis, dermatomyositis, diabetes mellitus, emphysema, erythroblastosis fetalis, erythema nodosum, atrophic gastritis, glomerulonephritis, Goodpasture's syndrome, gout, Graves' disease, Hashimoto's thyroiditis, paroxysmal nocturnal hemoglobinuria, hepatitis, hypereosinophilia, irritable bowel 10 syndrome, episodic lymphopenia with lymphocytotoxins, mixed connective tissue disease (MCTD), multiple sclerosis, myasthenia gravis, myocardial or pericardial inflammation, myelofibrosis, osteoarthritis, osteoporosis, pancreatitis, polycythemia vera, polymyositis, psoriasis, Reiter's syndrome, rheumatoid arthritis, scleroderma, Sjögren's syndrome, systemic anaphylaxis, systemic lupus erythematosus, systemic sclerosis, primary thrombocythemia, thrombocytopenic purpura, 15 ulcerative colitis, uveitis, Werner syndrome, complications of cancer, hemodialysis, and extracorporeal circulation, trauma, and hematopoietic cancer including lymphoma, leukemia, and myeloma; a neurological disorder such as epilepsy, ischemic cerebrovascular disease, stroke, cerebral neoplasms, Alzheimer's disease, Pick's disease, Huntington's disease, dementia, Parkinson's disease 20 and other extrapyramidal disorders, amyotrophic lateral sclerosis and other motor neuron disorders, progressive neural muscular atrophy, retinitis pigmentosa, hereditary ataxias, multiple sclerosis and other demyelinating diseases, bacterial and viral meningitis, brain abscess, subdural empyema, epidural abscess, suppurative intracranial thrombophlebitis, myelitis and radiculitis, viral central nervous system disease, prion diseases including kuru, Creutzfeldt-Jakob disease, and Gerstmann- 25 Straussler-Scheinker syndrome, fatal familial insomnia, nutritional and metabolic diseases of the nervous system, neurofibromatosis, tuberous sclerosis, cerebelloretinal hemangioblastomatosis, encephalotrigeminal syndrome, mental retardation and other developmental disorders of the central nervous system, cerebral palsy, neuroskeletal disorders, autonomic nervous system disorders, cranial nerve disorders, spinal cord diseases, muscular dystrophy and other neuromuscular disorders, 30 peripheral nervous system disorders, dermatomyositis and polymyositis, inherited, metabolic, endocrine, and toxic myopathies, myasthenia gravis, periodic paralysis, mental disorders including mood, anxiety, and schizophrenic disorders, akathisia, amnesia, catatonia, diabetic neuropathy, tardive dyskinesia, dystonias, paranoid psychoses, postherpetic neuralgia, and Tourette's disorder; and a cell proliferative disorder such as actinic keratosis, arteriosclerosis, atherosclerosis, bursitis, 35 cirrhosis, hepatitis, mixed connective tissue disease (MCTD), myelofibrosis, paroxysmal nocturnal

hemoglobinuria, polycythemia vera, psoriasis, primary thrombocythemia, and cancers including adenocarcinoma, leukemia, lymphoma, melanoma, myeloma, sarcoma, teratocarcinoma, and, in particular, cancers of the adrenal gland, bladder, bone, bone marrow, brain, breast, cervix, gall bladder, ganglia, gastrointestinal tract, heart, kidney, liver, lung, muscle, ovary, pancreas, parathyroid, penis, prostate, salivary glands, skin, spleen, testis, thymus, thyroid, and uterus. The polynucleotide sequences encoding NuABP may be used in Southern or northern analysis, dot blot, or other membrane-based technologies; in PCR technologies; in dipstick, pin, and multiformat ELISA-like assays; and in microarrays utilizing fluids or tissues from patients to detect altered NuABP expression. Such qualitative or quantitative methods are well known in the art.

In a particular aspect, the nucleotide sequences encoding NuABP may be useful in assays that detect the presence of associated disorders, particularly those mentioned above. The nucleotide sequences encoding NuABP may be labeled by standard methods and added to a fluid or tissue sample from a patient under conditions suitable for the formation of hybridization complexes. After a suitable incubation period, the sample is washed and the signal is quantified and compared with a standard value. If the amount of signal in the patient sample is significantly altered in comparison to a control sample then the presence of altered levels of nucleotide sequences encoding NuABP in the sample indicates the presence of the associated disorder. Such assays may also be used to evaluate the efficacy of a particular therapeutic treatment regimen in animal studies, in clinical trials, or to monitor the treatment of an individual patient.

In order to provide a basis for the diagnosis of a disorder associated with expression of NuABP, a normal or standard profile for expression is established. This may be accomplished by combining body fluids or cell extracts taken from normal subjects, either animal or human, with a sequence, or a fragment thereof, encoding NuABP, under conditions suitable for hybridization or amplification. Standard hybridization may be quantified by comparing the values obtained from normal subjects with values from an experiment in which a known amount of a substantially purified polynucleotide is used. Standard values obtained in this manner may be compared with values obtained from samples from patients who are symptomatic for a disorder. Deviation from standard values is used to establish the presence of a disorder.

Once the presence of a disorder is established and a treatment protocol is initiated, hybridization assays may be repeated on a regular basis to determine if the level of expression in the patient begins to approximate that which is observed in the normal subject. The results obtained from successive assays may be used to show the efficacy of treatment over a period ranging from several days to months.

With respect to cancer, the presence of an abnormal amount of transcript (either under- or overexpressed) in biopsied tissue from an individual may indicate a predisposition for the

development of the disease, or may provide a means for detecting the disease prior to the appearance of actual clinical symptoms. A more definitive diagnosis of this type may allow health professionals to employ preventative measures or aggressive treatment earlier thereby preventing the development or further progression of the cancer.

5 Additional diagnostic uses for oligonucleotides designed from the sequences encoding NuABP may involve the use of PCR. These oligomers may be chemically synthesized, generated enzymatically, or produced in vitro. Oligomers will preferably contain a fragment of a polynucleotide encoding NuABP, or a fragment of a polynucleotide complementary to the polynucleotide encoding NuABP, and will be employed under optimized conditions for identification of a specific gene or
10 condition. Oligomers may also be employed under less stringent conditions for detection or quantification of closely related DNA or RNA sequences.

Methods which may also be used to quantify the expression of NuABP include radiolabeling or biotinylating nucleotides, coamplification of a control nucleic acid, and interpolating results from standard curves. (See, e.g., Melby, P.C. et al. (1993) J. Immunol. Methods 159:235-244; Duplaa, C.
15 et al. (1993) Anal. Biochem. 212:229-236.) The speed of quantitation of multiple samples may be accelerated by running the assay in a high-throughput format where the oligomer of interest is presented in various dilutions and a spectrophotometric or colorimetric response gives rapid quantitation.

In further embodiments, oligonucleotides or longer fragments derived from any of the
20 polynucleotide sequences described herein may be used as targets in a microarray. The microarray can be used to monitor the expression level of large numbers of genes simultaneously and to identify genetic variants, mutations, and polymorphisms. This information may be used to determine gene function, to understand the genetic basis of a disorder, to diagnose a disorder, and to develop and monitor the activities of therapeutic agents.

25 Microarrays may be prepared, used, and analyzed using methods known in the art. (See, e.g., Brennan, T.M. et al. (1995) U.S. Patent No. 5,474,796; Schena, M. et al. (1996) Proc. Natl. Acad. Sci. USA 93:10614-10619; Baldeschweiler et al. (1995) PCT application WO95/251116; Shalon, D. et al. (1995) PCT application WO95/35505; Heller, R.A. et al. (1997) Proc. Natl. Acad. Sci. USA 94:2150-2155; and Heller, M.J. et al. (1997) U.S. Patent No. 5,605,662.)

30 In another embodiment of the invention, nucleic acid sequences encoding NuABP may be used to generate hybridization probes useful in mapping the naturally occurring genomic sequence. The sequences may be mapped to a particular chromosome, to a specific region of a chromosome, or to artificial chromosome constructions, e.g., human artificial chromosomes (HACs), yeast artificial chromosomes (YACs), bacterial artificial chromosomes (BACs), bacterial P1 constructions, or single
35 chromosome cDNA libraries. (See, e.g., Harrington, J.J. et al. (1997) Nat. Genet. 15:345-355; Price,

C.M. (1993) Blood Rev. 7:127-134; and Trask, B.J. (1991) Trends Genet. 7:149-154.)

Fluorescent in situ hybridization (FISH) may be correlated with other physical chromosome mapping techniques and genetic map data. (See, e.g., Heinz-Ulrich, et al. (1995) in Meyers, supra, pp. 965-968.) Examples of genetic map data can be found in various scientific journals or at the

5 Online Mendelian Inheritance in Man (OMIM) World Wide Web site. Correlation between the location of the gene encoding NuABP on a physical chromosomal map and a specific disorder, or a predisposition to a specific disorder, may help define the region of DNA associated with that disorder. The nucleotide sequences of the invention may be used to detect differences in gene sequences among normal, carrier, and affected individuals.

10 In situ hybridization of chromosomal preparations and physical mapping techniques, such as linkage analysis using established chromosomal markers, may be used for extending genetic maps. Often the placement of a gene on the chromosome of another mammalian species, such as mouse, may reveal associated markers even if the number or arm of a particular human chromosome is not known. New sequences can be assigned to chromosomal arms by physical mapping. This provides

15 valuable information to investigators searching for disease genes using positional cloning or other gene discovery techniques. Once the disease or syndrome has been crudely localized by genetic linkage to a particular genomic region, e.g., ataxia-telangiectasia to 11q22-23, any sequences mapping to that area may represent associated or regulatory genes for further investigation. (See, e.g., Gatti, R.A. et al. (1988) Nature 336:577-580.) The nucleotide sequence of the subject invention may also be

20 used to detect differences in the chromosomal location due to translocation, inversion, etc., among normal, carrier, or affected individuals.

In another embodiment of the invention, NuABP, its catalytic or immunogenic fragments, or oligopeptides thereof can be used for screening libraries of compounds in any of a variety of drug screening techniques. The fragment employed in such screening may be free in solution, affixed to a

25 solid support, borne on a cell surface, or located intracellularly. The formation of binding complexes between NuABP and the agent being tested may be measured.

Another technique for drug screening provides for high throughput screening of compounds having suitable binding affinity to the protein of interest. (See, e.g., Geysen, et al. (1984) PCT application WO84/03564.) In this method, large numbers of different small test compounds are

30 synthesized on a solid substrate. The test compounds are reacted with NuABP, or fragments thereof, and washed. Bound NuABP is then detected by methods well known in the art. Purified NuABP can also be coated directly onto plates for use in the aforementioned drug screening techniques. Alternatively, non-neutralizing antibodies can be used to capture the peptide and immobilize it on a solid support.

35 In another embodiment, one may use competitive drug screening assays in which neutralizing

antibodies capable of binding NuABP specifically compete with a test compound for binding NuABP. In this manner, antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with NuABP.

In additional embodiments, the nucleotide sequences which encode NuABP may be used in any molecular biology techniques that have yet to be developed, provided the new techniques rely on properties of nucleotide sequences that are currently known, including, but not limited to, such properties as the triplet genetic code and specific base pair interactions.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The disclosures of all patents, applications, and publications mentioned above and below, in particular U.S. Ser. No. 60/117,905 and U.S. Ser. No. 60/117,904, are hereby expressly incorporated by reference.

20

EXAMPLES

I. Construction of cDNA Libraries

RNA was purchased from Clontech or isolated from tissues described in Table 4. Some tissues were homogenized and lysed in guanidinium isothiocyanate, while others were homogenized and lysed in phenol or in a suitable mixture of denaturants, such as TRIZOL (Life Technologies), a monophasic solution of phenol and guanidine isothiocyanate. The resulting lysates were centrifuged over CsCl cushions or extracted with chloroform. RNA was precipitated from the lysates with either isopropanol or sodium acetate and ethanol, or by other routine methods.

Phenol extraction and precipitation of RNA were repeated as necessary to increase RNA purity. In some cases, RNA was treated with DNase. For most libraries, poly(A⁺) RNA was isolated using oligo d(T)-coupled paramagnetic particles (Promega), OLIGOTEX latex particles (QIAGEN, Chatsworth CA), or an OLIGOTEX mRNA purification kit (QIAGEN). Alternatively, RNA was isolated directly from tissue lysates using other RNA isolation kits, e.g., the POLY(A)PURE mRNA purification kit (Ambion, Austin TX).

In some cases, Stratagene was provided with RNA and constructed the corresponding cDNA libraries. Otherwise, cDNA was synthesized and cDNA libraries were constructed with the UNIZAP

vector system (Stratagene) or SUPERScript plasmid system (Life Technologies), using the recommended procedures or similar methods known in the art. (See, e.g., Ausubel, 1997, supra, units 5.1-6.6.) Reverse transcription was initiated using oligo d(T) or random primers. Synthetic oligonucleotide adapters were ligated to double stranded cDNA, and the cDNA was digested with the appropriate restriction enzyme or enzymes. For most libraries, the cDNA was size-selected (300-1000 bp) using SEPHACRYL S1000, SEPHAROSE CL2B, or SEPHAROSE CL4B column chromatography (Amersham Pharmacia Biotech) or preparative agarose gel electrophoresis. cDNAs were ligated into compatible restriction enzyme sites of the polylinker of a suitable plasmid, e.g., PBLUEScript plasmid (Stratagene), PSPORT1 plasmid (Life Technologies), or pINCy (Incyte Pharmaceuticals, Palo Alto CA). Recombinant plasmids were transformed into competent E. coli cells including XL1-Blue, XL1-BlueMRF, or SOLR from Stratagene or DH5 α , DH10B, or ElectroMAX DH10B from Life Technologies.

II. Isolation of cDNA Clones

Plasmids were recovered from host cells by in vivo excision using the UNIZAP vector system (Stratagene) or by cell lysis. Plasmids were purified using at least one of the following: a Magic or WIZARD Minipreps DNA purification system (Promega); an AGTC Miniprep purification kit (Edge Biosystems, Gaithersburg MD); and QIAWELL 8 Plasmid, QIAWELL 8 Plus Plasmid, QIAWELL 8 Ultra Plasmid purification systems or the R.E.A.L. PREP 96 plasmid purification kit from QIAGEN. Following precipitation, plasmids were resuspended in 0.1 ml of distilled water and stored, with or without lyophilization, at 4°C.

Alternatively, plasmid DNA was amplified from host cell lysates using direct link PCR in a high-throughput format (Rao, V.B. (1994) Anal. Biochem. 216:1-14). Host cell lysis and thermal cycling steps were carried out in a single reaction mixture. Samples were processed and stored in 384-well plates, and the concentration of amplified plasmid DNA was quantified fluorometrically using PICOGREEN dye (Molecular Probes, Eugene OR) and a FLUOROSKAN II fluorescence scanner (Labsystems Oy, Helsinki, Finland).

III. Sequencing and Analysis

cDNA sequencing reactions were processed using standard methods or high-throughput instrumentation such as the ABI CATALYST 800 (Perkin-Elmer) thermal cycler or the PTC-200 thermal cycler (MJ Research) in conjunction with the HYDRA microdispenser (Robbins Scientific) or the MICROLAB 2200 (Hamilton) liquid transfer system. cDNA sequencing reactions were prepared using reagents provided by Amersham Pharmacia Biotech or supplied in ABI sequencing kits such as the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer). Electrophoretic separation of cDNA sequencing reactions and detection of labeled polynucleotides were carried out using the MEGABACE 1000 DNA sequencing system (Molecular Dynamics); the

ABI PRISM 373 or 377 sequencing system (Perkin-Elmer) in conjunction with standard ABI protocols and base calling software; or other sequence analysis systems known in the art. Reading frames within the cDNA sequences were identified using standard methods (reviewed in Ausubel, 1997, supra, unit 7.7). Some of the cDNA sequences were selected for extension using the techniques disclosed in Example V.

The polynucleotide sequences derived from cDNA sequencing were assembled and analyzed using a combination of software programs which utilize algorithms well known to those skilled in the art. Table 5 summarizes the tools, programs, and algorithms used and provides applicable descriptions, references, and threshold parameters. The first column of Table 5 shows the tools, programs, and algorithms used, the second column provides brief descriptions thereof, the third column presents appropriate references, all of which are incorporated by reference herein in their entirety, and the fourth column presents, where applicable, the scores, probability values, and other parameters used to evaluate the strength of a match between two sequences (the higher the score, the greater the homology between two sequences). Sequences were analyzed using MACDNASIS PRO software (Hitachi Software Engineering, South San Francisco CA) and LASERGENE software (DNASTAR). Polynucleotide and polypeptide sequence alignments were generated using the default parameters specified by the clustal algorithm as incorporated into the MEGALIGN multisequence alignment program (DNASTAR), which also calculates the percent identity between aligned sequences.

The polynucleotide sequences were validated by removing vector, linker, and polyA sequences and by masking ambiguous bases, using algorithms and programs based on BLAST, dynamic programming, and dinucleotide nearest neighbor analysis. The sequences were then queried against a selection of public databases such as the GenBank primate, rodent, mammalian, vertebrate, and eukaryote databases, and BLOCKS, PRINTS, DOMO, PRODOM, and PFAM to acquire annotation using programs based on BLAST, FASTA, and BLIMPS. The sequences were assembled into full length polynucleotide sequences using programs based on Phred, Phrap, and Consed, and were screened for open reading frames using programs based on GeneMark, BLAST, and FASTA. The full length polynucleotide sequences were translated to derive the corresponding full length amino acid sequences, and these full length sequences were subsequently analyzed by querying against databases such as the GenBank databases (described above), SwissProt, BLOCKS, PRINTS, DOMO, PRODOM, Prosite, and Hidden Markov Model (HMM)-based protein family databases such as PFAM. HMM is a probabilistic approach which analyzes consensus primary structures of gene families. (See, e.g., Eddy, S.R. (1996) Curr. Opin. Struct. Biol. 6:361-365.)

The programs described above for the assembly and analysis of full length polynucleotide and amino acid sequences were also used to identify polynucleotide sequence fragments from SEQ ID

NO:56-110. Fragments from about 20 to about 4000 nucleotides which are useful in hybridization and amplification technologies were described in The Invention section above.

IV. Northern Analysis

Northern analysis is a laboratory technique used to detect the presence of a transcript of a gene and involves the hybridization of a labeled nucleotide sequence to a membrane on which RNAs from a particular cell type or tissue have been bound. (See, e.g., Sambrook, supra, ch. 7; Ausubel, 1995, supra, ch. 4 and 16.)

Analogous computer techniques applying BLAST were used to search for identical or related molecules in nucleotide databases such as GenBank or LIFESEQ (Incyte Pharmaceuticals). This analysis is much faster than multiple membrane-based hybridizations. In addition, the sensitivity of the computer search can be modified to determine whether any particular match is categorized as exact or similar. The basis of the search is the product score, which is defined as:

$$\frac{\% \text{ sequence identity} \times \% \text{ maximum BLAST score}}{100}$$

100

The product score takes into account both the degree of similarity between two sequences and the length of the sequence match. For example, with a product score of 40, the match will be exact within a 1% to 2% error, and, with a product score of 70, the match will be exact. Similar molecules are usually identified by selecting those which show product scores between 15 and 40, although lower scores may identify related molecules.

The results of northern analyses are reported as a percentage distribution of libraries in which the transcript encoding NuABP occurred. Analysis involved the categorization of cDNA libraries by organ/tissue and disease. The organ/tissue categories included cardiovascular, dermatologic, developmental, endocrine, gastrointestinal, hematopoietic/immune, musculoskeletal, nervous, reproductive, and urologic. The disease/condition categories included cancer, inflammation, trauma, cell proliferation, neurological, and pooled. For each category, the number of libraries expressing the sequence of interest was counted and divided by the total number of libraries across all categories. Percentage values of tissue-specific and disease- or condition-specific expression are reported in Table 3.

V. Extension of NuABP Encoding Polynucleotides

The full length nucleic acid sequences of SEQ ID NO:56-110 were produced by extension of an appropriate fragment of the full length molecule using oligonucleotide primers designed from this fragment. One primer was synthesized to initiate 5' extension of the known fragment, and the other primer, to initiate 3' extension of the known fragment. The initial primers were designed using OLIGO 4.06 software (National Biosciences), or another appropriate program, to be about 22 to 30 nucleotides in length, to have a GC content of about 50% or more, and to anneal to the target

sequence at temperatures of about 68°C to about 72°C. Any stretch of nucleotides which would result in hairpin structures and primer-primer dimerizations was avoided.

Selected human cDNA libraries were used to extend the sequence. If more than one extension was necessary or desired, additional or nested sets of primers were designed.

5 High fidelity amplification was obtained by PCR using methods well known in the art. PCR was performed in 96-well plates using the PTC-200 thermal cycler (MJ Research, Inc.). The reaction mix contained DNA template, 200 nmol of each primer, reaction buffer containing Mg^{2+} , $(NH_4)_2SO_4$, and β -mercaptoethanol, Taq DNA polymerase (Amersham Pharmacia Biotech), ELONGASE enzyme (Life Technologies), and Pfu DNA polymerase (Stratagene), with the following parameters for primer pair PCI A and PCI B: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C. In the alternative, the parameters for primer pair T7 and SK+ were as follows: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 57°C, 1 min; Step 4: 68°C, 2 min; Step 5: Steps 2, 3, and 4 repeated 20 times; Step 6: 68°C, 5 min; Step 7: storage at 4°C.

15 The concentration of DNA in each well was determined by dispensing 100 μ l PICOGREEN quantitation reagent (0.25% (v/v) PICOGREEN; Molecular Probes, Eugene OR) dissolved in 1X TE and 0.5 μ l of undiluted PCR product into each well of an opaque fluorimeter plate (Corning Costar, Acton MA), allowing the DNA to bind to the reagent. The plate was scanned in a Fluoroskan II (Labsystems Oy, Helsinki, Finland) to measure the fluorescence of the sample and to quantify the concentration of DNA. A 5 μ l to 10 μ l aliquot of the reaction mixture was analyzed by electrophoresis on a 1 % agarose mini-gel to determine which reactions were successful in extending the sequence.

The extended nucleotides were desalted and concentrated, transferred to 384-well plates, digested with CviJI cholera virus endonuclease (Molecular Biology Research, Madison WI), and sonicated or sheared prior to religation into pUC 18 vector (Amersham Pharmacia Biotech). For 25 shotgun sequencing, the digested nucleotides were separated on low concentration (0.6 to 0.8%) agarose gels, fragments were excised, and agar digested with Agar ACE (Promega). Extended clones were religated using T4 ligase (New England Biolabs, Beverly MA) into pUC 18 vector (Amersham Pharmacia Biotech), treated with Pfu DNA polymerase (Stratagene) to fill-in restriction site overhangs, and transfected into competent *E. coli* cells. Transformed cells were selected on 30 antibiotic-containing media, individual colonies were picked and cultured overnight at 37°C in 384-well plates in LB/2x carb liquid media.

The cells were lysed, and DNA was amplified by PCR using Taq DNA polymerase (Amersham Pharmacia Biotech) and Pfu DNA polymerase (Stratagene) with the following 35 parameters: Step 1: 94°C, 3 min; Step 2: 94°C, 15 sec; Step 3: 60°C, 1 min; Step 4: 72°C, 2 min;

Step 5: steps 2, 3, and 4 repeated 29 times; Step 6: 72°C, 5 min; Step 7: storage at 4°C. DNA was quantified by PICOGREEN reagent (Molecular Probes) as described above. Samples with low DNA recoveries were reamplified using the same conditions as described above. Samples were diluted with 20% dimethylsulfoxide (1:2, v/v), and sequenced using DYENAMIC energy transfer sequencing primers and the DYENAMIC DIRECT kit (Amersham Pharmacia Biotech) or the ABI PRISM BIGDYE Terminator cycle sequencing ready reaction kit (Perkin-Elmer).

In like manner, the nucleotide sequences of SEQ ID NO:56-110 are used to obtain 5' regulatory sequences using the procedure above, oligonucleotides designed for such extension, and an appropriate genomic library.

10 VI. Labeling and Use of Individual Hybridization Probes

Hybridization probes derived from SEQ ID NO:56-110 are employed to screen cDNAs, genomic DNAs, or mRNAs. Although the labeling of oligonucleotides, consisting of about 20 base pairs, is specifically described, essentially the same procedure is used with larger nucleotide fragments. Oligonucleotides are designed using state-of-the-art software such as OLIGO 4.06 software (National Biosciences) and labeled by combining 50 pmol of each oligomer, 250 μ Ci of [γ -³²P] adenosine triphosphate (Amersham Pharmacia Biotech), and T4 polynucleotide kinase (DuPont NEN, Boston MA). The labeled oligonucleotides are substantially purified using a SEPHADEX G-25 superfine size exclusion dextran bead column (Amersham Pharmacia Biotech). An aliquot containing 10⁷ counts per minute of the labeled probe is used in a typical membrane-based hybridization analysis of human genomic DNA digested with one of the following endonucleases: Ase I, Bgl II, Eco RI, Pst I, Xba I, or Pvu II (DuPont NEN).

The DNA from each digest is fractionated on a 0.7% agarose gel and transferred to nylon membranes (Nytran Plus, Schleicher & Schuell, Durham NH). Hybridization is carried out for 16 hours at 40°C. To remove nonspecific signals, blots are sequentially washed at room temperature under conditions of up to, for example, 0.1 x saline sodium citrate and 0.5% sodium dodecyl sulfate. Hybridization patterns are visualized using autoradiography or an alternative imaging means and compared.

25 VII. Microarrays

A chemical coupling procedure and an ink jet device can be used to synthesize array elements on the surface of a substrate. (See, e.g., Baldeschweiler, *supra*.) An array analogous to a dot or slot blot may also be used to arrange and link elements to the surface of a substrate using thermal, UV, chemical, or mechanical bonding procedures. A typical array may be produced by hand or using available methods and machines and contain any appropriate number of elements. After hybridization, nonhybridized probes are removed and a scanner used to determine the levels and patterns of fluorescence. The degree of complementarity and the relative abundance of each probe

which hybridizes to an element on the microarray may be assessed through analysis of the scanned images.

Full-length cDNAs, Expressed Sequence Tags (ESTs), or fragments thereof may comprise the elements of the microarray. Fragments suitable for hybridization can be selected using software well known in the art such as LASERGENE software (DNASTAR). Full-length cDNAs, ESTs, or fragments thereof corresponding to one of the nucleotide sequences of the present invention, or selected at random from a cDNA library relevant to the present invention, are arranged on an appropriate substrate, e.g., a glass slide. The cDNA is fixed to the slide using, e.g., UV cross-linking followed by thermal and chemical treatments and subsequent drying. (See, e.g., Schena, M. et al. (1995) Science 270:467-470; Shalon, D. et al. (1996) Genome Res. 6:639-645.) Fluorescent probes are prepared and used for hybridization to the elements on the substrate. The substrate is analyzed by procedures described above.

VIII. Complementary Polynucleotides

Sequences complementary to the NuABP-encoding sequences, or any parts thereof, are used to detect, decrease, or inhibit expression of naturally occurring NuABP. Although use of oligonucleotides comprising from about 15 to 30 base pairs is described, essentially the same procedure is used with smaller or with larger sequence fragments. Appropriate oligonucleotides are designed using OLIGO 4.06 software (National Biosciences) and the coding sequence of NuABP. To inhibit transcription, a complementary oligonucleotide is designed from the most unique 5' sequence and used to prevent promoter binding to the coding sequence. To inhibit translation, a complementary oligonucleotide is designed to prevent ribosomal binding to the NuABP-encoding transcript.

IX. Expression of NuABP

Expression and purification of NuABP is achieved using bacterial or virus-based expression systems. For expression of NuABP in bacteria, cDNA is subcloned into an appropriate vector containing an antibiotic resistance gene and an inducible promoter that directs high levels of cDNA transcription. Examples of such promoters include, but are not limited to, the *trp-lac (tac)* hybrid promoter and the T5 or T7 bacteriophage promoter in conjunction with the *lac* operator regulatory element. Recombinant vectors are transformed into suitable bacterial hosts, e.g., BL21(DE3).

Antibiotic resistant bacteria express NuABP upon induction with isopropyl beta-D-

thiogalactopyranoside (IPTG). Expression of NuABP in eukaryotic cells is achieved by infecting insect or mammalian cell lines with recombinant Autographica californica nuclear polyhedrosis virus (AcMNPV), commonly known as baculovirus. The nonessential polyhedrin gene of baculovirus is replaced with cDNA encoding NuABP by either homologous recombination or bacterial-mediated transposition involving transfer plasmid intermediates. Viral infectivity is maintained and the strong polyhedrin promoter drives high levels of cDNA transcription. Recombinant baculovirus is used to

infect Spodoptera frugiperda (Sf9) insect cells in most cases, or human hepatocytes, in some cases. Infection of the latter requires additional genetic modifications to baculovirus. (See Engelhard, E.K. et al. (1994) Proc. Natl. Acad. Sci. USA 91:3224-3227; Sandig, V. et al. (1996) Hum. Gene Ther. 7:1937-1945.)

5 In most expression systems, NuABP is synthesized as a fusion protein with, e.g., glutathione S-transferase (GST) or a peptide epitope tag, such as FLAG or 6-His, permitting rapid, single-step, affinity-based purification of recombinant fusion protein from crude cell lysates. GST, a 26-kilodalton enzyme from Schistosoma japonicum, enables the purification of fusion proteins on immobilized glutathione under conditions that maintain protein activity and antigenicity (Amersham
10 Pharmacia Biotech). Following purification, the GST moiety can be proteolytically cleaved from NuABP at specifically engineered sites. FLAG, an 8-amino acid peptide, enables immunoaffinity purification using commercially available monoclonal and polyclonal anti-FLAG antibodies (Eastman Kodak). 6-His, a stretch of six consecutive histidine residues, enables purification on metal-chelate resins (QIAGEN). Methods for protein expression and purification are discussed in Ausubel (1995,
15 supra, ch. 10 and 16). Purified NuABP obtained by these methods can be used directly in the following activity assay.

X. Demonstration of NuABP Activity

NuABP activity is measured by its ability to stimulate transcription of a reporter gene (Liu, H.Y. et al. (1997) EMBO J. 16(17):5289-5298.) The assay entails the use of a well characterized
20 reporter gene construct, LexA_{op}-LacZ, that consists of LexA DNA transcriptional control elements (LexA_{op}) fused to sequences encoding the E. coli LacZ enzyme. The methods for constructing and expressing fusions genes, introducing them into cells, and measuring LacZ enzyme activity, are well known to those skilled in the art. Sequences encoding NuABP are cloned into a plasmid that directs the synthesis of a fusion protein, LexA-NuABP, consisting of NuABP and a DNA binding domain
25 derived from the LexA transcription factor. The resulting plasmid, encoding a LexA-NuABP fusion protein, is introduced into yeast cells along with a plasmid containing the LexA_{op}-LacZ reporter gene. The amount of LacZ enzyme activity associated with LexA-NuABP transfected cells, relative to control cells, is proportional to the amount of transcription stimulated by the NuABP.

XI. Functional Assays

30 NuABP function is assessed by expressing the sequences encoding NuABP at physiologically elevated levels in mammalian cell culture systems. cDNA is subcloned into a mammalian expression vector containing a strong promoter that drives high levels of cDNA expression. Vectors of choice include pCMV SPORT (Life Technologies) and pCR3.1 (Invitrogen, Carlsbad CA), both of which contain the cytomegalovirus promoter. 5-10 μ g of recombinant vector
35 are transiently transfected into a human cell line, for example, an endothelial or hematopoietic cell

line, using either liposome formulations or electroporation. 1-2 μ g of an additional plasmid containing sequences encoding a marker protein are co-transfected. Expression of a marker protein provides a means to distinguish transfected cells from nontransfected cells and is a reliable predictor of cDNA expression from the recombinant vector. Marker proteins of choice include, e.g., Green
5 Fluorescent Protein (GFP; Clontech), CD64, or a CD64-GFP fusion protein. Flow cytometry (FCM), an automated, laser optics-based technique, is used to identify transfected cells expressing GFP or CD64-GFP and to evaluate the apoptotic state of the cells and other cellular properties. FCM detects and quantifies the uptake of fluorescent molecules that diagnose events preceding or coincident with cell death. These events include changes in nuclear DNA content as measured by staining of DNA
10 with propidium iodide; changes in cell size and granularity as measured by forward light scatter and 90 degree side light scatter; down-regulation of DNA synthesis as measured by decrease in bromodeoxyuridine uptake; alterations in expression of cell surface and intracellular proteins as measured by reactivity with specific antibodies; and alterations in plasma membrane composition as measured by the binding of fluorescein-conjugated Annexin V protein to the cell surface. Methods in
15 flow cytometry are discussed in Ormerod, M.G. (1994) Flow Cytometry, Oxford, New York NY.

The influence of NuABP on gene expression can be assessed using highly purified populations of cells transfected with sequences encoding NuABP and either CD64 or CD64-GFP. CD64 and CD64-GFP are expressed on the surface of transfected cells and bind to conserved regions of human immunoglobulin G (IgG). Transfected cells are efficiently separated from nontransfected
20 cells using magnetic beads coated with either human IgG or antibody against CD64 (DYNAL, Lake Success NY). mRNA can be purified from the cells using methods well known by those of skill in the art. Expression of mRNA encoding NuABP and other genes of interest can be analyzed by northern analysis or microarray techniques.

XII. Production of NuABP Specific Antibodies

25 NuABP substantially purified using polyacrylamide gel electrophoresis (PAGE; see, e.g., Harrington, M.G. (1990) *Methods Enzymol.* 182:488-495), or other purification techniques, is used to immunize rabbits and to produce antibodies using standard protocols.

Alternatively, the NuABP amino acid sequence is analyzed using LASERGENE software (DNASTAR) to determine regions of high immunogenicity, and a corresponding oligopeptide is
30 synthesized and used to raise antibodies by means known to those of skill in the art. Methods for selection of appropriate epitopes, such as those near the C-terminus or in hydrophilic regions are well described in the art. (See, e.g., Ausubel, 1995, supra, ch. 11.)

Typically, oligopeptides of about 15 residues in length are synthesized using an ABI 431A peptide synthesizer (Perkin-Elmer) using fmoc-chemistry and coupled to KLH (Sigma-Aldrich, St.
35 Louis MO) by reaction with N-maleimidobenzoyl-N-hydroxysuccinimide ester (MBS) to increase

immunogenicity. (See, e.g., Ausubel, 1995, supra.) Rabbits are immunized with the oligopeptide-KLH complex in complete Freund's adjuvant. Resulting antisera are tested for antipeptide and anti-NuABP activity by, for example, binding the peptide or NuABP to a substrate, blocking with 1% BSA, reacting with rabbit antisera, washing, and reacting with radio-iodinated goat anti-rabbit IgG.

5 **XIII. Purification of Naturally Occurring NuABP Using Specific Antibodies**

Naturally occurring or recombinant NuABP is substantially purified by immunoaffinity chromatography using antibodies specific for NuABP. An immunoaffinity column is constructed by covalently coupling anti-NuABP antibody to an activated chromatographic resin, such as CNBr-activated SEPHAROSE (Amersham Pharmacia Biotech). After the coupling, the resin is
10 blocked and washed according to the manufacturer's instructions.

Media containing NuABP are passed over the immunoaffinity column, and the column is washed under conditions that allow the preferential absorbance of NuABP (e.g., high ionic strength buffers in the presence of detergent). The column is eluted under conditions that disrupt antibody/NuABP binding (e.g., a buffer of pH 2 to pH 3, or a high concentration of a chaotrope, such
15 as urea or thiocyanate ion), and NuABP is collected.

XIV. Identification of Molecules Which Interact with NuABP

NuABP, or biologically active fragments thereof, are labeled with ¹²⁵I Bolton-Hunter reagent. (See, e.g., Bolton A.E. and W.M. Hunter (1973) Biochem. J. 133:529-539.) Candidate molecules previously arrayed in the wells of a multi-well plate are incubated with the labeled NuABP, washed,
20 and any wells with labeled NuABP complex are assayed. Data obtained using different concentrations of NuABP are used to calculate values for the number, affinity, and association of NuABP with the candidate molecules.

Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention.
25 Although the invention has been described in connection with certain embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the following claims.

TABLE 1

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
1	56	025733	SPLNFET01	025733H1 and 025733X307D2 (SPLNFET01), 1519809F1 and 1519809T1 (BLADTUT04), 1526288T6 (UCMCL5T01), 1595557X16C1 and 1595557X19C1 (BRAINT014), 1903359F6 (OVARNOT07), 2417225F6 (HNT3AGT01), 4283542H1 (LIVRDIR01)
2	57	079702	SYNORAB01	002190H1 (U937NOT01), 079479F1 and 079702H1 (SYNORAB01), 125223T6 (LUNGNOT01), 371006R6 (LUNGNOT02), 2129460T6 (KIDNNOT05), 2999366H1 (TLYMNOT06), 3031905F6 (TLYMNOT05), 4949863H1 (SINTNOT25)
3	58	116208	KIDNNOT01	116208H1 and 116208R1 (KIDNNOT01), 2293058R6 (BRAINO01), 3731418H1 and 3731418T6 (SMCCNON03)
4	59	179261	PLACNOB01	179261H1 (PLACNOB01), 3666231F6 and 3666231T6 (PANCNOT16)
5	60	259161	HNT2RAT01	259161H1 (HNT2RAT01), 1005021R6 (BRSTNOT03), 2634660H1 (COLNTUT15), 2894335H1 (KIDNTUT14), 2924845H1 (SININOT04), 3659440H1 (ENDPNOT02), SBMA02955F1, SBMA03577F1, SBMA01445F1, SBMA00985F1, SBMA01499F1
6	61	320087	EOSIHET02	016657F1 (HUELPOB1), 320087H1 (EOSIHET02), 824110R1 (PROSNOT06), 987467H1 (LVENNOT03), 1235752F1 (LUNGNET03), 1361280F1 (LUNGNOT12), 1389740H1 (EOSINOT01), 1534332F1 (SPLNNOT04), 1813754F6 (SKINBIT01), 4184915H1 (BRSTNOT31), 5306522H1 (MONOTXT02)
7	62	491271	HNT2AGT01	363816X3 and 363816X9 (PROSNOT01), 491271F1, 491271H1 and 491271T6 (HNT2AGT01), 967354R6, 967354X15 and 967354X27 (BRSTNOT05), 2733444H1 (OVARTUT04)
8	63	585172	PROSNOT02	395188R6 (TMLR2DT01), 585172H1 (PROSNOT02), 864269T1 (BRAITUT03), 1417965F1 (KIDNNOT09)
9	64	615200	COLNTUT02	615200H1 and 615200R6 (COLNTUT02), 1213980R1 (BRSTTUT01), SBPA02731D1, SBPA00184D1
10	65	997067	KIDNTUT01	125981X3 (LUNGNOT01), 997067H1, 997067R6 and 997067T6 (KIDNTUT01), 1448201H1 (PLACNOT02), 1663447H1 (BRSTNOT09), 1889314H1 (BLADTUT07), 1918706H1 (PROSNOT06), 2699956H1 (OVARTUT10), 2702585H1 (OVARTUT10), 2900479H1 (DRGCNOT01), 3595727T6 (FIBPNOT01), 4309131H1 (BRAUNOT01)
11	66	144326 2	THYRN0T03	1257005F1 (MENITUT03), 1443262H1 (THYRN0T03), 1618906F6 (BRAITUT12), 2474133T6 (THP1NOT03), 3594075H1 (FIBPNOT01), 4914442H1 (LIVRFET05)

TABLE 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
12	67	152164 8	BLADTUT04	292172H1 (TMLR3DT01), 819925R6 (KERANOT02), 1353913H1 (LATRTUT02), 1521648H1 and 1522364F1 (BLADTUT04), 1963178H1 (BRSTNOT04), 2342505F6 (TESTTUT02), 4899970H1 (OVARDT01), 5043585H2 (PLACFER01)
13	68	168549 4	PROSNOT15	903980X13, 903980X14 and 903980X17 (COLNNOT07), 1685494H1 (PROSNOT15), 4164127T6 (BRSTNOT32)
14	69	173082 9	BRSTTUT08	116146R1 (KIDNNOT01), 836856R1 (PROSNOT07), 1730829H1, 1730829X11C1, 1730829X12C1 and 1730829X13C1 (BRSTTUT08), 1959889R6 (BRSTNOT04), 2188079H1 (PROSNOT26), 3384625H1 (ESOGNOT04)
15	70	186464 1	PROSNOT19	1844972H1 (COLNNOT08), 1864641F6 and 1864641H1 (PROSNOT19), 3090702T6 (BRSTNOT19), 3411665H1 (BRSTTUS08), 5152366H1 (HEARFET03), 5166179H1 (MUSCDMT01)
16	71	244460 4	THP1NOT03	1506658F1 (BRAITUT07), 1532034F1 (SPLNNOT04), 2444604H1 (THP1NOT03)
17	72	244500 8	THP1NOT03	605598X12 (BRSTTUT01), 628644H1 (KIDNNOT05), 732124R1 (LUNGNOT03), 819194R1 (KERANOT02), 1259467H1 (MENITUT03), 1363205F6 (LUNGNOT12), 1901312T6 (BLADTUT06), 2445008H1 (THP1NOT03), 2681125H1 (SINIUCT01)
18	73	257246 2	HIPOAZT01	396323R6 (PITUNOT02), 863622H1 (BRAITUT03), 1848956F6 and 1848956T6 (LUNGFET03), 2345947H1 (TESTTUT02), 2396384F6 (THP1AZT01), 2572462H1 (HIPOAZT01), 2650980F6 (LUNGUT02), 2814325H1 (OVARNOT10), 5076051H1 (COLCTUT03)
19	74	257289 2	HIPOAZT01	030596X15R1 (THPINOB01), 539564X11 (LNODNOT02), 1275514F1 and 1275514T6 (TESTTUT02), 2112383H1 (BRAITUT03), 2572892H1 (HIPOAZT01), 2986518H1 (CARGDIT01)
20	75	278567 4	BRSTNOT13	261399H1 (HNT2AGT01), 1274739F1 (TESTTUT02), 2785674H1 (BRSTNOT13)
21	76	279747 9	NPOLNOT01	302614X13 (TESTNOT04), 2797479H1 (NPOLNOT01), SAIA02597F1, SAIA00739F1, SAIA02537F1
22	77	296064 0	ADRENOT09	027211R1, 027211X1 and 027211X3 (SPLNFET01), 1401538F6 (BRAITUT08), 2496984F6 (ADRETUT05), 2960640H1 (ADRENOT09), 3211036T6 (BLADNOT08)
23	78	345405 1	SPLNNOT11	279331R6 (LIVRNUT02), 2515972T6 and 2516010T6 (LIVRTUT04), 2910726F6 (KIDNTUT15), 3454051H1 (SPLNNOT11)

TABLE 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
24	79	351064 0	CONCNOT01	556354H1 (MPHGLPT02), 581085H1 (BRAVXT05), 990636R6 (COLNNOT11), 1799185F6 and 1799185T6 (COLNNOT27), 3510640H1 (CONCNOT01), 4326648H1 (TLYMUNT01), SBHA01099F1
25	80	381508 3	TONSNOT03	2026951R6 and 2026951T6 (KERANOT02), 2300211R6 (BRSTNOT05), 2505283F6 (CONUTUT01), 3187267R6 (THYMNON04), 3815083H1 and 3815083T6 (TONSNOT03)
26	81	398845 7	LUNGNON03	609622R6 (COLNNOT01), 1710465F6 (PROSNOT16), 3988457H1 (LUNGNON03), SAQA00089F1, SAQA03055F1, SAQB00279F1
27	82	131890	BMARNOT02	131890H1 (BMARNOT02), 131890T6 (BMARNOT02), 132849R6 (BMARNOT02), 3357071F6 (PROSTUT16)
28	83	238642	SINTNOT02	238642H1 (SINTNOT02), 1620593F6 (BRAITUT13), 1620593H1 (BRAITUT13), 1620593T6 (BRAITUT13), 2534087F6 (BRAINOT18)
29	84	669862	CRBLNOT01	347231X7 (THYMNOT02), 669862H1 (CRBLNOT01), 2244458R6 (HIPONON02), 2244458T6 (HIPONON02), 2622610T6 (KERANOT02), 3536262H1 (KIDNNOT25), 4204212H1 (BRAITUT29)
30	85	100366 3	BRSTNOT03	850478T1 (NGANNOT01), 1003663H1 (BRSTNOT03), 1252179F2 (LUNGFET03), 1293336F1 (PGANNOT03), 1813002F6 (PROSTUT12), 2101974R6 (BRAITUT02)
31	86	143255 7	BEPINON01	1314545F6 (BLADTUT02), 1432557H1 (BEPINON01), 1443311R1 (THYRNOT03), 1705738F6 (DUODNOT02), 2182184F6 (SININOT01)
32	87	144177 0	THYRNOT03	035105R1 (HUVENOB01), 1441770H1 (THYRNOT03), 1500943F1 (SINTBST01), 2542840H1 (UTRSNOT11), 4533672H1 (OVARNOT12)
33	88	145668 4	COLNFET02	1456684F6 (COLNFET02), 1456684H1 (COLNFET02), 1456684T6 (COLNFET02), 1992143H1 (CORPNOT02), 2687476F6 (LUNGNOT23), 3139175F6 (SMCCNOT02), 4746319H1 (SMCRUNT01)
34	89	160291 6	BLADNOT03	3397976X305D2 (UTRSNOT16)
35	90	169281 6	COLNNOT23	999017R6 (KIDNTUT01), 1342490T1 (COLNTUT03), 1421981F1 (KIDNNOT09), 1692816H1 (COLNNOT23), 2176832F6 (ENDCNOT03), 2451404F6 (ENDANOT01)
36	91	196819 1	BRSTNOT04	1968191H1 (BRSTNOT04), 1968191T6 (BRSTNOT04), 2752967R6 (THPLAZS08), 3281163T6 (STOMFET02), 3748607H1 (UTRSNOT18)
37	92	205206 1	LIVRFET02	003803X8 (HMC1NOT01), 027044X1 (SPLNFET01), 027044X101 (SPLNFET01), 2052061H1 (LIVRFET02), 3931936F6 (PROSTUT09)

TABLE 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
38	93	205620 7	BEPINOT01	071525F1 (PLACNOB01), 162720R1 (ADENINB01), 270498H1 (HNT2NOT01), 1477853T1 (CORPNOT02), 1931058F6 (COLNTUT03), 2056207H1 (BEPINOT01), 2056207X11R1 (BEPINOT01), 2231060F6 (PROSNOT16), 2420063X309D4 (SCORNONOT2), 3424630H1 (BRSTNOR01), 3873760F6 (HEARNOT06), 3873760T6 (HEARNOT06), SBQA00627D1, SCJA02192V1, SCJA02089V1
39	94	210180 3	BRAITUT02	399584R6 (PITUNOT02), 399584T6 (PITUNOT02), 1649058F6 (PROSTUT09), 1902809F6 (OVARNOT07), 2101803H1 (BRAITUT02), 2101803R6 (BRAITUT02), 3098623H1 (CERVNOT03)
40	95	211236 2	BRAITUT03	948628R1 (PANCNOT05), 1209447T1 (BRSTNOT02), 1814624F6 (PROSNOT20), 2112362H1 (BRAITUT03), 2945621H1 (BRAITUT23), 3285663H1 (HEAONOT05), 3526403H1 (ESOGTUN01), 5033729H1 (ENDIUNT01), 5099417H1 (PROSTUS20)
41	96	211734 6	BRSTTUT02	487994R6 (HNT2AGT01), 952855R1 (SCORNONOT1), 952855T1 (SCORNONOT1), 2117346H1 (BRSTTUT02), 2458342F6 (ENDANOT01), 2731585H1 (OVARNOT04), 3475780H1 (LUNGNOT27), 3538525F6 (SEMNOT04)
42	97	211991 7	BRSTTUT02	2119917H1 (BRSTTUT02), 2791421F6 (COLNTUT16), 2794083F6 (COLNTUT16), 5006921H1 (STOMNOT08)
43	98	212345 6	BRSTNOT07	484031H1 (HNT2RAT01), 617559F1 (PGANNOT01), 617559R1 (PGANNOT01), 1575977F1 (LNODNOT03), 2123456H1 (BRSTNOT07), 2958712H1 (ADRENOT09), 3764961H1 (BRSTNOT24)
44	99	214879 2	BRAINOT09	1732781F6 (BRSTTUT08), 2050885F6 (LIVRFET02), 2148792H1 (BRAINOT09), 2590822H1 (LUNGNOT22), 2972368T6 (HEAONOT02), SBQA00396D1, SBQA03678D1, SBQA02120D1, SBQA03269D1
45	100	275194 3	THPLAZS08	1720187X16C1 (BLADNOT06), 2751943H1 (THPLAZS08), 3492378H1 (ADRETUT07)
46	101	312891 3	LUNGUTUT12	2551859F6 (LUNGUTUT06), 3128913H1 (LUNGUTUT12), SBMA01861F1, SBMA02298F1, SBMA01013F1, SBMA02403F1, SBMA01362F1
47	102	328294 1	HEAONOT05	154741R6 (THP1PLB02), 155904R6 (THP1PLB02), 157816R1 (THP1PLB02), 979920H1 (TONGTUT01), 1233933T6 (LUNGFTET03), 1657077F6 (URETTUT01), 2445017F6 (THPINOT03), 3282941H1 (HEAONOT05), 3341633H1 (SPLNNOT09), 3517140H1 (LUNGNONOT03)
48	103	328665 6	HEAONOT05	898123H1 (BRSTNOT05), 3286656H1 (HEAONOT05), 3641429T6 (LUNGNOT30), 3657668F6 (ENDPNOT02)

TABLE 1 (cont.)

Protein SEQ ID NO:	Nucleotide SEQ ID NO:	Clone ID	Library	Fragments
49	104	349080 2	EPIGNOT01	2238441H1 (PANCYTUT02), 2700133F6 (OVARTUT10), 2700133T6 (OVARTUT10), 3490226H1 (EPIGNOT01), 3490802H1 (EPIGNOT01), 4822929H1 (PROSTUT17)
50	105	350736 6	CONCNOT01	2130284H1 (KIDNNOT05), 3507366H1 (CONCNOT01), 3557087F6 (LUNGNOT31), 4241774H1 (SYNWDIT01)
51	105	357306 0	BRONNOT01	3573060F6 (BRONNOT01), 3573060H1 (BRONNOT01), 3573060T6 (BRONNOT01), 3867263H1 (BRAITUT07), 5013346H1 (BRAXNOT03)
52	107	357366 1	BRONNOT01	3028034F6 (HEARFET02), 3152642H1 (ADRENON04), 3573661F6 (BRONNOT01), 3573661H1 (BRONNOT01), 3577568F6 (BRONNOT01)
53	108	363342 2	LIVRNOT03	033412R6 (THP1NOB01), 074123F1 (THP1PEB01), 263241H1 (HNT2AGT01), 748567R1 (BRAITUT01), 1292088T1 (PGANNOT03), 1517449T1 (PANCYTUT01), 3633422H1 (LIVRNOT03)
54	109	399337 7	LUNGNON03	3003233H1 (TLYMNNOT06), 3993377H1 (LUNGNON03), 3993377T6 (LUNGNON03), 4251662F6 (BRADDIR01), SBSA02001V1
55	110	471793 6	BRAIHCT02	4717936F6 (BRAIHCT02), 4717936H1 (BRAIHCT02), 4717936T6 (BRAIHCT02)

TABLE 2

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
1	754	S6 T15 S22 S26 S28 S30 S60 Y84 S98 S102 S103 S112 S121 S146 T166 T183 T184 S231 Y253 S303 T304 S308 T327 T361 T393 T394 T399 T448 S496 Y583 S586 T608 S635 S672 S673 S682 S691 S711 S713 S751		L435 - P474 PFAM	DNA polymerase	PFAM, BLOCKS, MOTIFS
2	593	S32 Y62 T65 S83 S141 T159 T160 S185 S254 S288 T311 S314 S333 T368 S380 S401 T410 S426 S452 S461 S479 T483 T485 T576		E102 - A157 Q201 - C252	PHD finger DNA binding protein (GI 3342452)	BLAST, PFAM, MOTIFS
3	534	T39 S52 S301 T344 S373 T404 S425 S438 S439 S440 S473 T480 S490 T527		I17 - D57 PFAM	DNA polymerase	PFAM, BLOCKS, MOTIFS
4	255	S10 S26 Y35 Y113 S149 T168 Y169 T248		Y58 - C86 F141 - H163 Y169 - H191 Y197 - H219 Y225 - H247 PFAM	C2H2-type zinc finger protein (GI 498721)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS

TABLE 2 (cont.)

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
5	562	T29 T43 S76 S142 S165 S202 T214 Y302 S305 Y349 S385 S500 T526 S527		Y235 - D312 PFAM	DNA helicase (GI 531243; SEQ ID NO:113)	BLAST, PFAM, BLOCKS, MOTIFS
6	432	S33 S58 T166 T172 S197 T230 T261 S275 S286 S290 S298 S338 T362 S376 T407 T409 T419		E329 - A355 BLOCKS	CCAAT-box-binding transcription factor	BLOCKS, MOTIFS
7	799	T24 S33 T43 S73 T88 S91 S110 S147 T219 S262 S323 Y380 S532 S586 S756 T795		H250 - H291 Y324 - H346 Y253 - H374 Y380 - H402 PFAM	C2H2-type zinc finger protein (GI 498727)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
8	137	S3 T38 S74 S75 S118		R85 - L97 PFAM	BTB domain/C2H2-type zinc finger protein	PFAM, PRINTS, MOTIFS
9	230	T178 S187			sirtuin type 3 (GI 5225322)	BLAST, MOTIFS
10	446	T3 S28 S32 T52 T94 T96 S135 S143 T159 T165 S171 S433		H200 - H222 Y228 - H250 Y256 - H278 Y284 - H306	zinc finger protein ZFP113 (GI 5640017)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
11	428	S72 S92 S101 S118 S120 S125 T245 T277 S289 S315 S317 Y326 S409			Skeletal muscle BOP2 (GI 5870834; SEQ ID NO:117)	BLAST, MOTIFS

TABLE 2 (cont.)

Polypeptide Seq ID No:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
12	590	S45 Y52 T60 S83 S90 T95 T116 T145 T233 T330 S391 S410 S411 T420 T439 T490 S521			Methyl-CpG binding protein (GI 2239126)	BLAST, MOTIFS
13	479	S15 Y29 S30 S118 T173 T183 S203 S217 Y232 S235 T255 S352 S362 Y451		Y232 - H254 H283 - H305 Y311 - H333 Y339 - H361 PFAM	SRE-ZBP (GI g36603)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
14	433	S92 S96 T250 S319 T322 T327 S335 T344		C380 - C421 PFAM	C3HC4-type zinc finger protein	PFAM, BLOCKS, MOTIFS
15	320	T6 S27 T125 T172 S229 S232 T239 S248 S259 S266 Y267 S291			Zinc finger factor (GI 3150148)	BLAST, MOTIFS
16	179	S11 T21 S46 S140			Single-stranded DNA binding protein (csdp) (GI 1562534)	BLAST, MOTIFS
17	494	T73 Y80 S104 Y116 T192 S289 S297 T329 T364 T376 S387		C13 - H41 BLOCKS	Zinc finger transcription factor (GI 2895870)	BLAST, BLOCKS, MOTIFS
18	401	S4 S82 S97 T166 S188 S249 S279 S289 S290 S294 S319 S368 S371 S372 S378 T392 S396		P5 - K81 A114 - S179 G186 - P262 PFAM	HP1-BP74 (GI 1480112)	PFAM, BLOCKS, PRINTS, MOTIFS

TABLE 2 (cont.)

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
19	264	S11 S25 S76 S82 S90 S92 S96 S119 T229		F154 - H176 C180 - H202 F208 - H230 Y236 - C259 PFAM	C2H2-type zinc finger protein (GI 429188)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
20	153	T23 S40 T44 S110 S120 T124		R42 - E141 PFAM	High mobility group-like nuclear protein (GI 2822179)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
21	243	S20 S21 S76 S100 S104 S160 T194 S196 S212 T222 Y229		Y90 - H112 H118 - H140 Y146 - H168 Y174 - H195 PFAM	C2H2-type zinc finger protein (GI 38015)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
22	485	T29 S34 S104 S147 T162 T248 S249 S256 S347 S452 S477		S309 - H331 H337 - H359 Y365 - H387 Y393 - H415 PFAM	BTB domain/C2H2-type zinc finger protein (GI 2843171)	BLAST, PFAM, BLOCKS, PRINTS, MOTIFS
23	160	S118		C5 - F62 C80 - F137 PFAM	LIM domain protein/CRP2 (GI 487284)	BLAST, PFAM, BLOCKS, PROFILES, PRINTS, MOTIFS
24	511	S10 T36 S75 S90 S222 T245 T259 S399 S405 Y443 S500		Y171 - P223 Y267 - K294 BLOCKS	2'-5'oligoadenylate synthetase-related protein p56 (GI 4731857)	BLOCKS, MOTIFS
25	310	S24 S39 T69 Y104 S185 T282 T296			SIR2 family transcriptional regulatory protein (GI 2648874)	BLAST, MOTIFS

TABLE 2 (cont.)

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
26	331	T29 S40 S74 S257 T270 S301		V112 - R134 BLOCKS	Histone protein	BLOCKS, MOTIFS
27	200	T43 S123 T129 S167 S183 S184			Zinc finger protein (g1373394)	MOTIFS BLAST
28	100	S44 Y25 Y98		Transcription anti-terminator; bglG family; E47-I100	transcription elongation factor (g4336506)	MOTIFS BLAST BLOCKS
29	528	S204 T487 S29 S34 T48 T227 S327 T367 T423 S483 Y39 Y44 Y112 Y163	N24 N52 N100 N481	C2H2 zinc fingers: Y191-H213 Y247-H269 Y275-H297 Y331-H353 Y387-H409 Y415-H437 Y443-H465 Y471-H493	Zinc finger protein (g498721)	MOTIFS BLAST BLOCKS PFAM
30	350	T264 S305	N33 N79	C3HC4 RING finger: C230-C271	C3HC4/RING zinc finger protein (g1321818)	MOTIFS BLAST BLOCKS PFAM PROFILESCAN
31	315	S51 T94 S121 S123 S142 S143 T184 S232 S252 T36 T46 S159 S163 S168			Similar to CCAAT/enhancer-binding protein (g1947129)	MOTIFS BLAST
32	120	S36 S56 T93 S104	N98 N103	bZIP transcription factor: P21-P85		MOTIFS PFAM BLOCKS

TABLE 2 (cont.)

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
33	326	S59 S38 T207 S284 T319 T43 S80 T137 T155 T211 S238 T239	N209	C2H2 zinc fingers: C143-C171 Y169-H191 F197-H219	Zinc finger protein (g220643)	MOTIFS BLAST PFAM BLOCKS
34	106	S80 T89 T39 T53		Homeobox: R14-K70	CHOX M product; homeobox protein (g62701)	MOTIFS BLAST BLOCKS PFAM PROFILES SCAN
35	209	S176 T180 S184 T193 S201 S4 T25 S49	N18	bZIP transcription factor: K115-E140	geminin (g3219357)	MOTIFS BLAST PFAM
36	212	S79 S107 T127 T202 S45 S56 S124 T152 Y35	N92	HMG box: M1-Q36	Smardcel-related protein (GI 4321968)	MOTIFS BLAST PFAM
37	359	T329 T50 S125 S224 S230 S235 S344 S31 S215 S312 Y42	N45 N340	C2H2 zinc fingers: F278-C306 Y304-H328 F334-H356	BKLF; CACCC-box binding protein (g1244515)	MOTIFS BLAST BLOCKS PFAM
38	445	S68 T87 S153 S339 S405 S55 T105 S315 S422 Y419	N337 N374 N388	C3HC4 RING fingers: C74-P120 N228-C235	ARI (RING finger) protein (g2058299)	MOTIFS BLAST BLOCKS PFAM
39	433	S283 T44 T57 T123 S136 T185 T220 S239 T268 S313 S330 T105 T109 S125 T216	N169 N206		skm-BOP2 zinc finger protein (g1809327)	MOTIFS BLAST

TABLE 2 (cont.)

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
40	355	T72 T84 T184 S191 T244 S88 T162 T229 T294 S330	N308 N324		Sir2 family protein (g5353746)	MOTIFS BLAST
41	443	T76 T106 S148 T309 S12 T253 S299 S357 T373 T427 Y61 Y114	N366 N425	Myb-like DNA binding domain: D72-F118	ADA2 transcriptional adaptor protein (g170991)	MOTIFS BLAST BLOCKS PFAM PROFLESCAN
42	164	S37 T55 T64 S83 S22 Y146	N117	mutT domain: V29-L70		MOTIFS BLOCKS PFAM
43	215	S79 S127 T12 T45		HMG box: M1-Q36	Sry-related protein (g211510)	MOTIFS BLAST PFAM
44	539	T15 S24 S66 S83 T97 T105 T109 S128 T149 S153 S198 T203 T225 T238 T296 S466 Y258	N465	C2H2 zinc fingers: H230-H252 Y258-H280 Y286-H308 Y314-H336 Y342-H364 F370-H392 Y398-H420 Y426-H448 Y482-H504 Y510-H532	C2H2 zinc finger protein (g5757625)	MOTIFS BLAST PFAM BLOCKS
45	182	T59 S112 S120 S100 S139 Y64			Transcriptional regulator (g2621798)	MOTIFS BLAST

TABLE 2 (cont.)

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
46	534	S494 S31 S44 S117 T123 S185 S216 S476 S504 S176 S182 S211 T249 S293 S323 S409 T489 Y76 Y285	N29 N39 N250 N351	C2H2 zinc fingers: Y285-H307 Y313-H335 Y341-H363 C369-H391 Y397-H419 Y425-H447 Y453-H475 Y481-H503 Y509-H531	Zinc finger protein (g1373394)	MOTIFS BLAST PFAM BLOCKS
47	206	S5 S7 S40 S45 S46 S100 S144 S26 S107 T148 S185 Y38	N44 N177	Myc-type HLH domain: Q108-R160	Musculin (g3599519)	MOTIFS BLAST PFAM BLOCKS PROFILES CAN
48	172	T5 S87 S96 S115 T124 S22 T64			KRAB zinc finger protein (g1049295)	MOTIFS BLAST
49	275	S185 S14 S48 T54 S118 T139 T161 T189 T217 Y256	N210 N214 N238 N260	C2H2 zinc fingers: F172-H194 Y200-H222 Y228-H250	Repressor transcriptional factor (g1017722)	MOTIFS BLAST PFAM BLOCKS
50	236	S157 S42 T167 T222 T81 Y213	N40	C3HC4 RING finger: P126-L150	Ariadne-2 RING finger protein (g3445441)	MOTIFS BLAST BLOCKS
51	214	S7 S8 S116 T127 S154 S191 T31 S41 T204	N2	Chromodomain: V113-E134	Nucleoplasmin (g833629)	MOTIFS BLAST BLOCKS

TABLE 2 (cont.)

Polypeptide Seq ID NO:	Amino Acid Residues	Potential Phosphorylation Sites	Potential Glycosylation Sites	Signature Sequence(s)	Identification/ Homologous Sequence	Analytical Methods and Databases
52	396	T348 T392 S118 T193 T201 S270 S294 S80 S112 S206 S260 T313 T355 S375 S387	N2	C3HC4 RING finger: C26-C50	Midline 1/ cerebellar isoform 1 RING finger protein (g3462503)	MOTIFS BLAST PFAM BLOCKS PROFESCAN
53	486	S29 T58 S155 S239 T292 T379 S146 T271 S425			5'-nucleotidase (g633071)	MOTIFS BLAST
54	555	S432 T502 S68 S195 T199 T226 S315 T379 T441 T534 S170 S248 S282 S291 T327 T336 S391 S422 T481 Y257 Y274	N25 N66 N246 N364	ATP/GTP binding site (P-loop): A434-T441	Transcription termination factor I (TTF-I) interacting peptide 5 isoform (g2183083)	MOTIFS BLAST
55	61	T34 T42 Y48			Putative leucine-rich DNA-binding protein (g555991)	MOTIFS BLAST

TABLE 3

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
56	169-215	Reproductive (0.224) Nervous (0.198) Cardiovascular (0.112)	Cell Proliferative (0.725) Inflammation (0.190)	PBLUESCRIPT
57	551-595	Hematopoietic/Immune (0.240) Reproductive (0.180) Gastrointestinal (0.120)	Cell Proliferative (0.700) Inflammation (0.360)	PBLUESCRIPT
58	541-585	Nervous (0.286) Reproductive (0.286) Cardiovascular (0.214)	Cell Proliferative (0.643) Trauma (0.214)	PBLUESCRIPT
59	109-153	Reproductive (1.000)	Cell Proliferative (1.000) Inflammation (1.000)	PBLUESCRIPT
60	435-479	Hematopoietic/Immune (0.211) Gastrointestinal (0.183) Reproductive (0.183)	Cell Proliferative (0.620) Inflammation (0.338)	PBLUESCRIPT
61	1195-1239	Reproductive (0.248) Cardiovascular (0.174) Nervous (0.157)	Cell Proliferative (0.637) Inflammation (0.256)	PBLUESCRIPT
62	217-261	Reproductive (0.429) Nervous (0.238) Cardiovascular (0.095)	Cell Proliferative (0.667) Inflammation (0.143) Trauma (0.095)	PBLUESCRIPT
63	919-963	Reproductive (0.265) Nervous (0.235) Cardiovascular (0.088)	Cell Proliferative (0.618) Inflammation (0.206)	PSPORT1
64	823-876	Reproductive (0.382) Nervous (0.176) Gastrointestinal (0.118)	Cell Proliferative (0.794)	PSPORT1
65	380-424	Reproductive (0.346) Nervous (0.154) Gastrointestinal (0.135)	Cell Proliferative (0.750) Inflammation (0.231)	PSPORT1

TABLE 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
66	757-801	Nervous (0.222) Hematopoietic/Immune (0.167) Reproductive (0.167)	Cell Proliferative (0.778) Inflammation (0.222)	pINCY
67	812-856	Reproductive (0.246) Nervous (0.180) Gastrointestinal (0.148)	Cell Proliferative (0.639) Inflammation (0.246)	pINCY
68	326-370	Reproductive (0.500) Nervous (0.200) Gastrointestinal (0.150)	Cell Proliferative (0.700) Trauma (0.150)	pINCY
69	703-747	Reproductive (0.278) Hematopoietic/Immune (0.204) Nervous (0.148)	Cell Proliferative (0.777) Inflammation (0.222)	pINCY
70	759-803	Reproductive (0.261) Gastrointestinal (0.217) Nervous (0.174)	Cell Proliferative (0.565) Trauma (0.130)	pINCY
71	110-154	Nervous (0.250) Developmental (0.208) Gastrointestinal (0.167)	Cell Proliferative (0.583) Trauma (0.167)	pINCY
72	529-573	Reproductive (0.186) Gastrointestinal (0.168) Hematopoietic/Immune (0.138)	Cell Proliferative (0.700) Inflammation (0.251)	pINCY
73	1784-1828	Reproductive (0.286) Hematopoietic/Immune (0.190) Nervous (0.167)	Cell Proliferative (0.667) Inflammation (0.286)	PSPORT1
74	111-155	Reproductive (0.316) Nervous (0.211) Hematopoietic/Immune (0.158)	Cell Proliferative (0.632) Inflammation (0.211)	PSPORT1
75	543-587	Reproductive (0.258) Nervous (0.206) Gastrointestinal (0.134)	Cell Proliferative (0.608) Inflammation (0.196)	pINCY

TABLE 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
76	272-316	Reproductive (0.246) Nervous (0.180) Hematopoietic/Immune (0.148)	Cell Proliferative (0.606) Inflammation (0.279)	pINCY
77	227-271	Hematopoietic/Immune (0.222) Endocrine (0.167) Cardiovascular (0.111)	Cell Proliferative (0.666) Inflammation (0.333)	pINCY
78	487-531	Gastrointestinal (0.375) Reproductive (0.250)	Cell Proliferative (0.500) Inflammation (0.250)	pINCY
79	111-155	Gastrointestinal (0.280) Hematopoietic/Immune (0.240) Reproductive (0.120)	Cell Proliferative (0.640) Inflammation (0.440)	pINCY
80	595-639	Reproductive (0.211) Gastrointestinal (0.158) Urologic (0.158)	Cell Proliferative (0.684) Inflammation (0.263)	pINCY
81	425-469	Reproductive (0.222) Gastrointestinal (0.160) Nervous (0.148)	Cell Proliferative (0.568) Inflammation (0.259)	PSPORT1
82	774-818	Gastrointestinal (0.200) Hematopoietic/Immune (0.200) Nervous (0.200)	Cancer (0.600) Trauma (0.200) Inflammation (0.200)	PBLUESCRIPT
83	517-561	Nervous (0.526) Reproductive (0.132) Cardiovascular (0.105)	Cancer (0.342) Fetal (0.158) Inflammation (0.158)	PBLUESCRIPT
84	1944-1988	Nervous (0.250) Reproductive (0.188) Endocrine (0.125)	Cancer (0.438) Fetal (0.250) Trauma (0.250)	PSPORT1
85	1027-1071	Reproductive (0.219) Nervous (0.206) Hematopoietic/Immune (0.116)	Cancer (0.458) Inflammation (0.232) Fetal (0.181)	PSPORT1

TABLE 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
86	658-702	Reproductive (0.227) Hematopoietic/Immune (0.182) Gastrointestinal (0.167)	Cancer (0.424) Inflammation (0.318) Fetal (0.167)	PT7T3
87	488-532	Nervous (0.200) Reproductive (0.200) Musculoskeletal (0.120)	Cancer (0.320) Fetal (0.320) Inflammation (0.320)	pINCY
88	379-423	Cardiovascular (0.250) Nervous (0.250) Reproductive (0.250)	Cancer (0.417) Fetal (0.167) Neurological (0.167)	pINCY
89	632-676	Reproductive (0.417) Cardiovascular (0.167) Gastrointestinal (0.167)	Cancer (0.333) Fetal (0.167) Inflammation (0.167)	pINCY
90	258-302	Reproductive (0.294) Nervous (0.137) Hematopoietic/Immune (0.118)	Cancer (0.569) Fetal (0.431) Inflammation (0.176)	pINCY
91	433-477	Reproductive (0.750) Nervous (0.250)	Cancer (0.500) Inflammation (0.500)	PSPORT1
92	542-586	Gastrointestinal (0.273) Hematopoietic/Immune (0.273) Developmental (0.182)	Cancer (0.455) Inflammation (0.364) Fetal (0.182)	pINCY
93	218-262	Reproductive (0.272) Nervous (0.204) Cardiovascular (0.126)	Cancer (0.447) Inflammation (0.214) Fetal (0.155)	PSPORT1
94	541-585	Reproductive (0.273) Nervous (0.250) Cardiovascular (0.159)	Cancer (0.364) Fetal (0.205) Inflammation (0.205)	PSPORT1
95	111-155	Reproductive (0.250) Gastrointestinal (0.173) Nervous (0.154)	Cancer (0.481) Fetal (0.231) Inflammation (0.212)	PSPORT1
96	597-641	Reproductive (0.261) Cardiovascular (0.217) Nervous (0.130)	Cancer (0.391) Fetal (0.304) Inflammation (0.130)	PSPORT1

TABLE 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
97	434-478	Cardiovascular (0.222) Endocrine (0.222) Gastrointestinal (0.222)	Cancer (0.667) Fetal (0.111) Neurological (0.111)	PSPORT1
98	218-247 920-964	Reproductive (0.333) Gastrointestinal (0.129) Hematopoietic/Immune (0.118)	Cancer (0.559) Inflammation (0.204) Fetal (0.183)	pINCY
99	327-371	Gastrointestinal (0.211) Reproductive (0.211) Cardiovascular (0.158)	Cancer (0.421) Fetal (0.316) Inflammation (0.158)	pINCY
100	596-625	Reproductive (0.230) Nervous (0.164) Gastrointestinal (0.131)	Cancer (0.590) Inflammation (0.246) Fetal (0.082)	PSPORT
101	487-531	Cardiovascular (0.235) Reproductive (0.235) Hematopoietic/Immune (0.176)	Cancer (0.588) Inflammation (0.176) Trauma (0.118)	pINCY
102	218-247 542-586	Gastrointestinal (0.241) Hematopoietic/Immune (0.207) Cardiovascular (0.138)	Cancer (0.448) Fetal (0.276) Inflammation (0.276)	pINCY
103	219-263	Reproductive (0.500) Cardiovascular (0.250) Hematopoietic/Immune (0.250)	Cancer (0.500) Inflammation (0.250) Trauma (0.250)	pINCY
104	111-140 327-371	Hematopoietic/Immune (0.286) Nervous (0.238) Reproductive (0.143)	Cancer (0.333) Fetal (0.286) Inflammation (0.286)	pINCY
105	243-281	Musculoskeletal (0.286) Nervous (0.286) Gastrointestinal (0.143)	Inflammation (0.429) Fetal (0.286) Cancer (0.286)	pINCY
106	271-315	Nervous (0.800) Reproductive (0.100) Cardiovascular (0.100)	Cancer (0.400) Inflammation (0.200) Trauma (0.200)	pINCY

TABLE 3 (cont.)

Nucleotide Seq ID NO:	Selected Fragment	Tissue Expression (Fraction of Total)	Disease or Condition (Fraction of Total)	Vector
107	489-533	Cardiovascular (0.364) Gastrointestinal (0.182) Reproductive (0.182)	Cancer (0.273) Trauma (0.273) Inflammation (0.182)	pINCY
108	156-200	Nervous (0.256) Reproductive (0.256) Hematopoietic/Immune (0.128)	Cancer (0.465) Fetal (0.291) Inflammation (0.186)	pINCY
109	1459-1503	Cardiovascular (0.250) Hematopoietic/Immune (0.250) Nervous (0.167)	Inflammation (0.417) Cancer (0.333) Trauma (0.167)	PSPORT1
110	164-208	Nervous (1.000)	Neurological (1.000)	pINCY

TABLE 4

Nucleotide SEQ ID NO:	Library	Library Comment
56	SPLNFETO 1	Library was constructed at Stratagene, using RNA isolated from a pool of fetal spleen tissue. Following vector packaging, 2x10 ⁶ primary clones were then amplified to stabilize the library for long-term storage. Amplification may significantly skew sequence abundances.
57	SYNORABO 1	Library was constructed using RNA isolated from the synovial membrane tissue of a 68-year-old Caucasian female with rheumatoid arthritis.
58	KIDNNOTO 1	Library was constructed using RNA isolated from the kidney tissue of a 64-year-old Caucasian female, who died from an intracranial bleed. Patient history included rheumatoid arthritis and tobacco use.
59	PLACNOBO 1	Library was constructed using RNA isolated from placenta.
60	HNT2RATO 1	Library was constructed at Stratagene (STR937231), using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated with retinoic acid for 24 hours.
61	EOSIHETO 2	Library was constructed using RNA isolated from peripheral blood cells apheresed from a 48-year-old Caucasian male. Patient history included hypereosinophilia. The cell population was determined to be greater than 77% eosinophils by Wright's staining.
62	HNT2AGTO 1	Library was constructed at Stratagene (STR937233), using RNA isolated from the hNT2 cell line (derived from a human teratocarcinoma that exhibited properties characteristic of a committed neuronal precursor). Cells were treated with retinoic acid for 5 weeks and with mitotic inhibitors for two weeks and allowed to mature for an additional 4 weeks in conditioned medium.
63	PROSNOTO 2	Library was constructed using RNA isolated from the diseased prostate tissue removed from a 50-year-old Caucasian male during a retropubic prostatectomy. Pathology indicated adenofibromatous hyperplasia was present. Pathology for the associated tumor tissue indicated adenocarcinoma Gleason grade 3+3. Patient history included dysuria, carcinoma in situ of prostate, coronary atherosclerosis, and hyperlipidemia.
64	COLNTUTO 2	Library was constructed using RNA isolated from colon tumor tissue removed from a 75-year-old Caucasian male during a hemicolectomy. Pathology indicated invasive grade 3 adenocarcinoma arising in a tubulovillous adenoma, which was distal to the ileocecal valve in the cecum. The tumor penetrated deeply into the muscularis propria but not through it.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
65	KIDNTUT0 1	Library was constructed using RNA isolated from the kidney tumor tissue removed from an 8-month-old female during nephroureterectomy. Pathology indicated Wilms' tumor (nephroblastoma), which involved 90 percent of the renal parenchyma. Prior to surgery, the patient was receiving heparin anticoagulant therapy.
66	THYRNOT0 3	Library was constructed using RNA isolated from thyroid tissue removed from the left thyroid of a 28-year-old Caucasian female during a complete thyroidectomy. Pathology indicated a small nodule of adenomatous hyperplasia present in the left thyroid. Pathology for the associated tumor tissue indicated dominant follicular adenoma, forming a well-encapsulated mass in the left thyroid.
67	BLADTUT0 4	Library was constructed using RNA isolated from bladder tumor tissue removed from a 60-year-old Caucasian male during a radical cystectomy, prostatectomy, and vasectomy. Pathology indicated grade 3 transitional cell carcinoma in the left bladder wall. Carcinoma in-situ was identified in the dome and trigone. Patient history included tobacco use. Family history included type I diabetes, a malignant neoplasm of the stomach, atherosclerotic coronary artery disease, and an acute myocardial infarction.
68	PROSNOT1 5	Library was constructed using RNA isolated from diseased prostate tissue removed from a 66-year-old Caucasian male during radical prostatectomy and regional lymph node excision. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 2+3). The patient presented with elevated prostate specific antigen (PSA). Family history included prostate cancer, secondary bone cancer, and benign hypertension.
69	BRSTTUT0 8	Library was constructed using RNA isolated from breast tumor tissue removed from a 45-year-old Caucasian female during unilateral extended simple mastectomy. Pathology indicated invasive nuclear grade 2-3 adenocarcinoma, ductal type, with 3 of 23 lymph nodes positive for metastatic disease. Greater than 50% of the tumor volume was in situ, both comedo and non-comedo types. Immunostains were positive for estrogen/progesterone receptors, and uninvolved tissue showed proliferative changes. The patient concurrently underwent a total abdominal hysterectomy. Patient history included valvuloplasty of mitral valve without replacement, rheumatic mitral insufficiency, and rheumatic heart disease. Family history included acute myocardial infarction, atherosclerotic coronary artery disease, and type II diabetes.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
70	PROSNOT1 9	Library was constructed using RNA isolated from diseased prostate tissue removed from a 59-year-old Caucasian male during a radical prostatectomy with regional lymph node excision. Pathology indicated adenofibromatous hyperplasia. Pathology for the associated tumor tissue indicated an adenocarcinoma (Gleason grade 3+3). The patient presented with elevated prostate-specific antigen (PSA). Patient history included colon diverticuli, asbestosis, and thrombophlebitis. Previous surgeries included a partial colectomy. Family history included benign hypertension, multiple myeloma, hyperlipidemia and rheumatoid arthritis.
71	THP1NOT0 3	Library was constructed using RNA isolated from untreated THP-1 cells. THP-1 (ATCC TIB 202) is a human promonocyte line derived from the peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia (ref: Int. J. Cancer (1980) 26:171).
72	THP1NOT0 3	Library was constructed using RNA isolated from untreated THP-1 cells. THP-1 (ATCC TIB 202) is a human promonocyte line derived from the peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia (ref: Int. J. Cancer (1980) 26:171).
73	HIPOAZT0 1	Library was constructed from RNA isolated from diseased hippocampus tissue removed from the brain of a 74-year-old Caucasian male who died from Alzheimer's disease.
74	HIPOAZT0 1	Library was constructed from RNA isolated from diseased hippocampus tissue removed from the brain of a 74-year-old Caucasian male who died from Alzheimer's disease.
75	BRSTNOT1 3	Library was constructed using RNA isolated from breast tissue removed from the left medial lateral breast of a 36-year-old Caucasian female during bilateral simple mastectomy and total breast reconstruction. Pathology indicated benign breast tissue. Patient history included a breast neoplasm, depressive disorder, hyperlipidemia, chronic stomach ulcer, and an ectopic pregnancy. Family history included myocardial infarction, cerebrovascular disease, atherosclerotic coronary artery disease, hyperlipidemia, skin cancer, breast cancer, depressive disorder, esophageal cancer, bone cancer, Hodgkin's lymphoma, bladder cancer, and heart condition.
76	NPOLNOT0 1	Library was constructed using RNA isolated from nasal polyp tissue removed from a 78-year-old Caucasian male during a nasal polypectomy. Pathology indicated a nasal polyp and striking eosinophilia. Patient history included asthma and nasal polyps.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
77	ADRENOT0 9	Library was constructed using RNA isolated from left adrenal gland tissue removed from a 43-year-old Caucasian male during nephroureterectomy, regional lymph node excision, and unilateral left adrenalectomy. Pathology indicated no diagnostic abnormalities of the adrenal gland. Pathology for the associated tumor tissue indicated a grade 2 renal cell carcinoma mass in the posterior lower pole of the left kidney with invasion into the renal pelvis.
78	SPLNNOT1 1	Library was constructed using RNA isolated from diseased spleen tissue removed from a 14-year-old Asian male during a total splenectomy. Pathology indicated changes consistent with idiopathic thrombocytopenic purpura. The patient presented with bruising.
79	CONCNOT0 1	Library was constructed using RNA isolated from chest wall soft tissue removed from a 63-year-old Caucasian male during a chest wall lesion destruction. Pathology indicated surgical margins were free of tumor. Pathology for the associated tumor tissue indicated invasive grade 3 adenocarcinoma, forming a mass that extended through the visceral pleura to involve parietal pleura. Patient history included MEN (multiple endocrine neoplasia) syndrome type I, abnormal secretion of gastrin, alcohol and tobacco abuse, calcium metabolism disease, chronic stomach ulcer with hemorrhage, lung cancer, and calculus of the kidney. Family history included prostate cancer, benign hypertension, stroke, atherosclerotic coronary artery disease, type II diabetes, hyperlipidemia, and cancer of an unspecified location.
80	TONSNOT0 3	Library was constructed using RNA isolated from diseased left tonsil tissue removed from a 6-year-old Caucasian male during adenotonsillectomy. Pathology indicated reactive lymphoid hyperplasia, bilaterally. Family history included benign hypertension, myocardial infarction, and atherosclerotic coronary artery disease.
81	LUNGNO0 3	This normalized library was constructed from 2.56×10^6 independent clones from a lung tissue library. RNA was made from lung tissue removed from the left lobe a 58-year-old Caucasian male during a segmental lung resection. Pathology for the associated tumor tissue indicated a metastatic grade 3 (of 4) osteosarcoma. Patient history included soft tissue cancer, secondary cancer of the lung, prostate cancer, and an acute duodenal ulcer with hemorrhage. Patient also received radiation therapy to the retroperitoneum. Family history included prostate cancer, breast cancer, and acute leukemia. The normalization and hybridization conditions were adapted from Soares et al., PNAS (1994) 91:9228; Swaroop et al., NAR (1991) 19:1954; and Bonaldo et al., Genome Research (1996) 6:791.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
82	BMARNOT0 2	The library was constructed using Clontech RNA isolated from the bone marrow of 24 male and female Caucasian donors, 16 to 70 years old.
83	SINTNOT0 2	The library was constructed using RNA isolated from the small intestine of a 55-year-old Caucasian female, who died from a subarachnoid hemorrhage. Serologies were positive for cytomegalovirus (CMV).
84	CRBLNOT0 1	The library was constructed using RNA isolated from the cerebellum tissue of a 69-year-old Caucasian male who died from chronic obstructive pulmonary disease. Patient history included myocardial infarction, hypertension, and osteoarthritis.
85	BRSTNOT0 3	The library was constructed using RNA isolated from diseased breast tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy. Pathology for the associated tumor tissue indicated residual invasive grade 3 mammary ductal adenocarcinoma. Patient history included kidney infection and condyloma acuminatum. Family history included benign hypertension, hyperlipidemia and a malignant neoplasm of the colon.
86	BEPINON0 1	The normalized bronchial epithelium library was constructed from 5.12 million independent clones from a bronchial epithelium library. RNA was isolated from a bronchial epithelium primary cell line derived from a 54-year-old Caucasian male. The normalization and hybridization conditions were adapted from Soares et al., PNAS (1994) 91:9228, using a 24-hour reannealing hybridization period.
87	THYRNOT0 3	The library was constructed using RNA isolated from thyroid tissue removed from the left thyroid of a 28-year-old Caucasian female during a complete thyroidectomy. Pathology indicated a small nodule of adenomatous hyperplasia present in the left thyroid. Pathology for the associated tumor tissue indicated dominant follicular adenoma, forming a well-encapsulated mass in the left thyroid.
88	COLNFETO 2	The library was constructed using RNA isolated from the colon tissue of a Caucasian female fetus who died at 20 weeks' gestation.
89	BLADNOT0 3	The library was constructed using RNA isolated from bladder tissue removed from an 80-year-old Caucasian female during a radical cystectomy and lymph node excision. Pathology for the associated tumor tissue indicated grade 3 invasive transitional cell carcinoma. Patient history included malignant neoplasm of the uterus, atherosclerosis, and atrial fibrillation. Family history included acute renal failure, osteoarthritis, and atherosclerosis.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
90	COLNNOT2 3	The library was constructed using RNA isolated from diseased colon tissue removed from a 16-year-old Caucasian male during a total colectomy with abdominal/perineal resection. Pathology indicated gastritis and pancolitis consistent with the acute phase of ulcerative colitis. Inflammation was more severe in the transverse colon with inflammation confined to the mucosa. The ascending and sigmoid colon was mildly involved. Family history included irritable bowel syndrome.
91	BRSTNOT0 4	The library was constructed using RNA isolated from breast tissue removed from a 62-year-old East Indian female during a unilateral extended simple mastectomy. Pathology for the associated tumor tissue indicated an invasive grade 3 ductal carcinoma. Patient history included benign hypertension, hyperlipidemia, and hematuria. Family history included cerebrovascular and cardiovascular disease, hyperlipidemia, and liver cancer.
92	LIVRFETO 2	The library was constructed using RNA isolated from liver tissue removed from a Caucasian female fetus who died at 20 weeks' gestation. Family history included seven days of erythromycin treatment for bronchitis in the mother during the first trimester.
93	BEPINOT0 1	The library was constructed using RNA isolated from a bronchial epithelium primary cell line derived from a 54-year-old Caucasian male.
94	BRAITUTO 2	The library was constructed using RNA isolated from brain tumor tissue removed from the frontal lobe of a 58-year-old Caucasian male during excision of a cerebral meningeal lesion. Pathology indicated a grade 2 metastatic hypernephroma. Patient history included a grade 2 renal cell carcinoma, insomnia, and chronic airway obstruction. Family history included a malignant neoplasm of the kidney.
95	BRAITUTO 3	The library was constructed using RNA isolated from brain tumor tissue removed from the left frontal lobe of a 17-year-old Caucasian female during excision of a cerebral meningeal lesion. Pathology indicated a grade 4 fibrillary giant and small-cell astrocytoma. Family history included benign hypertension and cerebrovascular disease.
96	BRSTTUTO 2	The library was constructed using RNA isolated from breast tumor tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy with reconstruction. Pathology indicated residual invasive grade 3 mammary ductal adenocarcinoma. The remaining breast parenchyma exhibited proliferative fibrocystic changes without atypia. One of 10 axillary lymph nodes had metastatic tumor. Patient history included kidney infection and condyloma acuminatum. Family history included benign hypertension, hyperlipidemia, and a malignant colon neoplasm.

TABLE 4 (cont.)

Nucleotide SEQ ID NO:	Library	Library Comment
97	BRSTTUT0 2	The library was constructed using RNA isolated from breast tumor tissue removed from a 54-year-old Caucasian female during a bilateral radical mastectomy with reconstruction. Pathology indicated residual invasive grade 3 mammary ductal adenocarcinoma. The remaining breast parenchyma exhibited proliferative fibrocystic changes without atypia. One of 10 axillary lymph nodes had metastatic tumor. Patient history included kidney infection and condyloma acuminatum. Family history included benign hypertension, hyperlipidemia, and a malignant colon neoplasm.
98	BRSTNOT0 7	The library was constructed using RNA isolated from diseased breast tissue removed from a 43-year-old Caucasian female during a unilateral extended simple mastectomy. Pathology indicated mildly proliferative fibrocystic changes with epithelial hyperplasia, papillomatosis, and duct ectasia. Pathology for the associated tumor tissue indicated invasive grade 4, nuclear grade 3 mammary adenocarcinoma with extensive comedo necrosis. Family history included epilepsy, cardiovascular disease, and type II diabetes.
99	BRAINOT0 9	The library was constructed using RNA isolated from brain tissue removed from a Caucasian male fetus who died at 23 weeks' gestation.
100	THP1AZS0 8	The library was constructed using RNA isolated from 5.76 million clones from a 5-aza-2'-deoxycytidine treated THP-1 cell library. The library was subjected to subtractive hybridization using 5 million clones from an untreated THP-1 cell library. Hybridization conditions were adapted from Swaroop et al., NAR (1991) 19:1954; and Bonaldo et al., Genome Research (1996) 6:791. THP-1 (ATCC TIB 202) is a human promonocyte cell line derived from peripheral blood of a 1-year-old Caucasian male with acute monocytic leukemia.
101	LUNGUTUT1 2	The library was constructed using RNA isolated from tumorous lung tissue removed from a 70-year-old Caucasian female during a lung lobectomy of the left upper lobe. Pathology indicated grade 3 (of 4) adenocarcinoma and vascular invasion. Patient history included tobacco abuse, depressive disorder, anxiety state, and skin cancer. Family history included cerebrovascular disease, congestive heart failure, colon cancer, depressive disorder, and primary liver.
102	HEAONOT0 5	The library was constructed using RNA isolated from aortic tissue removed from a 17-year-old Hispanic female who died from a gunshot wound.
103	HEAONOT0 5	The library was constructed using RNA isolated from aortic tissue removed from a 17-year-old Hispanic female who died from a gunshot wound.

Table 5

Program	Description	Reference	Parameter Threshold
ABI FACTURA	A program that removes vector sequences and masks ambiguous bases in nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
ABI/PARACEL FDF	A Fast Data Finder useful in comparing and annotating amino acid or nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA; Paracel Inc., Pasadena, CA.	Mismatch <50%
ABI AutoAssembler	A program that assembles nucleic acid sequences.	Perkin-Elmer Applied Biosystems, Foster City, CA.	
BLAST	A Basic Local Alignment Search Tool useful in sequence similarity search for amino acid and nucleic acid sequences. BLAST includes five functions: blastp, blastn, blastx, tblastn, and tblastx.	Altschul, S.F. et al. (1990) <i>J. Mol. Biol.</i> 215:403-410; Altschul, S.F. et al. (1997) <i>Nucleic Acids Res.</i> 25: 3389-3402.	ESTs: Probability value= 1.0E-8 or less Full Length sequences: Probability value= 1.0E-10 or less
FASTA	A Pearson and Lipman algorithm that searches for similarity between a query sequence and a group of sequences of the same type. FASTA comprises at least five functions: fasta, tfasta, fastx, tfastx, and ssearch.	Pearson, W.R. and D.J. Lipman (1988) <i>Proc. Natl. Acad. Sci.</i> 85:2444-2448; Pearson, W.R. (1990) <i>Methods Enzymol.</i> 183: 63-98; and Smith, T.F. and M. S. Waterman (1981) <i>Adv. Appl. Math.</i> 2:482-489.	ESTs: fasta E value=1.06E-6 Assembled ESTs: fasta Identity= 95% or greater and Match length=200 bases or greater. fastx E value= 1.0E-8 or less Full Length sequences: fastx score=100 or greater
BLIMPS	A BLOCKS IMPROVED Searcher that matches a sequence against those in BLOCKS, PRINTS, DOMO, PRODOM, and PFAM databases to search for gene families, sequence homology, and structural fingerprint regions.	Henikoff, S. and J.G. Henikoff, <i>Nucl. Acid Res.</i> , 19:6565-72, 1991. J.G. Henikoff and S. Henikoff (1996) <i>Methods Enzymol.</i> 266:88-105; and Attwood, T.K. et al. (1997) <i>J. Chem. Inf. Comput. Sci.</i> 37: 417-424.	Score=1000 or greater; Ratio of Score/Strength = 0.75 or larger; and, if applicable, Probability value= 1.0E-3 or less
IIMMER	An algorithm for searching a query sequence against hidden Markov model (HMM)-based databases of protein family consensus sequences, such as PFAM.	Krogh, A. et al. (1994) <i>J. Mol. Biol.</i> , 235:1501-1531; Sonnhammer, E.L.L. et al. (1988) <i>Nucleic Acids Res.</i> 26:320-322.	Score=10-50 bits for PFAM hits, depending on individual protein families

Table 5 (cont.)

Program	Description	Reference	Parameter Threshold
ProfileScan	An algorithm that searches for structural and sequence motifs in protein sequences that match sequence patterns defined in Prosite.	Gribskov, M. et al. (1988) CABIOS 4:61-66; Gribskov, et al. (1989) Methods Enzymol. 183:146-159; Bairoch, A. et al. (1997) Nucleic Acids Res. 25: 217-221.	Normalized quality score>GCG-specified "HIGH" value for that particular Prosite motif. Generally, score=1.4-2.1.
Phred	A base-calling algorithm that examines automated sequencer traces with high sensitivity and probability.	Ewing, B. et al. (1998) Genome Res. 8:175-185; Ewing, B. and P. Green (1998) Genome Res. 8:186-194.	
Phrap	A Phils Revised Assembly Program including SWAT and CrossMatch, programs based on efficient implementation of the Smith-Waterman algorithm, useful in searching sequence homology and assembling DNA sequences.	Smith, T.F. and M. S. Waterman (1981) Adv. Appl. Math. 2:482-489; Smith, T.F. and M. S. Waterman (1981) J. Mol. Biol. 147:195-197; and Green, P., University of Washington, Seattle, WA.	Score= 120 or greater; Match length= 56 or greater
Consed	A graphical tool for viewing and editing Phrap assemblies	Gordon, D. et al. (1998) Genome Res. 8:195-202.	
SPScan	A weight matrix analysis program that scans protein sequences for the presence of secretory signal peptides.	Nielson, H. et al. (1997) Protein Engineering 10:1-6; Claverie, J.M. and S. Audic (1997) CABIOS 12: 431-439.	Score=3.5 or greater
Motifs	A program that searches amino acid sequences for patterns that matched those defined in Prosite.	Bairoch et al. <i>supra</i> ; Wisconsin Package Program Manual, version 9, page M51-59, Genetics Computer Group, Madison, WI.	

What is claimed is:

1. An isolated polypeptide comprising:
 - a) an amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID
5 NO:3-5, SEQ ID NO:7-14, SEQ ID NO:16-31, SEQ ID NO:33-34, SEQ ID NO:36-40, SEQ ID
NO:42-48, SEQ ID NO:50-55.
 - b) a naturally occurring amino acid sequence having at least 90% sequence identity to an
amino acid sequence selected from the group consisting of SEQ ID NO:1, SEQ ID NO:3-5, SEQ ID
NO:7-14, SEQ ID NO:16-31, SEQ ID NO:33-34, SEQ ID NO:36-40, SEQ ID NO:42-48, SEQ ID
10 NO:50-55,
 - c) a biologically active fragment of an amino acid sequence selected from the group
consisting of SEQ ID NO:1, SEQ ID NO:3-5, SEQ ID NO:7-14, SEQ ID NO:16-31, SEQ ID NO:33-
34, SEQ ID NO:36-40, SEQ ID NO:42-48, SEQ ID NO:50-55, or
 - d) an immunogenic fragment of an amino acid sequence selected from the group consisting
15 of SEQ ID NO:1, SEQ ID NO:3-5, SEQ ID NO:7-14, SEQ ID NO:16-31, SEQ ID NO:33-34, SEQ ID
NO:36-40, SEQ ID NO:42-48, SEQ ID NO:50-55.
2. An isolated polypeptide of claim 1, having an amino acid sequence selected from the
group consisting of SEQ ID NO:1, SEQ ID NO:3-5, SEQ ID NO:7-14, SEQ ID NO:16-31, SEQ ID
20 NO:33-34, SEQ ID NO:36-40, SEQ ID NO:42-48, SEQ ID NO:50-55.
3. An isolated polynucleotide encoding a polypeptide of claim 1.
4. An isolated polynucleotide of claim 3, having a sequence selected from the group
25 consisting of SEQ ID NO:56-110.
5. A recombinant polynucleotide comprising a promoter sequence operably linked to a
polynucleotide of claim 3.
- 30 6. A cell transformed with a recombinant polynucleotide of claim 5.
7. A transgenic organism comprising a polynucleotide of claim 5.
8. A method for producing a polypeptide of claim 1, the method comprising:
 - a) culturing a cell under conditions suitable for expression of the polypeptide, wherein said
35

cell is transformed with a recombinant polynucleotide, and said recombinant polynucleotide comprises a promoter sequence operably linked to a polynucleotide encoding the polypeptide of claim 1, and

b) recovering the polypeptide so expressed.

5

9. An isolated antibody which specifically binds to a polypeptide of claim 1.

10. An isolated polynucleotide comprising:

10

- a) a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110,
- b) a naturally occurring polynucleotide sequence having at least 90% sequence identity to a polynucleotide sequence selected from the group consisting of SEQ ID NO:56-110,
- c) a polynucleotide sequence complementary to a), or
- d) a polynucleotide sequence complementary to b).

15

11. An isolated polynucleotide comprising at least 60 contiguous nucleotides of a polynucleotide of claim 10.

12. A method for detecting a target polynucleotide in a sample, said target polynucleotide having a sequence of a polynucleotide of claim 10, the method comprising:

20

a) hybridizing the sample with a probe comprising at least 16 contiguous nucleotides comprising a sequence complementary to said target polynucleotide in the sample, and which probe specifically hybridizes to said target polynucleotide, under conditions whereby a hybridization complex is formed between said probe and said target polynucleotide, and

25

b) detecting the presence or absence of said hybridization complex, and, optionally, if present, the amount thereof.

13. A method of claim 12, wherein the probe comprises at least 30 contiguous nucleotides.

30

14. A method of claim 12, wherein the probe comprises at least 60 contiguous nucleotides.

15. A pharmaceutical composition comprising an effective amount of a polypeptide of claim 1 and a pharmaceutically acceptable excipient.

35

16. A method of treating a disease or condition associated with decreased expression of

functional NuABP, comprising administering to a patient in need of such treatment the pharmaceutical composition of claim 15.

17. A method for screening a compound for effectiveness as an agonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting agonist activity in the sample.

18. A pharmaceutical composition comprising an agonist compound identified by a method of claim 17 and a pharmaceutically acceptable excipient.

19. A method of treating a disease or condition associated with decreased expression of functional NuABP, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 18.

15

20. A method for screening a compound for effectiveness as an antagonist of a polypeptide of claim 1, the method comprising:

- a) exposing a sample comprising a polypeptide of claim 1 to a compound, and
- b) detecting antagonist activity in the sample.

20

21. A pharmaceutical composition comprising an antagonist compound identified by a method of claim 20 and a pharmaceutically acceptable excipient.

22. A method for treating a disease or condition associated with overexpression of functional NuABP, comprising administering to a patient in need of such treatment a pharmaceutical composition of claim 21.

25

23. A method for screening a compound for effectiveness in altering expression of a target polynucleotide, wherein said target polynucleotide comprises a sequence of claim 4, the method comprising:

30

- a) exposing a sample comprising the target polynucleotide to a compound, and
- b) detecting altered expression of the target polynucleotide.

SEQUENCE LISTING

<110> INCYTE PHARMACEUTICALS, INC.

TANG, Y. Tom

LAL, Preeti

HILLMAN, Jennifer L.

YUE, Henry

AZIMZAI, Yalda

LU, Aina M.D.

BAUGHN, Mariah R.

TRAN, Bao

SHIH, Leo L.

AU-YOUNG, Janice

<120> NUCLEIC ACID-BINDING PROTEINS

<130> PF-0662 PCT

<140> To Be Assigned

<141> Herewith

<150> 60/117,905; 60/117,904

<151> 1999-01-29; 1999-01-29

<160> 110

<170> PERL Program

<210> 1

<211> 754

<212> PRT

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 025733CD1

<400> 1

```

Met Ala Ala Ala Gly Ser Arg Lys Arg Arg Leu Ala Glu Leu Thr
  1          5          10          15
Val Asp Glu Phe Leu Ala Ser Gly Phe Asp Ser Glu Ser Glu Ser
          20          25          30
Glu Ser Glu Asn Ser Pro Gln Ala Glu Thr Arg Glu Ala Arg Glu
          35          40          45
Ala Ala Arg Ser Pro Asp Lys Pro Gly Gly Ser Pro Ser Ala Ser
          50          55          60
Arg Arg Lys Gly Arg Ala Ser Glu His Lys Asp Gln Leu Ser Arg
          65          70          75
Leu Lys Asp Arg Asp Pro Glu Phe Tyr Lys Phe Leu Gln Glu Asn
          80          85          90
Asp Gln Ser Leu Leu Asn Phe Ser Asp Ser Asp Ser Ser Glu Glu
          95          100          105
Glu Glu Gly Pro Phe His Ser Leu Pro Asp Val Leu Glu Glu Ala
          110          115          120
Ser Glu Glu Glu Asp Gly Ala Glu Glu Gly Glu Asp Gly Asp Arg
          125          130          135
Val Pro Arg Gly Leu Lys Gly Lys Lys Asn Ser Val Pro Val Thr
          140          145          150
Val Ala Met Val Glu Arg Trp Lys Gln Ala Ala Lys Gln Arg Leu
          155          160          165

```

Thr	Pro	Lys	Leu	Phe	His	Glu	Val	Val	Gln	Ala	Phe	Arg	Ala	Ala	
				170					175					180	
Val	Ala	Thr	Thr	Arg	Gly	Asp	Gln	Glu	Ser	Ala	Glu	Ala	Asn	Lys	
				185					190					195	

Phe	Gln	Val	Thr	Asp	Ser	Ala	Ala	Phe	Asn	Ala	Leu	Val	Thr	Phe	
				200					205					210	
Cys	Ile	Arg	Asp	Leu	Ile	Gly	Cys	Leu	Gln	Lys	Leu	Leu	Phe	Gly	
				215					220					225	
Lys	Val	Ala	Lys	Asp	Ser	Ser	Arg	Met	Leu	Gln	Pro	Ser	Ser	Ser	
				230					235					240	
Pro	Leu	Trp	Gly	Lys	Leu	Arg	Val	Asp	Ile	Lys	Ala	Tyr	Leu	Gly	
				245					250					255	
Ser	Ala	Ile	Gln	Leu	Val	Ser	Cys	Leu	Ser	Glu	Thr	Thr	Val	Leu	
				260					265					270	
Ala	Ala	Val	Leu	Arg	His	Ile	Ser	Val	Leu	Val	Pro	Cys	Phe	Leu	
				275					280					285	
Thr	Phe	Pro	Lys	Gln	Cys	Arg	Met	Leu	Leu	Lys	Arg	Met	Val	Val	
				290					295					300	
Val	Trp	Ser	Thr	Gly	Glu	Glu	Ser	Leu	Arg	Val	Leu	Ala	Phe	Leu	
				305					310					315	
Val	Leu	Ser	Arg	Val	Cys	Arg	His	Lys	Lys	Asp	Thr	Phe	Leu	Gly	
				320					325					330	
Pro	Val	Leu	Lys	Gln	Met	Tyr	Ile	Thr	Tyr	Val	Arg	Asn	Cys	Lys	
				335					340					345	
Phe	Thr	Ser	Pro	Gly	Ala	Leu	Pro	Phe	Ile	Ser	Phe	Met	Gln	Trp	
				350					355					360	
Thr	Leu	Thr	Glu	Leu	Leu	Ala	Leu	Glu	Pro	Gly	Val	Ala	Tyr	Gln	
				365					370					375	
His	Ala	Phe	Leu	Tyr	Ile	Arg	Gln	Leu	Ala	Ile	His	Leu	Arg	Asn	
				380					385					390	
Ala	Met	Thr	Thr	Arg	Lys	Lys	Glu	Thr	Tyr	Gln	Ser	Val	Tyr	Asn	
				395					400					405	
Trp	Gln	Tyr	Val	His	Cys	Leu	Phe	Leu	Trp	Cys	Arg	Val	Leu	Ser	
				410					415					420	
Thr	Ala	Gly	Pro	Ser	Glu	Ala	Leu	Gln	Pro	Leu	Val	Tyr	Pro	Leu	
				425					430					435	
Ala	Gln	Val	Ile	Ile	Gly	Cys	Ile	Lys	Leu	Ile	Pro	Thr	Ala	Arg	
				440					445					450	
Phe	Tyr	Pro	Leu	Arg	Met	His	Cys	Ile	Arg	Ala	Leu	Thr	Leu	Leu	
				455					460					465	
Ser	Gly	Ser	Ser	Gly	Ala	Phe	Ile	Pro	Val	Leu	Pro	Phe	Ile	Leu	
				470					475					480	
Glu	Met	Phe	Gln	Gln	Val	Asp	Phe	Asn	Arg	Lys	Pro	Gly	Arg	Met	
				485					490					495	
Ser	Ser	Lys	Pro	Ile	Asn	Phe	Ser	Val	Ile	Leu	Lys	Leu	Ser	Asn	
				500					505					510	
Val	Asn	Leu	Gln	Glu	Lys	Ala	Tyr	Arg	Asp	Gly	Leu	Val	Glu	Gln	
				515					520					525	
Leu	Tyr	Asp	Leu	Thr	Leu	Glu	Tyr	Leu	His	Ser	Gln	Ala	His	Cys	
				530					535					540	
Ile	Gly	Phe	Pro	Glu	Leu	Val	Leu	Pro	Val	Val	Leu	Gln	Leu	Lys	
				545					550					555	
Ser	Phe	Leu	Arg	Glu	Cys	Lys	Val	Ala	Asn	Tyr	Cys	Arg	Gln	Val	
				560					565					570	
Gln	Gln	Leu	Leu	Gly	Lys	Val	Gln	Glu	Asn	Ser	Ala	Tyr	Ile	Cys	
				575					580					585	
Ser	Arg	Arg	Gln	Arg	Val	Ser	Phe	Gly	Val	Ser	Glu	Gln	Gln	Ala	
				590					595					600	
Val	Glu	Ala	Trp	Glu	Lys	Leu	Thr	Arg	Glu	Glu	Gly	Thr	Pro	Leu	
				605					610					615	

Thr	Leu	Tyr	Tyr	Ser	His	Trp	Arg	Lys	Leu	Arg	Asp	Arg	Glu	Ile
				620					625					630
Gln	Leu	Glu	Ile	Ser	Gly	Lys	Glu	Arg	Leu	Glu	Asp	Leu	Asn	Phe
				635					640					645
Pro	Glu	Ile	Lys	Arg	Arg	Lys	Met	Ala	Asp	Arg	Lys	Asp	Glu	Asp
				650					655					660
Arg	Lys	Gln	Phe	Lys	Asp	Leu	Phe	Asp	Leu	Asn	Ser	Ser	Glu	Glu
				665					670					675
Asp	Asp	Thr	Glu	Gly	Phe	Ser	Glu	Arg	Gly	Ile	Leu	Arg	Pro	Leu
				680					685					690
Ser	Thr	Arg	His	Gly	Val	Glu	Asp	Asp	Glu	Glu	Asp	Glu	Glu	Glu
				695					700					705
Gly	Glu	Glu	Asp	Ser	Ser	Asn	Ser	Glu	Gly	Glu	Trp	Ser	Trp	Asp
				710					715					720
Gly	Asp	Pro	Asp	Ala	Glu	Ala	Gly	Leu	Ala	Pro	Gly	Glu	Leu	Gln
				725					730					735
Gln	Leu	Ala	Gln	Gly	Pro	Glu	Asp	Glu	Leu	Glu	Asp	Leu	Gln	Leu
				740					745					750
Ser	Glu	Asp	Asp											

<210> 2
 <211> 593
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 079702CD1

<400> 2

Met	Arg	Asp	Ser	Thr	Gly	Ala	Gly	Asn	Ser	Leu	Val	His	Lys	Arg
1				5					10					15
Ser	Pro	Leu	Arg	Arg	Asn	Gln	Lys	Thr	Pro	Thr	Ser	Leu	Thr	Lys
				20					25					30
Leu	Ser	Leu	Gln	Asp	Gly	His	Lys	Ala	Lys	Lys	Pro	Ala	Cys	Lys
				35					40					45
Phe	Glu	Glu	Gly	Gln	Asp	Val	Leu	Ala	Arg	Trp	Ser	Asp	Gly	Leu
				50					55					60
Phe	Tyr	Leu	Gly	Thr	Ile	Lys	Lys	Ile	Asn	Ile	Leu	Lys	Gln	Ser
				65					70					75
Cys	Phe	Ile	Ile	Phe	Glu	Asp	Ser	Ser	Lys	Ser	Trp	Val	Leu	Trp
				80					85					90
Lys	Asp	Ile	Gln	Thr	Gly	Ala	Thr	Gly	Ser	Gly	Glu	Met	Val	Cys
				95					100					105
Thr	Ile	Cys	Gln	Glu	Glu	Tyr	Ser	Glu	Ala	Pro	Asn	Glu	Met	Val
				110					115					120
Ile	Cys	Asp	Lys	Cys	Gly	Gln	Gly	Tyr	His	Gln	Leu	Cys	His	Thr
				125					130					135
Pro	His	Ile	Asp	Ser	Ser	Val	Ile	Asp	Ser	Asp	Glu	Lys	Trp	Leu
				140					145					150
Cys	Arg	Gln	Cys	Val	Phe	Ala	Thr	Thr	Thr	Lys	Arg	Gly	Gly	Ala
				155					160					165
Leu	Lys	Lys	Gly	Pro	Asn	Ala	Lys	Ala	Leu	Gln	Val	Met	Lys	Gln
				170					175					180
Thr	Leu	Pro	Tyr	Ser	Val	Ala	Asp	Leu	Glu	Trp	Asp	Ala	Gly	His
				185					190					195
Lys	Thr	Asn	Val	Gln	Gln	Cys	Tyr	Cys	Tyr	Cys	Gly	Gly	Pro	Gly
				200					205					210
Asp	Trp	Tyr	Leu	Lys	Met	Leu	Gln	Cys	Cys	Lys	Cys	Lys	Gln	Trp

	215		220		225
Phe His Glu Ala Cys Val Gln Cys Leu		Gln Lys Pro Met Leu Phe			
	230		235		240
Gly Asp Arg Phe Tyr Thr Phe Ile Cys		Ser Val Cys Ser Ser Gly			
	245		250		255
Pro Glu Tyr Leu Lys Arg Leu Pro Leu		Gln Trp Val Asp Ile Ala			
	260		265		270
His Leu Cys Leu Tyr Asn Leu Ser Val		Ile His Lys Lys Lys Tyr			
	275		280		285
Phe Asp Ser Glu Leu Glu Leu Met Thr		Tyr Ile Asn Glu Asn Trp			
	290		295		300
Asp Arg Leu His Pro Gly Glu Leu Ala		Asp Thr Pro Lys Ser Glu			
	305		310		315
Arg Tyr Glu His Val Leu Glu Ala Leu		Asn Asp Tyr Lys Thr Met			
	320		325		330
Phe Met Ser Gly Lys Glu Ile Lys Lys		Lys Lys His Leu Phe Gly			
	335		340		345
Leu Arg Ile Arg Val Pro Pro Val Pro		Pro Asn Val Ala Phe Lys			
	350		355		360
Ala Glu Lys Glu Pro Glu Gly Thr Ser		His Glu Phe Lys Ile Lys			
	365		370		375
Gly Arg Lys Ala Ser Lys Pro Ile Ser		Asp Ser Arg Glu Val Ser			
	380		385		390
Asn Gly Ile Glu Lys Lys Lys Lys Lys		Lys Ser Val Gly Arg Pro			
	395		400		405
Pro Gly Pro Tyr Thr Arg Lys Met Ile		Gln Lys Thr Ala Glu Pro			
	410		415		420
Leu Leu Asp Lys Glu Ser Ile Ser Glu		Asn Pro Thr Leu Asp Leu			
	425		430		435
Pro Cys Ser Ile Gly Arg Thr Glu Gly		Thr Ala His Ser Ser Asn			
	440		445		450
Thr Ser Asp Val Asp Phe Thr Gly Ala		Ser Ser Ala Lys Glu Thr			
	455		460		465
Thr Ser Ser Ser Ile Ser Arg His Tyr		Gly Leu Ser Asp Ser Arg			
	470		475		480
Lys Arg Thr Arg Thr Gly Arg Ser Trp		Pro Ala Ala Ile Pro His			
	485		490		495
Leu Arg Arg Arg Arg Gly Arg Leu Pro		Arg Arg Ala Leu Gln Thr			
	500		505		510
Gln Asn Ser Glu Ile Val Lys Asp Asp		Glu Gly Lys Glu Asp Tyr			
	515		520		525
Gln Phe Asp Glu Leu Asn Thr Glu Ile		Leu Asn Asn Leu Ala Asp			
	530		535		540
Gln Glu Leu Gln Leu Asn His Leu Lys		Asn Ser Ile Thr Ser Tyr			
	545		550		555
Phe Gly Ala Ala Gly Arg Ile Ala Cys		Gly Glu Lys Tyr Arg Val			
	560		565		570
Leu Ala Arg Arg Val Thr Leu Asp Gly		Lys Val Gln Tyr Leu Val			
	575		580		585
Glu Trp Glu Gly Ala Thr Ala Ser					
	590				

<210> 3
 <211> 534
 <212> PRT
 <213> Homo sapiens

 <220>

<221> misc-feature

<223> Incyte ID No.: 116208CD1

<400> 3

```

Met Arg Ala Leu His Leu Leu Lys Ser Gly Cys Ser Pro Ala Val
 1           5           10           15
Gln Ile Lys Ile Arg Glu Leu Tyr Arg Arg Arg Tyr Pro Arg Thr
           20           25           30
Leu Glu Gly Leu Ser Asp Leu Ser Thr Ile Lys Ser Ser Val Phe
           35           40           45
Ser Leu Asp Gly Gly Ser Ser Pro Val Glu Pro Asp Leu Ala Val
           50           55           60
Ala Gly Ile His Ser Leu Pro Ser Thr Ser Val Thr Pro His Ser
           65           70           75
Pro Ser Ser Pro Val Gly Ser Val Leu Leu Gln Asp Thr Lys Pro
           80           85           90
Thr Phe Glu Met Gln Gln Pro Ser Pro Pro Ile Pro Pro Val His
           95          100          105
Pro Asp Val Gln Leu Lys Asn Leu Pro Phe Tyr Asp Val Leu Asp
          110          115          120
Val Leu Ile Lys Pro Thr Ser Leu Val Gln Ser Ser Ile Gln Arg
          125          130          135
Phe Gln Glu Lys Phe Phe Ile Phe Ala Leu Thr Pro Gln Gln Val
          140          145          150
Arg Glu Ile Cys Ile Ser Arg Asp Phe Leu Pro Gly Gly Arg Arg
          155          160          165
Asp Tyr Thr Val Gln Val Gln Leu Arg Leu Cys Leu Ala Glu Thr
          170          175          180
Ser Cys Pro Gln Glu Asp Asn Tyr Pro Asn Ser Leu Cys Ile Lys
          185          190          195
Val Asn Gly Lys Leu Phe Pro Leu Pro Gly Tyr Ala Pro Pro Pro
          200          205          210
Lys Asn Gly Ile Glu Gln Lys Arg Pro Gly Arg Pro Leu Asn Ile
          215          220          225
Thr Ser Leu Val Arg Leu Ser Ser Ala Val Pro Asn Gln Ile Ser
          230          235          240
Ile Ser Trp Ala Ser Glu Ile Gly Lys Asn Tyr Ser Met Ser Val
          245          250          255
Tyr Leu Val Arg Gln Leu Thr Ser Ala Met Leu Leu Gln Arg Leu
          260          265          270
Lys Met Lys Gly Ile Arg Asn Pro Asp His Ser Arg Ala Leu Ser
          275          280          285
Lys Glu Lys Leu Thr Ala Asp Pro Asp Ser Glu Ile Ala Thr Thr
          290          295          300
Ser Leu Arg Val Ser Leu Met Cys Pro Leu Gly Lys Met Arg Leu
          305          310          315
Thr Ile Pro Cys Arg Ala Val Thr Cys Thr His Leu Gln Cys Phe
          320          325          330
Asp Ala Ala Leu Tyr Leu Gln Met Asn Glu Lys Lys Pro Thr Trp
          335          340          345
Ile Cys Pro Val Cys Asp Lys Lys Ala Ala Tyr Glu Ser Leu Ile
          350          355          360
Leu Asp Gly Leu Phe Met Glu Ile Leu Asn Asp Cys Ser Asp Val
          365          370          375
Asp Glu Ile Lys Phe Gln Glu Asp Gly Ser Trp Cys Pro Met Arg
          380          385          390
Pro Lys Lys Glu Ala Met Lys Val Ser Ser Gln Pro Cys Thr Lys
          395          400          405
Ile Glu Ser Ser Ser Val Leu Ser Lys Pro Cys Ser Val Thr Val
          410          415          420
Ala Ser Glu Ala Ser Lys Lys Lys Val Asp Val Ile Asp Leu Thr
          425          430          435

```

```

Ile Glu Ser Ser Ser Asp Glu Glu Glu Asp Pro Pro Ala Lys Arg
      440      445      450
Lys Cys Ile Phe Met Ser Glu Thr Gln Ser Ser Pro Thr Lys Gly
      455      460      465
Val Leu Met Tyr Gln Pro Ser Ser Val Arg Val Pro Ser Val Thr
      470      475      480
Ser Val Asp Pro Ala Ala Ile Pro Pro Ser Leu Thr Asp Tyr Ser
      485      490      495
Val Pro Phe His His Thr Pro Ile Ser Ser Met Ser Ser Asp Leu
      500      505      510
Pro Gly Glu Gln Arg Arg Asn Asp Ile Asn Asn Glu Leu Lys Leu
      515      520      525
Gly Thr Ser Ser Asp Thr Val Gln Gln
      530

```

<210> 4
 <211> 255
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc feature
 <223> Incyte ID No.: 179261CD1

```

<400> 4
Met Lys Ile Ile Ser Glu Asn Val Pro Ser Tyr Lys Thr His Glu
  1      5      10
Ser Leu Thr Leu Pro Arg Arg Thr His Asp Ser Glu Lys Pro Tyr
      20      25      30
Glu Tyr Lys Glu Tyr Glu Lys Val Phe Ser Cys Asp Leu Glu Phe
      35      40      45
Asp Glu Tyr Gln Lys Ile His Thr Gly Gly Lys Asn Tyr Glu Cys
      50      55      60
Asn Gln Cys Trp Lys Thr Phe Gly Ile Asp Asn Ser Ser Met Leu
      65      70      75
Gln Leu Asn Ile His Thr Gly Val Lys Pro Cys Lys Tyr Met Glu
      80      85      90
Tyr Gly Asn Thr Cys Ser Phe Tyr Lys Asp Phe Asn Val Tyr Gln
      95      100      105
Lys Ile His Asn Glu Lys Phe Tyr Lys Cys Lys Glu Tyr Arg Arg
      110      115      120
Thr Phe Glu Arg Val Gly Lys Val Thr Pro Leu Gln Arg Val His
      125      130      135
Asp Gly Glu Lys His Phe Glu Cys Ser Phe Cys Gly Lys Ser Phe
      140      145      150
Arg Val His Ala Gln Leu Thr Arg His Gln Lys Ile His Thr Asp
      155      160      165
Glu Lys Thr Tyr Lys Cys Met Glu Cys Gly Lys Asp Phe Arg Phe
      170      175      180
His Ser Gln Leu Thr Glu His Gln Arg Ile His Thr Gly Glu Lys
      185      190      195
Pro Tyr Lys Cys Met His Cys Glu Lys Val Phe Arg Ile Ser Ser
      200      205      210
Gln Leu Ile Glu His Gln Arg Ile His Thr Gly Glu Lys Pro Tyr
      215      220      225
Ala Cys Lys Glu Cys Gly Lys Ala Phe Gly Val Cys Arg Glu Leu
      230      235      240
Ala Arg His Gln Arg Ile His Thr Gly Lys Tyr Cys Gly Trp Ile
      245      250      255

```

<210> 5
 <211> 562
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc feature
 <223> Incyte ID No.: 259161CD1

<400> 5
 Met Ala Ser Val Ser Ala Leu Thr Glu Glu Leu Asp Ser Ile Thr
 1 5 10 15
 Ser Glu Leu His Ala Val Glu Ile Gln Ile Gln Glu Leu Thr Glu
 20 25 30
 Arg Gln Gln Glu Leu Ile Gln Lys Lys Val Leu Thr Lys Lys
 35 40 45
 Ile Lys Gln Cys Leu Glu Asp Ser Asp Ala Gly Ala Ser Asn Glu
 50 55 60
 Tyr Asp Ser Ser Pro Ala Ala Trp Asn Lys Glu Asp Phe Pro Trp
 65 70 75
 Ser Gly Lys Val Lys Asp Ile Leu Gln Asn Val Phe Lys Leu Glu
 80 85 90
 Lys Phe Arg Pro Leu Gln Leu Glu Thr Ile Asn Val Thr Met Ala
 95 100 105
 Gly Lys Glu Val Phe Leu Val Met Pro Thr Gly Gly Gly Lys Ser
 110 115 120
 Leu Cys Tyr Gln Leu Pro Ala Leu Cys Ser Asp Gly Phe Thr Leu
 125 130 135
 Val Ile Cys Pro Leu Ile Ser Leu Met Glu Asp Gln Leu Met Val
 140 145 150
 Leu Lys Gln Leu Gly Ile Ser Ala Thr Met Leu Asn Ala Ser Ser
 155 160 165
 Ser Lys Glu His Val Lys Trp Val His Ala Glu Met Val Asn Lys
 170 175 180
 Asn Ser Glu Leu Lys Leu Ile Tyr Val Thr Pro Glu Lys Ile Ala
 185 190 195
 Lys Ser Lys Met Phe Met Ser Arg Leu Glu Lys Ala Tyr Glu Ala
 200 205 210
 Arg Arg Phe Thr Arg Ile Ala Val Asp Glu Val His Cys Cys Ser
 215 220 225
 Gln Trp Gly His Asp Phe Arg Pro Asp Tyr Lys Ala Leu Gly Ile
 230 235 240
 Leu Lys Arg Gln Phe Pro Asn Ala Ser Leu Ile Gly Leu Thr Ala
 245 250 255
 Thr Ala Thr Asn His Val Leu Thr Asp Ala Gln Lys Ile Leu Cys
 260 265 270
 Ile Glu Lys Cys Phe Thr Phe Thr Ala Ser Phe Asn Lys Pro Asp
 275 280 285
 Val Arg Phe Val Ile His His Ser Met Ser Lys Ser Met Glu Asn
 290 295 300
 Tyr Tyr Gln Glu Ser Gly Arg Ala Gly Arg Asp Asp Met Lys Ala
 305 310 315
 Asp Cys Ile Leu Tyr Tyr Gly Phe Gly Asp Ile Phe Arg Ile Ser
 320 325 330
 Ser Met Val Val Met Glu Asn Val Gly Gln Gln Lys Leu Tyr Glu
 335 340 345
 Met Val Ser Tyr Cys Gln Asn Ile Ser Lys Cys Arg Arg Val Leu
 350 355 360
 Met Ala Gln His Phe Asp Glu Val Trp Asn Ser Glu Ala Cys Asn

	365		370		375
Lys Met Cys Asp	Asn Cys Cys Lys Asp	Ser Ala Phe Glu Arg Lys			
	380		385		390
Asn Ile Thr Glu	Tyr Cys Arg Asp Leu	Ile Lys Ile Leu Lys Gln			
	395		400		405
Ala Glu Glu Leu	Asn Glu Lys Leu Thr	Pro Leu Lys Leu Ile Asp			
	410		415		420
Ser Trp Met Gly	Lys Gly Ala Ala Lys	Leu Arg Val Ala Gly Val			
	425		430		435
Val Ala Pro Thr	Leu Pro Arg Glu Asp	Leu Glu Lys Ile Ile Ala			
	440		445		450
His Phe Leu Ile	Gln Gln Tyr Leu Lys	Glu Asp Tyr Ser Phe Thr			
	455		460		465
Ala Tyr Ala Thr	Ile Ser Tyr Leu Lys	Ile Gly Pro Lys Ala Asn			
	470		475		480
Leu Leu Asn Asn	Glu Ala His Ala Ile	Thr Met Gln Val Thr Lys			
	485		490		495
Ser Thr Gln Asn	Ser Phe Arg Ala Glu	Ser Ser Gln Thr Cys His			
	500		505		510
Ser Glu Gln Gly	Asp Lys Lys Ile Gly	Gly Lys Lys Phe Gln Ala			
	515		520		525
Thr Ser Arg Arg	Arg Leu Gln Thr Cys	Phe Ser Asn Leu Val Leu			
	530		535		540
Arg Ile Gln Glu	Leu Arg Lys Glu Lys	Ser Met Met Pro Asp Met			
	545		550		555
Thr Val Thr Lys	Phe Ser Asn				
	560				

<210> 6

<211> 432

<212> PRT

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 320087CD1

<400> 6

Met Ser Glu Leu Lys	Asp Cys Pro Leu Gln	Phe His Asp Phe Lys
1	5	10
Ser Val Asp His Leu	Lys Val Cys Pro Arg	Tyr Thr Ala Val Leu
	20	25
Ala Arg Ser Glu Asp	Asp Gly Ile Gly Ile	Glu Glu Leu Asp Thr
	35	40
Leu Gln Leu Glu Leu	Glu Thr Leu Leu Ser	Ser Ala Ser Arg Arg
	50	55
Leu Arg Val Leu Glu	Ala Glu Thr Gln Ile	Leu Thr Asp Trp Gln
	65	70
Asp Lys Lys Gly Asp	Arg Arg Phe Leu Lys	Leu Gly Arg Asp His
	80	85
Glu Leu Gly Ala Pro	Lys His Gly Lys Pro	Lys Lys Gln Lys
	95	100
Leu Glu Gly Lys Ala	Gly His Gly Pro Gly	Pro Gly Pro Gly Arg
	110	115
Pro Lys Ser Lys Asn	Leu Gln Pro Lys Ile	Gln Glu Tyr Glu Phe
	125	130
Thr Asp Asp Pro Ile	Asp Val Pro Arg Ile	Pro Lys Asn Asp Ala
	140	145
		150

```

Pro Asn Arg Phe Trp Ala Ser Val Glu Pro Tyr Cys Ala Asp Ile
155 160 165
Thr Ser Glu Glu Val Arg Thr Leu Glu Glu Leu Leu Lys Pro Pro
170 175 180
Glu Asp Glu Ala Glu His Tyr Lys Ile Pro Pro Leu Gly Lys His
185 190 195
Tyr Ser Gln Arg Trp Ala Gln Glu Asp Leu Leu Glu Glu Gln Lys
200 205 210
Asp Gly Ala Arg Ala Ala Ala Val Ala Asp Lys Lys Lys Gly Leu
215 220 225
Met Gly Pro Leu Thr Glu Leu Asp Thr Lys Asp Val Asp Ala Leu
230 235 240
Leu Lys Lys Ser Glu Ala Gln His Glu Gln Pro Glu Asp Gly Cys
245 250 255
Pro Phe Gly Ala Leu Thr Gln Arg Leu Leu Gln Ala Leu Val Glu
260 265 270
Glu Asn Ile Ile Ser Pro Met Glu Asp Ser Pro Ile Pro Asp Met
275 280 285
Ser Gly Lys Glu Ser Gly Ala Asp Gly Ala Ser Thr Ser Pro Arg
290 295 300
Asn Gln Asn Lys Pro Phe Ser Val Pro His Thr Lys Ser Leu Glu
305 310 315
Ser Arg Ile Lys Glu Glu Leu Ile Ala Gln Gly Leu Leu Glu Ser
320 325 330
Glu Asp Arg Pro Ala Glu Asp Ser Glu Asp Glu Val Leu Ala Glu
335 340 345
Leu Arg Lys Arg Gln Ala Glu Leu Lys Ala Leu Ser Ala His Asn
350 355 360
Arg Thr Lys Lys His Asp Leu Leu Arg Leu Ala Lys Glu Glu Val
365 370 375
Ser Arg Gln Glu Leu Arg Gln Arg Val Arg Met Ala Asp Asn Glu
380 385 390
Val Met Asp Ala Phe Arg Lys Ile Met Ala Ala Arg Gln Lys Lys
395 400 405
Arg Thr Pro Thr Lys Lys Glu Lys Asp Gln Ala Trp Lys Thr Leu
410 415 420
Lys Glu Arg Glu Ser Ile Leu Lys Leu Leu Asp Gly
425 430

```

<210> 7
 <211> 799
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> 491271CD1

```

<400> 7
Met Pro Ser Gln Asn Tyr Asp Leu Pro Gln Lys Lys Gln Glu Lys
1 5 10 15
Met Thr Lys Phe Gln Glu Ala Val Thr Phe Lys Asp Val Ala Val
20 25 30
Val Phe Ser Arg Glu Glu Leu Arg Leu Leu Asp Leu Thr Gln Arg
35 40 45
Lys Leu Tyr Arg Asp Val Met Val Glu Asn Phe Lys Asn Leu Val
50 55 60
Ala Val Gly His Leu Pro Phe Gln Pro Asp Met Val Ser Gln Leu
65 70 75

```

Glu	Ala	Glu	Glu	Lys	Leu	Trp	Met	Met	Glu	Thr	Glu	Thr	Gln	Arg
				80					85					90
Ser	Ser	Lys	His	Gln	Asn	Lys	Met	Glu	Thr	Leu	Gln	Lys	Phe	Ala
				95					100					105
Leu	Lys	Tyr	Leu	Ser	Asn	Gln	Glu	Leu	Ser	Cys	Trp	Gln	Ile	Trp
				110					115					120
Lys	Gln	Val	Ala	Ser	Glu	Leu	Thr	Arg	Cys	Leu	Gln	Gly	Lys	Ser
				125					130					135
Ser	Gln	Leu	Leu	Gln	Gly	Asp	Ser	Ile	Gln	Val	Ser	Glu	Asn	Glu
				140					145					150
Asn	Asn	Ile	Met	Asn	Pro	Lys	Gly	Asp	Ser	Pro	Ile	Tyr	Ile	Glu
				155					160					165
Asn	Gln	Glu	Phe	Pro	Phe	Trp	Arg	Thr	Gln	His	Ser	Cys	Gly	Asn
				170					175					180
Thr	Tyr	Leu	Ser	Glu	Ser	Gln	Ile	Gln	Ser	Arg	Gly	Lys	Gln	Ile
				185					190					195
Asp	Val	Lys	Asn	Asn	Leu	Gln	Ile	Arg	Glu	Asp	Phe	Val	Lys	Lys
				200					205					210
Ser	Pro	Phe	His	Glu	His	Ile	Lys	Thr	Asp	Thr	Glu	Pro	Lys	Pro
				215					220					225
Cys	Lys	Gly	Asn	Glu	Tyr	Gly	Lys	Ile	Ile	Ser	Asp	Gly	Ser	Asn
				230					235					240
Gln	Lys	Leu	Pro	Leu	Gly	Glu	Lys	Pro	His	Pro	Cys	Gly	Glu	Cys
				245					250					255
Gly	Arg	Gly	Phe	Ser	Tyr	Ser	Pro	Arg	Leu	Pro	Leu	His	Pro	Asn
				260					265					270
Val	His	Thr	Gly	Glu	Lys	Cys	Phe	Ser	Gln	Ser	Ser	His	Leu	Arg
				275					280					285
Thr	His	Gln	Arg	Ile	His	Pro	Gly	Glu	Lys	Leu	Asn	Arg	Cys	His
				290					295					300
Glu	Ser	Gly	Asp	Cys	Phe	Asn	Lys	Ser	Ser	Phe	His	Ser	Tyr	Gln
				305					310					315
Ser	Asn	His	Thr	Gly	Glu	Lys	Ser	Tyr	Arg	Cys	Asp	Ser	Cys	Gly
				320					325					330
Lys	Gly	Phe	Ser	Ser	Ser	Thr	Gly	Leu	Ile	Ile	His	Tyr	Arg	Thr
				335					340					345
His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys	Cys	Glu	Glu	Cys	Gly	Lys	Cys
				350					355					360
Phe	Ser	Gln	Ser	Ser	Asn	Phe	Gln	Cys	His	Gln	Arg	Val	His	Thr
				365					370					375
Glu	Glu	Lys	Pro	Tyr	Lys	Cys	Glu	Glu	Cys	Gly	Lys	Gly	Phe	Gly
				380					385					390
Trp	Ser	Val	Asn	Leu	Arg	Val	His	Gln	Arg	Val	His	Arg	Gly	Glu
				395					400					405
Lys	Pro	Tyr	Lys	Cys	Glu	Glu	Cys	Gly	Lys	Gly	Phe	Thr	Gln	Ala
				410					415					420
Ala	His	Phe	His	Ile	His	Gln	Arg	Val	His	Thr	Gly	Glu	Lys	Pro
				425					430					435
Tyr	Lys	Cys	Asp	Val	Cys	Gly	Lys	Gly	Phe	Ser	His	Asn	Ser	Pro
				440					445					450
Leu	Ile	Cys	His	Arg	Arg	Val	His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys
				455					460					465
Cys	Glu	Ala	Cys	Gly	Lys	Gly	Phe	Thr	Arg	Asn	Thr	Asp	Leu	His
				470					475					480
Ile	His	Phe	Arg	Val	His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys	Cys	Lys
				485					490					495
Glu	Cys	Gly	Lys	Gly	Phe	Ser	Gln	Ala	Ser	Asn	Leu	Gln	Val	His
				500					505					510
Gln	Asn	Val	His	Thr	Gly	Glu	Lys	Arg	Phe	Lys	Cys	Glu	Thr	Cys
				515					520					525
Gly	Lys	Gly	Phe	Ser	Gln	Ser	Ser	Lys	Leu	Gln	Thr	His	Gln	Arg

	530		535		540
Val His Thr Gly	Glu Lys Pro Tyr Arg	Cys Asp Val Cys Gly Lys			
	545		550		555
Asp Phe Ser Tyr	Ser Ser Asn Leu Lys	Leu His Gln Val Ile His			
	560		565		570
Thr Gly Glu Lys	Pro Tyr Lys Cys Glu	Glu Cys Gly Lys Gly Phe			
	575		580		585
Ser Trp Arg Ser	Asn Leu His Ala His	Gln Arg Val His Ser Gly			
	590		595		600
Glu Lys Pro Tyr	Lys Cys Glu Gln Cys	Asp Lys Ser Phe Ser Gln			
	605		610		615
Ala Ile Asp Phe	Arg Val His Gln Arg	Val His Thr Gly Glu Lys			
	620		625		630
Pro Tyr Lys Cys	Gly Val Cys Gly Lys	Gly Phe Ser Gln Ser Ser			
	635		640		645
Gly Leu Gln Ser	His Gln Arg Val His	Thr Gly Glu Lys Pro Tyr			
	650		655		660
Lys Cys Asp Val	Cys Gly Lys Gly Phe	Arg Tyr Ser Ser Gln Phe			
	665		670		675
Ile Tyr His Gln	Arg Gly His Thr Gly	Glu Lys Pro Tyr Lys Cys			
	680		685		690
Glu Glu Cys Gly	Lys Gly Phe Gly Arg	Ser Leu Asn Leu Arg His			
	695		700		705
His Gln Arg Val	His Thr Gly Glu Lys	Pro His Ile Cys Glu Glu			
	710		715		720
Cys Gly Lys Ala	Phe Ser Leu Pro Ser	Asn Leu Arg Val His Leu			
	725		730		735
Gly Val His Thr	Arg Glu Lys Leu Phe	Lys Cys Glu Glu Cys Gly			
	740		745		750
Lys Gly Phe Ser	Gln Ser Ala Arg Leu	Glu Ala His Gln Arg Val			
	755		760		765
His Thr Gly Glu	Lys Pro Tyr Lys Cys	Asp Ile Cys Asp Lys Asp			
	770		775		780
Phe Arg His Arg	Ser Arg Leu Thr Tyr	His Gln Lys Val His Thr			
	785		790		795
Gly Lys Lys Leu					

<210> 8
 <211> 137
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 585172CD1

<400> 8	
Met Leu Ser Gly Arg Leu Val Leu Gly Leu Val Ser Met Ala Gly	
1 5 10 15	
Arg Val Cys Leu Cys Gln Gly Ser Ala Gly Ser Gly Ala Ile Gly	
20 25 30	
Pro Val Glu Ala Ala Ile Arg Thr Lys Leu Glu Glu Ala Leu Ser	
35 40 45	
Pro Glu Val Leu Glu Leu Arg Asn Glu Ser Gly Gly His Ala Val	
50 55 60	
Pro Pro Gly Ser Glu Thr His Phe Arg Val Ala Val Val Ser Ser	
65 70 75	
Arg Phe Glu Gly Leu Ser Pro Leu Gln Arg His Arg Leu Val His	

	80		85		90
Ala	Ala	Leu	Ala	Glu	Glu
		Leu	Gly	Gly	Pro
				Val	His
				Ala	Leu
				Ala	
	95				100
Ile	Gln	Ala	Arg	Thr	Pro
		Ala	Gln	Trp	Arg
				Glu	Asn
				Ser	Gln
				Leu	
	110				115
Asp	Thr	Ser	Pro	Pro	Cys
		Leu	Gly	Gly	Asn
				Lys	Lys
				Thr	Leu
				Gly	
	125				130
					135
Thr	Pro				

<210> 9
 <211> 230
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 615200CD1

<400> 9
 Met Val Gly Ala Gly Ile Ser Thr Pro Ser Gly Ile Pro Asp Phe
 1 5 10 15
 Arg Ser Pro Gly Ser Gly Leu Tyr Ser Asn Leu Gln Gln Tyr Asp
 20 25 30
 Leu Pro Tyr Pro Glu Ala Ile Phe Glu Leu Pro Phe Phe Phe His
 35 40 45
 Asn Pro Lys Pro Phe Phe Thr Leu Ala Lys Glu Leu Tyr Pro Gly
 50 55 60
 Asn Tyr Lys Pro Asn Ile Thr His Tyr Phe Leu Arg Leu Leu His
 65 70 75
 Asp Lys Gly Leu Leu Leu Arg Leu Tyr Thr Gln Asn Ile Asp Gly
 80 85 90
 Leu Glu Arg Val Ser Gly Ile Pro Ala Ser Lys Leu Val Glu Ala
 95 100 105
 His Gly Thr Phe Ala Ser Ala Thr Cys Thr Val Cys Gln Arg Pro
 110 115 120
 Phe Pro Gly Glu Asp Ile Arg Ala Asp Val Met Ala Asp Arg Val
 125 130 135
 Pro Arg Cys Pro Val Cys Thr Gly Val Val Lys Pro Asp Ile Val
 140 145 150
 Phe Phe Gly Glu Pro Leu Pro Gln Arg Phe Leu Leu His Val Val
 155 160 165
 Asp Phe Pro Met Ala Asp Leu Leu Leu Ile Leu Gly Thr Ser Leu
 170 175 180
 Glu Val Glu Pro Phe Ala Ser Leu Thr Glu Ala Val Arg Thr Gln
 185 190 195
 Phe Pro Asp Cys Ser Ser Thr Gly Thr Trp Trp Gly Pro Trp Leu
 200 205 210
 Gly Ile Leu Ala Ala Gly Thr Trp Pro Ser Trp Gly Thr Trp Phe
 215 220 225
 Thr Ala Trp Lys Ala
 230

<210> 10
 <211> 446
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 997067CD1

<400> 10
 Met Glu Thr Gln Ala Asp Leu Val Ser Gln Glu Pro Gln Ala Leu
 1 5 10 15
 Leu Asp Ser Ala Leu Pro Ser Lys Val Pro Ala Phe Ser Asp Lys
 20 25 30
 Asp Ser Leu Gly Asp Glu Met Leu Ala Ala Ala Leu Leu Lys Ala
 35 40 45
 Lys Ser Gln Glu Leu Val Thr Phe Glu Asp Val Ala Val Tyr Phe
 50 55 60
 Ile Arg Lys Glu Trp Lys Arg Leu Glu Pro Ala Gln Arg Asp Leu
 65 70 75
 Tyr Arg Asp Val Met Leu Glu Asn Tyr Gly Asn Val Phe Ser Leu
 80 85 90
 Asp Arg Glu Thr Arg Thr Glu Asn Asp Gln Glu Ile Ser Glu Asp
 95 100 105
 Thr Arg Ser His Gly Val Leu Leu Gly Arg Phe Gln Lys Asp Ile
 110 115 120
 Ser Gln Gly Leu Lys Phe Lys Glu Ala Tyr Glu Arg Glu Val Ser
 125 130 135
 Leu Lys Arg Pro Leu Gly Asn Ser Pro Gly Glu Arg Leu Asn Arg
 140 145 150
 Lys Met Pro Asp Phe Gly Gln Val Thr Val Glu Glu Lys Leu Thr
 155 160 165
 Pro Arg Gly Glu Arg Ser Glu Lys Tyr Asn Asp Phe Gly Asn Ser
 170 175 180
 Phe Thr Val Asn Ser Asn Leu Ile Ser His Gln Arg Leu Pro Val
 185 190 195
 Gly Asp Arg Pro His Lys Cys Asp Glu Cys Ser Lys Ser Phe Asn
 200 205 210
 Arg Thr Ser Asp Leu Ile Gln His Gln Arg Ile His Thr Gly Glu
 215 220 225
 Lys Pro Tyr Glu Cys Asn Glu Cys Gly Lys Ala Phe Ser Gln Ser
 230 235 240
 Ser His Leu Ile Gln His Gln Arg Ile His Thr Gly Glu Lys Pro
 245 250 255
 Tyr Glu Cys Ser Asp Cys Gly Lys Thr Phe Ser Cys Ser Ser Ala
 260 265 270
 Leu Ile Leu His Arg Arg Ile His Thr Gly Glu Lys Pro Tyr Glu
 275 280 285
 Cys Asn Glu Cys Gly Lys Thr Phe Ser Trp Ser Ser Thr Leu Thr
 290 295 300
 His His Gln Arg Ile His Thr Gly Glu Lys Pro Tyr Ala Cys Asn
 305 310 315
 Glu Cys Gly Lys Ala Phe Ser Arg Ser Ser Thr Leu Ile His His
 320 325 330
 Gln Arg Ile His Thr Gly Glu Lys Pro Tyr Glu Cys Asn Glu Cys
 335 340 345
 Gly Lys Ala Phe Ser Gln Ser Ser His Leu Tyr Gln His Gln Arg
 350 355 360
 Ile His Thr Gly Glu Lys Pro Tyr Glu Cys Met Glu Cys Gly Gly
 365 370 375
 Lys Phe Thr Tyr Ser Ser Gly Leu Ile Gln His Gln Arg Ile His
 380 385 390
 Thr Gly Glu Asn Pro Tyr Glu Cys Ser Glu Cys Gly Lys Ala Phe
 395 400 405
 Arg Tyr Ser Ser Ala Leu Val Arg His Gln Arg Ile His Thr Gly
 410 415 420
 Glu Lys Pro Leu Asn Gly Ile Gly Met Ser Lys Ser Ser Leu Arg

	425		430		435
Val Thr Thr Glu	Leu Asn Ile Arg Glu Ser Thr				
	440		445		

<210> 11
 <211> 428
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID NO.: 1443262CD1

<400> 11
 Met Glu Pro Leu Lys Val Glu Lys Phe Ala Thr Ala Asn Arg Gly
 1 5 10 15
 Asn Gly Leu Arg Ala Val Thr Pro Leu Arg Pro Gly Glu Leu Leu
 20 25 30
 Phe Arg Ser Asp Pro Leu Ala Tyr Thr Val Cys Lys Gly Ser Arg
 35 40 45
 Gly Val Val Cys Asp Arg Cys Leu Leu Gly Lys Glu Lys Leu Met
 50 55 60
 Arg Cys Ser Gln Cys Arg Val Ala Lys Tyr Cys Ser Ala Lys Cys
 65 70 75
 Gln Lys Lys Ala Trp Pro Asp His Lys Arg Glu Cys Lys Cys Leu
 80 85 90
 Lys Ser Cys Lys Pro Arg Tyr Pro Pro Asp Ser Val Arg Leu Leu
 95 100 105
 Gly Arg Val Val Phe Lys Leu Met Asp Gly Ala Pro Ser Glu Ser
 110 115 120
 Glu Lys Leu Tyr Ser Phe Tyr Asp Leu Glu Ser Asn Ile Asn Lys
 125 130 135
 Leu Thr Glu Asp Lys Lys Glu Gly Leu Arg Gln Leu Val Met Thr
 140 145 150
 Phe Gln His Phe Met Arg Glu Glu Ile Gln Asp Ala Ser Gln Leu
 155 160 165
 Pro Pro Ala Phe Asp Leu Phe Glu Ala Phe Ala Lys Val Ile Cys
 170 175 180
 Asn Ser Phe Thr Ile Cys Asn Ala Glu Met Gln Glu Val Gly Val
 185 190 195
 Gly Leu Tyr Pro Ser Ile Ser Leu Leu Asn His Ser Cys Asp Pro
 200 205 210
 Asn Cys Ser Ile Val Phe Asn Gly Pro His Leu Leu Leu Arg Ala
 215 220 225
 Val Arg Asp Ile Glu Val Gly Glu Glu Leu Thr Ile Cys Tyr Leu
 230 235 240
 Asp Met Leu Met Thr Ser Glu Glu Arg Arg Lys Gln Leu Arg Asp
 245 250 255
 Gln Tyr Cys Phe Glu Cys Asp Cys Phe Arg Cys Gln Thr Gln Asp
 260 265 270
 Lys Asp Ala Asp Met Leu Thr Gly Asp Glu Gln Val Trp Lys Glu
 275 280 285
 Val Gln Glu Ser Leu Lys Lys Ile Glu Glu Leu Lys Ala His Trp
 290 295 300
 Lys Trp Glu Gln Val Leu Ala Met Cys Gln Ala Ile Ile Ser Ser
 305 310 315
 Asn Ser Glu Arg Leu Pro Asp Ile Asn Ile Tyr Gln Leu Lys Val
 320 325 330

```

Leu Asp Cys Ala Met Asp Ala Cys Ile Asn Leu Gly Leu Leu Glu
335 340 345
Glu Ala Leu Phe Tyr Gly Thr Arg Thr Met Glu Pro Tyr Arg Ile
350 355 360
Phe Phe Pro Gly Ser His Pro Val Arg Gly Val Gln Val Met Lys
365 370 375
Val Gly Lys Leu Gln Leu His Gln Gly Met Phe Pro Gln Ala Met
380 385 390
Lys Asn Leu Arg Leu Ala Phe Asp Ile Met Arg Val Thr His Gly
395 400 405
Arg Glu His Ser Leu Ile Glu Asp Leu Ile Leu Leu Leu Glu Glu
410 415 420
Cys Asp Ala Asn Ile Arg Ala Ser
425

```

<210> 12
 <211> 590
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1521648CD1

```

<400> 12
Met Ala Glu Asp Trp Leu Asp Cys Pro Ala Leu Gly Pro Gly Trp
1 5 10 15
Lys Arg Arg Glu Val Phe Arg Lys Ser Gly Ala Thr Cys Gly Arg
20 25 30
Ser Asp Thr Tyr Tyr Gln Ser Pro Thr Gly Asp Arg Ile Arg Ser
35 40 45
Lys Val Glu Leu Thr Arg Tyr Leu Gly Pro Ala Cys Asp Leu Thr
50 55 60
Leu Phe Asp Phe Lys Gln Gly Ile Leu Cys Tyr Pro Ala Pro Lys
65 70 75
Ala His Pro Val Ala Val Ala Ser Lys Lys Arg Lys Lys Pro Ser
80 85 90
Arg Pro Ala Lys Thr Arg Lys Arg Gln Val Gly Pro Gln Ser Gly
95 100 105
Glu Val Arg Lys Glu Ala Pro Arg Asp Glu Thr Lys Ala Asp Thr
110 115 120
Asp Thr Ala Pro Ala Ser Phe Pro Ala Pro Gly Cys Cys Glu Asn
125 130 135
Cys Gly Ile Ser Phe Ser Gly Asp Gly Thr Gln Arg Gln Arg Leu
140 145 150
Lys Thr Leu Cys Lys Asp Cys Arg Ala Gln Arg Ile Ala Phe Asn
155 160 165
Arg Glu Gln Arg Met Phe Lys Arg Val Gly Cys Gly Glu Cys Ala
170 175 180
Ala Cys Gln Val Thr Glu Asp Cys Gly Ala Cys Ser Thr Cys Leu
185 190 195
Leu Gln Leu Pro His Asp Val Ala Ser Gly Leu Phe Cys Lys Cys
200 205 210
Glu Arg Arg Arg Cys Leu Arg Ile Val Glu Arg Ser Arg Gly Cys
215 220 225
Gly Val Cys Arg Gly Cys Gln Thr Gln Glu Asp Cys Gly His Cys
230 235 240
Pro Ile Cys Leu Arg Pro Pro Arg Pro Gly Leu Arg Arg Gln Trp
245 250 255

```

```

Lys Cys Val Gln Arg Arg Cys Leu Arg Gly Lys His Ala Arg Arg
260 265 270
Lys Gly Gly Cys Asp Ser Lys Met Ala Ala Arg Arg Arg Pro Gly
275 280 285
Ala Gln Pro Leu Pro Pro Pro Pro Pro Ser Gln Ser Pro Glu Pro
290 295 300
Thr Glu Pro His Pro Arg Ala Leu Ala Pro Ser Pro Pro Ala Glu
305 310 315
Phe Ile Tyr Tyr Cys Val Asp Glu Asp Glu Leu Gln Pro Tyr Thr
320 325 330
Asn Arg Arg Gln Asn Arg Lys Cys Gly Ala Cys Ala Ala Cys Leu
335 340 345
Arg Arg Met Asp Cys Gly Arg Cys Asp Phe Cys Cys Asp Lys Pro
350 355 360
Lys Phe Gly Gly Ser Asn Gln Lys Arg Gln Lys Cys Arg Trp Arg
365 370 375
Gln Cys Leu Gln Phe Ala Met Lys Arg Leu Leu Pro Ser Val Trp
380 385 390
Ser Glu Ser Glu Asp Gly Ala Gly Ser Pro Pro Pro Tyr Arg Arg
395 400 405

Arg Lys Arg Pro Ser Ser Ala Arg Arg His His Leu Gly Pro Thr
410 415 420
Leu Lys Pro Thr Leu Ala Thr Arg Thr Ala Gln Pro Asp His Thr
425 430 435
Gln Ala Pro Thr Lys Gln Glu Ala Gly Gly Gly Phe Val Leu Pro
440 445 450
Pro Pro Gly Thr Asp Leu Val Phe Leu Arg Glu Gly Ala Ser Ser
455 460 465
Pro Val Gln Val Pro Gly Pro Val Ala Ala Ser Thr Glu Ala Leu
470 475 480
Leu Gln Val Lys Gln Glu Lys Ala Asp Thr Gln Asp Glu Trp Thr
485 490 495
Pro Gly Thr Ala Val Leu Thr Ser Pro Val Leu Val Pro Gly Cys
500 505 510
Pro Ser Lys Ala Val Asp Pro Gly Leu Pro Ser Val Lys Gln Glu
515 520 525
Pro Pro Asp Pro Glu Glu Asp Lys Glu Glu Asn Lys Asp Asp Ser
530 535 540
Ala Ser Lys Leu Ala Pro Glu Glu Glu Ala Gly Gly Ala Gly Thr
545 550 555
Pro Val Ile Thr Glu Ile Phe Ser Leu Gly Gly Thr Arg Phe Arg
560 565 570
Asp Thr Ala Val Trp Leu Pro Arg Ser Lys Asp Leu Lys Lys Pro
575 580 585
Gly Ala Arg Lys Gln
590

```

<210> 13
 <211> 479
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1685494CD1

<400> 13
 Met Ala Thr Ala Leu Val Ser Ala His Ser Leu Ala Pro Leu Ser

1	5	10	15
Leu Lys Lys Glu Gly	Leu Arg Val Val Arg	Glu Asp His Tyr Ser	
20	25	30	
Thr Trp Glu Gln Gly	Phe Lys Leu Gln Gly	Asn Ser Lys Gly Leu	
35	40	45	
Gly Gln Glu Pro Leu	Cys Lys Gln Phe Arg	Gln Leu Arg Tyr Glu	
50	55	60	
Glu Thr Thr Gly Pro	Arg Glu Ala Leu Ser	Arg Leu Arg Glu Leu	
65	70	75	
Cys Gln Gln Trp Leu	Gln Pro Glu Thr His	Thr Lys Glu Gln Ile	
80	85	90	
Leu Glu Leu Leu Val	Leu Glu Gln Phe Leu	Ile Ile Leu Pro Lys	
95	100	105	
Glu Leu Gln Ala Arg	Val Gln Glu His His	Pro Glu Ser Arg Glu	
110	115	120	
Asp Val Val Val Val	Leu Glu Asp Leu Gln	Leu Asp Leu Gly Glu	
125	130	135	
Thr Gly Gln Gln Val	Asp Pro Asp Gln Pro	Lys Lys Gln Lys Ile	
140	145	150	
Leu Val Glu Glu Met	Ala Pro Leu Lys Gly	Val Gln Glu Gln Gln	
155	160	165	
Val Arg His Glu Cys	Glu Val Thr Lys Pro	Glu Lys Glu Lys Gly	
170	175	180	
Glu Glu Thr Arg Ile	Glu Asn Gly Lys Leu	Ile Val Val Thr Asp	
185	190	195	
Ser Cys Gly Arg Val	Glu Ser Ser Gly Lys	Ile Ser Glu Pro Met	
200	205	210	
Glu Ala His Asn Glu	Gly Ser Asn Leu Glu	Arg His Gln Ala Lys	
215	220	225	
Pro Lys Glu Lys Ile	Glu Tyr Lys Cys Ser	Glu Arg Glu Gln Arg	
230	235	240	
Phe Ile Gln His Leu	Asp Leu Ile Glu His	Ala Ser Thr His Thr	
245	250	255	
Gly Lys Lys Leu Cys	Glu Ser Asp Val Cys	Gln Ser Ser Ser Leu	
260	265	270	
Thr Gly His Lys Lys	Val Leu Ser Arg Glu	Lys Gly His Gln Cys	
275	280	285	
His Glu Cys Gly Lys	Ala Phe Gln Arg Ser	Ser His Leu Val Arg	
290	295	300	
His Gln Lys Ile His	Leu Gly Glu Lys Pro	Tyr Gln Cys Asn Glu	
305	310	315	
Cys Gly Lys Val Phe	Ser Gln Asn Ala Gly	Leu Leu Glu His Leu	
320	325	330	
Arg Ile His Thr Gly	Glu Lys Pro Tyr Leu	Cys Ile His Cys Gly	
335	340	345	
Lys Asn Phe Arg Arg	Ser Ser His Leu Asn	Arg His Gln Arg Ile	
350	355	360	
His Ser Gln Glu Glu	Pro Cys Glu Cys Lys	Glu Cys Gly Lys Thr	
365	370	375	
Phe Ser Gln Ala Leu	Leu Leu Leu Thr His	His Gln Arg Ile His Ser	
380	385	390	
His Ser Lys Ser His	Gln Cys Asn Glu Cys	Gly Lys Ala Phe Ser	
395	400	405	
Leu Thr Ser Asp Leu	Ile Arg His His Arg	Ile His Thr Gly Glu	
410	415	420	
Lys Pro Phe Lys Cys	Asn Ile Cys Gln Lys	Ala Phe Arg Leu Asn	
425	430	435	
Ser His Leu Ala Gln	His Val Arg Ile His	Asn Glu Glu Lys Pro	
440	445	450	
Tyr Gln Cys Ser Glu	Cys Gly Glu Ala Phe	Arg Gln Arg Ser Gly	
455	460	465	
Leu Phe Gln His Gln	Arg Tyr His His Lys	Asp Lys Leu Ala	

470

475

<210> 14
 <211> 433
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1730829CD1

<400> 14
 Met Glu Ala Val Tyr Leu Val Val Asn Gly Leu Gly Leu Val Leu
 1 5 10 15
 Asp Val Leu Thr Leu Val Leu Asp Leu Asn Phe Leu Leu Val Ser
 20 25 30
 Ser Leu Leu Ala Ser Leu Ala Trp Leu Leu Ala Phe Val Tyr Asn
 35 40 45
 Leu Pro His Thr Val Leu Thr Ser Leu Leu His Leu Gly Arg Gly
 50 55 60
 Val Leu Leu Ser Leu Leu Ala Leu Ile Glu Ala Val Val Arg Phe
 65 70 75
 Thr Cys Gly Gly Leu Gln Ala Leu Cys Thr Leu Leu Tyr Ser Cys
 80 85 90
 Cys Ser Gly Leu Glu Ser Leu Lys Leu Leu Gly His Leu Ala Ser
 95 100 105
 His Gly Ala Leu Arg Ser Arg Glu Ile Leu His Arg Gly Val Leu
 110 115 120
 Asn Val Val Ser Ser Gly His Ala Leu Leu Arg Gln Ala Cys Asp
 125 130 135
 Ile Cys Ala Ile Ala Met Ser Leu Val Ala Tyr Val Ile Asn Ser
 140 145 150
 Leu Val Asn Ile Cys Leu Ile Gly Thr Gln Asn Leu Phe Ser Leu
 155 160 165
 Val Leu Ala Leu Trp Asp Ala Val Thr Gly Pro Leu Trp Arg Met
 170 175 180
 Thr Asp Val Val Ala Ala Phe Leu Ala His Ile Ser Ser Ser Ala
 185 190 195
 Val Ala Met Ala Ile Leu Leu Trp Thr Pro Cys Gln Leu Ala Leu
 200 205 210
 Glu Leu Leu Ala Ser Ala Ala Arg Leu Leu Ala Ser Phe Val Leu
 215 220 225
 Val Asn Leu Thr Gly Leu Val Leu Leu Ala Cys Val Leu Ala Val
 230 235 240
 Thr Val Thr Val Leu His Pro Asp Phe Thr Leu Arg Leu Ala Thr
 245 250 255
 Gln Ala Leu Ser Gln Leu His Ala Arg Pro Ser Tyr His Arg Leu
 260 265 270
 Arg Glu Asp Val Met Arg Leu Ser Arg Leu Ala Leu Gly Ser Glu
 275 280 285
 Ala Trp Arg Arg Val Trp Ser Arg Ser Leu Gln Leu Ala Ser Trp
 290 295 300
 Pro Asn Arg Gly Gly Ala Pro Gly Ala Pro Gln Gly Asp Pro Met
 305 310 315
 Arg Val Phe Ser Val Arg Thr Arg Arg Gln Asp Thr Leu Pro Glu
 320 325 330
 Ala Gly Arg Arg Ser Glu Ala Glu Glu Glu Ala Arg Thr Ile
 335 340 345
 Arg Val Thr Pro Val Arg Gly Arg Glu Arg Leu Asn Glu Glu Glu

18/91

Pro	Pro	Gly	Gly	Gln	Asp	Pro	Trp	Lys	Leu	Leu	Lys	Glu	Gln	Glu	350	355	360
															365	370	375
Glu	Arg	Lys	Lys	Cys	Val	Ile	Cys	Gln	Asp	Gln	Ser	Lys	Thr	Val	380	385	390
Leu	Leu	Leu	Pro	Cys	Arg	His	Leu	Cys	Leu	Cys	Gln	Ala	Cys	Thr	395	400	405
Glu	Ile	Leu	Met	Arg	His	Pro	Val	Tyr	His	Arg	Asn	Cys	Pro	Leu	410	415	420
Cys	Arg	Arg	Gly	Ile	Leu	Gln	Thr	Leu	Asn	Val	Tyr	Leu			425	430	

<210> 15
 <211> 320
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc feature
 <223> Incyte ID No.: 1864641CD1

<400> 15

Met	Pro	Lys	Lys	Lys	Thr	Gly	Ala	Arg	Lys	Lys	Ala	Glu	Asn	Arg	1	5	10	15
Arg	Glu	Arg	Glu	Lys	Gln	Leu	Arg	Ala	Ser	Arg	Ser	Thr	Ile	Asp	20	25	30	35
Leu	Ala	Lys	His	Pro	Cys	Asn	Ala	Ser	Met	Glu	Cys	Asp	Lys	Cys	40	45	50	55
Gln	Arg	Arg	Gln	Lys	Asn	Arg	Ala	Phe	Cys	Tyr	Phe	Cys	Asn	Ser	60	65	70	75
Val	Gln	Lys	Leu	Pro	Ile	Cys	Ala	Gln	Cys	Gly	Lys	Thr	Lys	Cys	80	85	90	95
Met	Met	Lys	Ser	Ser	Asp	Cys	Val	Ile	Lys	His	Ala	Gly	Val	Tyr	100	105	110	115
Ser	Thr	Gly	Leu	Ala	Met	Val	Gly	Ala	Ile	Cys	Asp	Phe	Cys	Glu	120	125	130	135
Ala	Trp	Val	Cys	His	Gly	Arg	Lys	Cys	Leu	Ser	Thr	His	Ala	Cys	140	145	150	155
Ala	Cys	Pro	Leu	Thr	Asp	Ala	Glu	Cys	Val	Glu	Cys	Glu	Arg	Gly	160	165	170	175
Val	Trp	Asp	His	Gly	Gly	Arg	Ile	Phe	Ser	Cys	Ser	Phe	Cys	His	180	185	190	195
Asn	Phe	Leu	Cys	Glu	Asp	Asp	Gln	Phe	Glu	His	Gln	Ala	Ser	Cys	200	205	210	215
Gln	Val	Leu	Glu	Ala	Glu	Thr	Phe	Lys	Cys	Val	Ser	Cys	Asn	Arg	220	225	230	235
Leu	Gly	Gln	His	Ser	Cys	Leu	Arg	Cys	Lys	Ala	Cys	Phe	Cys	Asp	240	245	250	255
Asp	His	Thr	Arg	Ser	Lys	Val	Phe	Lys	Gln	Glu	Lys	Gly	Lys	Gln	260	265	270	
Pro	Pro	Cys	Pro	Lys	Cys	Gly	His	Glu	Thr	Gln	Glu	Thr	Lys	Asp				
Leu	Ser	Met	Ser	Thr	Arg	Ser	Leu	Lys	Phe	Gly	Arg	Gln	Thr	Gly				
Gly	Glu	Glu	Gly	Asp	Gly	Ala	Ser	Gly	Tyr	Asp	Ala	Tyr	Trp	Lys				
Asn	Leu	Ser	Ser	Asp	Lys	Tyr	Gly	Asp	Thr	Ser	Tyr	His	Asp	Glu				
Glu	Glu	Asp	Glu	Tyr	Glu	Ala	Glu	Asp	Asp	Glu	Glu	Glu	Glu	Asp				

	275		280		285
Glu Gly Arg Lys	Asp Ser Asp Thr Glu	Ser Ser Asp Leu Phe	Thr		
	290		295		300
Asn Leu Asn Leu	Gly Arg Thr Tyr Ala	Ser Gly Tyr Ala His	Tyr		
	305		310		315
Glu Glu Gln Glu	Asn				
	320				

<210> 16
 <211> 179
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2444604CD1

<400> 16

Met Ala Ala Gly	Phe Phe Gln Pro	Phe Met Ser Pro Arg	Phe Pro
1	5	10	15
Gly Gly Pro Arg	Pro Thr Leu Arg Met	Pro Ser Gln Pro Pro	Ala
	20	25	30
Cys Leu Pro Gly	Ser Gln Pro Leu Leu Pro	Gly Ala Met Glu	Pro
	35	40	45
Ser Pro Arg Ala	Gln Gly His Pro Ser Met	Gly Gly Pro Met	Gln
	50	55	60
Arg Val Thr Pro	Pro Arg Gly Met Ala Ser	Val Gly Pro Gln	Ser
	65	70	75
Tyr Gly Gly Gly	Met Arg Pro Pro Pro	Asn Ser Leu Ala Gly	Pro
	80	85	90
Gly Leu Pro Ala	Met Asn Met Gly Pro	Gly Val Arg Gly Pro	Trp
	95	100	105
Ala Ser Pro Ser	Gly Asn Ser Ile Pro Tyr	Ser Ser Ser Ser	Pro
	110	115	120
Gly Ser Tyr Thr	Gly Pro Pro Gly Gly	Gly Gly Pro Pro Gly	Thr
	125	130	135
Pro Ile Met Pro	Ser Pro Gly Asp Ser	Thr Asn Ser Ser Glu	Asn
	140	145	150
Met Tyr Thr Ile	Met Asn Pro Ile Gly	Gln Gly Ala Gly Arg	Ala
	155	160	165
Asn Phe Pro Leu	Gly Pro Gly Pro Glu	Gly Pro Trp Pro Pro	
	170	175	

<210> 17
 <211> 494
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2445008CD1

<400> 17

Met Gly Arg Lys	Lys Lys Lys Gln Leu	Lys Pro Trp Cys Trp	Tyr
1	5	10	15
Cys Asn Arg Asp	Phe Asp Asp Glu Lys	Ile Leu Ile Gln His	Gln

	20		25		30
Lys Ala Lys His Phe	Lys Cys His Ile Cys	His Lys Lys Leu Tyr			
	35		40		45
Thr Gly Pro Gly Leu	Ala Ile His Cys Met	Gln Val His Lys Glu			
	50		55		60
Thr Ile Asp Ala Val	Pro Asn Ala Ile Pro	Gly Arg Thr Asp Ile			
	65		70		75
Glu Leu Glu Ile Tyr	Gly Met Glu Gly Ile	Pro Glu Lys Asp Met			
	80		85		90
Asp Glu Arg Arg Arg	Leu Leu Glu Gln Lys	Thr Gln Glu Ser Gln			
	95		100		105
Lys Lys Lys Gln Gln	Asp Asp Ser Asp Glu	Tyr Asp Asp Asp Asp			
	110		115		120
Ser Ala Ala Ser Thr	Ser Phe Gln Pro Gln	Pro Val Gln Pro Gln			
	125		130		135
Gln Gly Tyr Ile Pro	Pro Met Ala Gln Pro	Gly Leu Pro Pro Val			
	140		145		150
Pro Gly Ala Pro Gly	Met Pro Pro Gly Ile	Pro Pro Leu Met Pro			
	155		160		165
Gly Val Pro Pro Leu	Met Pro Gly Met Pro	Pro Val Met Pro Gly			
	170		175		180
Met Pro Pro Gly Leu	His His Gln Arg Lys	Tyr Thr Gln Ser Phe			
	185		190		195
Cys Gly Glu Asn Ile	Met Met Pro Met Gly	Gly Met Met Pro Pro			
	200		205		210
Gly Pro Gly Ile Pro	Pro Pro Leu Met Pro	Gly Met Pro Pro Gly			
	215		220		225
Pro Pro Pro Val Pro	Arg Pro Gly Ile Pro	Pro Met Thr Gln Ala			
	230		235		240
Gln Ala Val Ser Ala	Pro Gly Ile Leu Asn	Arg Pro Pro Ala Pro			
	245		250		255
Thr Ala Thr Val Pro	Ala Pro Gln Pro Pro	Val Thr Lys Pro Leu			
	260		265		270
Phe Pro Ser Ala Gly	Gln Met Gly Thr Pro	Val Thr Ser Ser Ser			
	275		280		285
Thr Ala Ser Ser Asn	Ser Glu Ser Leu Ser	Ala Ser Ser Lys Ala			
	290		295		300
Leu Phe Pro Ser Thr	Ala Gln Ala Gln Ala	Ala Val Gln Gly Pro			
	305		310		315
Val Gly Thr Asp Phe	Lys Pro Leu Asn Ser	Thr Pro Ala Thr Thr			
	320		325		330
Thr Glu Pro Pro Lys	Pro Thr Phe Pro Ala	Tyr Thr Gln Ser Thr			
	335		340		345
Ala Ser Thr Thr Ser	Thr Thr Asn Ser Thr	Ala Ala Lys Pro Ala			
	350		355		360
Ala Ser Ile Thr Ser	Lys Pro Ala Thr Leu	Thr Thr Thr Ser Ala			
	365		370		375
Thr Ser Lys Leu Ile	His Pro Asp Glu Asp	Ile Ser Leu Glu Glu			
	380		385		390
Arg Arg Ala Gln Leu	Pro Lys Tyr Gln Arg	Asn Leu Pro Arg Pro			
	395		400		405
Gly Gln Ala Pro Ile	Gly Asn Pro Pro Val	Gly Pro Ile Gly Gly			
	410		415		420
Met Met Pro Pro Gln	Pro Gly Ile Pro Gln	Gln Gln Gly Met Arg			
	425		430		435
Pro Pro Met Pro Pro	His Gly Gln Tyr Gly	Gly His His Gln Gly			
	440		445		450
Met Pro Gly Tyr Leu	Pro Gly Ala Met Pro	Pro Tyr Gly Gln Gly			
	455		460		465
Pro Pro Met Val Pro	Pro Tyr Gln Gly Gly	Pro Pro Arg Pro Pro			
	470		475		480

Met Gly Met Arg Pro Pro Val Met Ser Gln Gly Gly Arg Tyr
 485 490

<210> 18
 <211> 401
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2572462CD1

<400> 18

Met	Ala	Ser	Ser	Pro	Arg	Pro	Lys	Met	Asp	Ala	Ile	Leu	Thr	Glu	
1				5					10					15	
Ala	Ile	Lys	Ala	Cys	Phe	Gln	Lys	Ser	Gly	Ala	Ser	Val	Val	Ala	
				20					25					30	
Ile	Arg	Lys	Tyr	Ile	Ile	His	Lys	Tyr	Pro	Ser	Leu	Glu	Leu	Glu	
				35					40					45	
Arg	Arg	Gly	Tyr	Leu	Leu	Lys	Gln	Ala	Leu	Lys	Arg	Glu	Leu	Asn	
				50					55					60	
Arg	Gly	Val	Ile	Lys	Gln	Val	Lys	Gly	Lys	Gly	Ala	Ser	Gly	Ser	
				65					70					75	
Phe	Val	Val	Val	Gln	Lys	Ser	Arg	Lys	Thr	Pro	Gln	Lys	Ser	Arg	
				80					85					90	
Asn	Arg	Lys	Asn	Arg	Ser	Ser	Ala	Val	Asp	Pro	Glu	Pro	Gln	Val	
				95					100					105	
Lys	Leu	Glu	Asp	Val	Leu	Pro	Leu	Ala	Phe	Thr	Arg	Leu	Cys	Glu	
				110					115					120	
Pro	Lys	Glu	Ala	Ser	Tyr	Ser	Leu	Ile	Arg	Lys	Tyr	Val	Ser	Gln	
				125					130					135	
Tyr	Tyr	Pro	Lys	Leu	Arg	Val	Asp	Ile	Arg	Pro	Gln	Leu	Leu	Lys	
				140					145					150	
Asn	Ala	Leu	Gln	Arg	Ala	Val	Glu	Arg	Gly	Gln	Leu	Glu	Gln	Ile	
				155					160					165	
Thr	Gly	Lys	Gly	Ala	Ser	Gly	Thr	Phe	Gln	Leu	Lys	Lys	Ser	Gly	
				170					175					180	
Glu	Lys	Pro	Leu	Leu	Gly	Gly	Ser	Leu	Met	Glu	Tyr	Ala	Ile	Leu	
				185					190					195	
Ser	Ala	Ile	Ala	Ala	Met	Asn	Glu	Pro	Lys	Thr	Cys	Ser	Thr	Thr	
				200					205					210	
Ala	Leu	Lys	Lys	Tyr	Val	Leu	Glu	Asn	His	Pro	Gly	Thr	Asn	Ser	
				215					220					225	
Asn	Tyr	Gln	Met	His	Leu	Leu	Lys	Lys	Thr	Leu	Gln	Lys	Cys	Glu	
				230					235					240	
Lys	Asn	Gly	Trp	Met	Glu	Gln	Ile	Ser	Gly	Lys	Gly	Phe	Ser	Gly	
				245					250					255	
Thr	Phe	Gln	Leu	Cys	Phe	Pro	Tyr	Tyr	Pro	Ser	Pro	Gly	Val	Leu	
				260					265					270	
Phe	Pro	Lys	Lys	Glu	Pro	Asp	Asp	Ser	Arg	Asp	Glu	Asp	Glu	Asp	
				275					280					285	
Glu	Asp	Glu	Ser	Ser	Glu	Glu	Asp	Ser	Glu	Asp	Glu	Glu	Pro	Pro	
				290					295					300	
Pro	Lys	Arg	Arg	Leu	Gln	Lys	Lys	Thr	Pro	Ala	Lys	Ser	Pro	Gly	
				305					310					315	
Lys	Ala	Ala	Ser	Val	Lys	Gln	Arg	Gly	Ser	Lys	Pro	Ala	Pro	Lys	
				320					325					330	
Val	Ser	Ala	Ala	Gln	Arg	Gly	Lys	Ala	Arg	Pro	Leu	Pro	Lys	Lys	
				335					340					345	

Ala	Pro	Pro	Lys	Ala	Lys	Thr	Pro	Ala	Lys	Lys	Thr	Arg	Pro	Ser
			350						355					360
Ser	Thr	Val	Ile	Lys	Lys	Pro	Ser	Gly	Gly	Ser	Ser	Lys	Lys	Pro
			365						370					375
Ala	Thr	Ser	Ala	Arg	Lys	Glu	Val	Lys	Leu	Pro	Gly	Lys	Gly	Lys
			380						385					390
Ser	Thr	Met	Lys	Lys	Ser	Phe	Arg	Val	Lys	Lys				
			395						400					

<210> 19
 <211> 264
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2572892CD1

<400> 19

Met	Pro	Arg	Ser	Phe	Leu	Val	Arg	Lys	Pro	Ser	Asp	Pro	Asn	Arg
1				5					10					15
Lys	Pro	Asn	Tyr	Ser	Glu	Leu	Gln	Asp	Ser	Asn	Pro	Glu	Phe	Thr
			20						25					30
Phe	Gln	Gln	Pro	Tyr	Asp	Gln	Ala	His	Leu	Leu	Ala	Ala	Ile	Pro
			35						40					45
Pro	Pro	Glu	Ile	Leu	Asn	Pro	Thr	Ala	Ser	Leu	Pro	Met	Leu	Ile
			50						55					60
Trp	Asp	Ser	Val	Leu	Ala	Pro	Gln	Ala	Gln	Pro	Ile	Ala	Trp	Ala
			65						70					75
Ser	Leu	Arg	Leu	Gln	Glu	Ser	Pro	Arg	Val	Ala	Glu	Leu	Thr	Ser
			80						85					90
Leu	Ser	Asp	Glu	Asp	Ser	Gly	Lys	Gly	Ser	Gln	Pro	Pro	Ser	Pro
			95						100					105
Pro	Ser	Pro	Ala	Pro	Ser	Ser	Phe	Ser	Ser	Thr	Ser	Ala	Ser	Ser
			110						115					120
Leu	Glu	Ala	Glu	Ala	Tyr	Ala	Ala	Phe	Pro	Gly	Leu	Gly	Gln	Val
			125						130					135
Pro	Lys	Gln	Leu	Ala	Gln	Leu	Ser	Glu	Ala	Lys	Asp	Leu	Gln	Ala
			140						145					150
Arg	Lys	Ala	Phe	Asn	Cys	Lys	Tyr	Cys	Asn	Lys	Glu	Tyr	Leu	Ser
			155						160					165
Leu	Gly	Ala	Leu	Lys	Met	His	Ile	Arg	Ser	His	Thr	Leu	Pro	Cys
			170						175					180
Val	Cys	Gly	Thr	Cys	Gly	Lys	Ala	Phe	Ser	Arg	Pro	Trp	Leu	Leu
			185						190					195
Gln	Gly	His	Val	Arg	Thr	His	Thr	Gly	Glu	Lys	Pro	Phe	Ser	Cys
			200						205					210
Pro	His	Cys	Ser	Arg	Ala	Phe	Ala	Asp	Arg	Ser	Asn	Leu	Arg	Ala
			215						220					225
His	Leu	Gln	Thr	His	Ser	Asp	Val	Lys	Lys	Tyr	Gln	Cys	Gln	Ala
			230						235					240
Cys	Ala	Arg	Thr	Phe	Ser	Arg	Met	Ser	Leu	Leu	His	Lys	His	Gln
			245						250					255
Glu	Ser	Gly	Cys	Ser	Gly	Cys	Pro	Arg						
			260											

<210> 20
 <211> 153
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2785674CD1

<400> 20
 Met Thr Lys Ile Lys Ala Asp Pro Asp Gly Pro Glu Ala Gln Ala
 1 5 10 15
 Glu Ala Cys Ser Gly Glu Arg Thr Tyr Gln Glu Leu Leu Val Asn
 20 25 30
 Gln Asn Pro Ile Ala Gln Pro Leu Ala Ser Arg Arg Leu Thr Arg
 35 40 45
 Lys Leu Tyr Lys Cys Ile Lys Lys Ala Val Lys Gln Lys Gln Ile
 50 55 60
 Arg Arg Gly Val Lys Glu Val Gln Lys Phe Val Asn Lys Gly Glu
 65 70 75
 Lys Gly Ile Met Val Leu Ala Gly Asp Thr Leu Pro Ile Glu Val
 80 85 90
 Tyr Cys His Leu Pro Val Met Cys Glu Asp Arg Asn Leu Pro Tyr
 95 100 105
 Val Tyr Ile Pro Ser Lys Thr Asp Leu Gly Ala Ala Ala Gly Ser
 110 115 120
 Lys Arg Pro Thr Cys Val Ile Met Val Lys Pro His Glu Glu Tyr
 125 130 135
 Gln Glu Ala Tyr Asp Glu Cys Leu Glu Glu Val Gln Ser Leu Pro
 140 145 150
 Leu Pro Leu

<210> 21
 <211> 243
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2797479CD1

<400> 21
 Met Gly Asp Asp Ile Ser Thr His Ile Ala Pro Lys Glu Leu Arg
 1 5 10 15
 His Lys His Pro Ser Ser Val Asp Glu Val Ala Gln Val Val Lys
 20 25 30
 Gln Leu Arg Ile Ile Leu Gln Gln Gln Val Arg Pro Gly Glu Ser
 35 40 45
 Thr Val Leu Ala Leu Arg Pro Asn Val Gln Gln Ile Glu Ala Pro
 50 55 60
 Asp Val Ser Arg Asp Pro Arg Val Leu Gly Phe Asp Phe Pro Gly
 65 70 75
 Ser Ala Arg Ile His Glu Gly Thr His Thr Leu Glu Lys Pro Tyr
 80 85 90
 Glu Cys Lys Gln Cys Gly Lys Leu Leu Ser His Arg Ser Ser Phe
 95 100 105
 Arg Arg His Met Ala His Thr Gly Asp Gly Pro His Lys Cys
 110 115 120
 Thr Val Cys Gly Lys Ala Phe Asp Ser Pro Ser Val Phe Gln Arg

	125		130		135
His Glu Arg Thr	His Thr Gly Glu Lys	Pro Tyr Glu Cys Lys	Gln		
	140		145		150
Cys Gly Lys Ala	Phe Arg Thr Ser Ser	Ser Leu Arg Lys His	Glu		
	155		160		165
Thr Thr His Thr	Gly Glu Gln Pro Tyr	Lys Cys Lys Cys Gly	Lys		
	170		175		180
Ala Phe Ser Asp	Leu Phe Ser Phe Gln	Ser His Glu Thr Thr	His		
	185		190		195
Ser Glu Glu Glu	Pro Tyr Glu Cys Lys	Glu Cys Gly Lys Ala	Phe		
	200		205		210
Ser Ser Phe Lys	Tyr Phe Cys Arg His	Glu Arg Thr His Ser	Glu		
	215		220		225
Glu Lys Ser Tyr	Glu Cys Gln Ile Cys	Gly Lys Leu Ser Val	Val		
	230		235		240
Ser Val Thr					

<210> 22
 <211> 485
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2960640CD1

<400> 22

Met Arg Asp Asn Arg Ala Val Ser Leu Cys Gln Gln Glu Trp Met		
1	5	10
Cys Pro Gly Pro Ala Gln Arg Ala Leu Tyr Arg Gly Ala Thr Gln		
	20	25
Arg Lys Asp Ser His Val Ser Leu Ala Thr Gly Val Pro Trp Gly		
	35	40
Tyr Glu Glu Thr Lys Thr Leu Leu Ala Ile Leu Ser Ser Ser Gln		
	50	55
Phe Tyr Gly Lys Leu Gln Thr Cys Gln Gln Asn Ser Gln Ile Tyr		
	65	70
Arg Ala Met Ala Glu Gly Leu Trp Glu Gln Gly Phe Leu Arg Thr		
	80	85
Pro Glu Gln Cys Arg Thr Lys Phe Lys Ser Leu Gln Leu Ser Tyr		
	95	100
Arg Lys Val Arg Arg Gly Arg Val Pro Glu Pro Cys Ile Phe Tyr		
	110	115
Glu Glu Met Asn Ala Leu Ser Gly Ser Trp Ala Ser Ala Pro Pro		
	125	130
Met Ala Ser Asp Ala Val Pro Gly Gln Glu Gly Ser Asp Ile Glu		
	140	145
Ala Gly Glu Leu Asn His Gln Asn Gly Glu Pro Thr Glu Val Glu		
	155	160
Asp Gly Thr Val Asp Gly Ala Asp Arg Asp Glu Lys Asp Phe Arg		
	170	175
Asn Pro Gly Gln Glu Val Arg Lys Leu Asp Leu Pro Val Leu Phe		
	185	190
Pro Asn Arg Leu Gly Phe Glu Phe Lys Asn Glu Ile Lys Lys Glu		
	200	205
Asn Leu Lys Trp Asp Asp Ser Glu Glu Val Glu Ile Asn Lys Ala		
	215	220
Leu Gln Arg Lys Ser Arg Gly Val Tyr Trp His Ser Glu Leu Gln		

Lys Gly Leu Glu Ser	230	Arg Arg Gln Cys Arg Asn	235	240
	245	Pro Ser Gln Glu Lys Met	250	255
Ser Pro Gly Glu Ser	260	Lys Ala Cys Thr His Ile	265	270
	275	Val His Ser Pro His Lys	280	285
Ser His Gln Ser Phe	290	Gln Cys Pro Glu Cys Gly	295	300
	305	Val Arg His Gln Arg Ile	310	315
Leu Cys Gly Lys Asn	320		325	330
Pro Ala Leu Lys Leu				
Lys Thr Phe Ser Arg				
His Thr Gly Glu Lys	335	Glu Cys Gly Lys Gly	340	345
	350	His Leu Arg Thr His Thr	355	360
Phe Ser Glu Arg Ser	365	Gly Lys Ser Phe Asn	370	375
	380	Thr His Thr Gly Glu	385	390
Gly Glu Arg Pro Tyr	395	Arg Phe Asn Asn Ser	400	405
	410	His Thr Gly Glu Ser Pro	415	420
Gln Ser Ser Ser Leu	425	Phe Asn Asn Ser Ser His	430	435
	440	Gly Glu Lys Pro Tyr Arg	445	450
Lys Pro Tyr Gln Cys	455	Lys Asn Ser Ala Leu Thr	460	465
	470	Val Leu Ser Ser Gln Glu	475	480
Ser Gln Phe Ser Ala				
Tyr Lys Cys Ala Val				
Phe Ser Ala His Arg				
Cys Ser His Cys Glu				
Arg His Gln Thr Val				
Gly Arg Asp Ala Leu				

<210> 23
 <211> 160
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3454051CD1

Met Ser Trp Thr Cys	1	Pro Arg Cys Gln Gln	10	Pro Val Phe Phe Ala	15
	5	Trp His Arg Phe Cys Leu	20		25
Glu Lys Val Ser Ser	20	Leu Ser Pro Gly Gly His	25		30
	25	Ala	30		35
Lys Cys Glu Arg Cys	35	His Ser Ile Leu Ser	40		45
	40	Pro Cys Tyr Gly Ala	45		50
Glu His Asn Gly Arg	50	Pro Tyr Cys His Lys	55		60
	55	Met Lys Thr Phe Thr Gly	60		65
Leu Phe Gly Pro Arg	65	Gly Pro Pro His	70		75
	70	Glu Pro Val Tyr Phe Ala	75		80
Glu Thr Ser Leu Cys	80		85		90

Glu	Lys	Val	Met	Ser	Leu	Gly	Arg	Asn	Trp	His	Arg	Pro	Cys	Leu	
				95					100					105	
Arg	Cys	Gln	Arg	Cys	His	Lys	Thr	Leu	Thr	Ala	Gly	Ser	His	Ala	
				110					115					120	
Glu	His	Asp	Gly	Val	Pro	Tyr	Cys	His	Val	Pro	Cys	Tyr	Gly	Tyr	
				125					130					135	
Leu	Phe	Gly	Pro	Lys	Gly	Val	Asn	Ile	Gly	Asp	Val	Gly	Cys	Tyr	
				140					145					150	
Ile	Tyr	Asp	Pro	Val	Lys	Ile	Lys	Phe	Lys						
				155					160						

<210> 24
 <211> 511
 <212> PRT
 <213> Homo sapiens

<220>
 <221>
 <223> Incyte ID No.: 3510640CD1

<400> 24

Met	Gln	Glu	Leu	Tyr	Ser	Thr	Pro	Ala	Ser	Arg	Leu	Asp	Ser	Phe	
1				5					10					15	
Val	Ala	Gln	Trp	Leu	Gln	Pro	His	Arg	Glu	Trp	Lys	Glu	Glu	Val	
				20					25					30	
Leu	Asp	Ala	Val	Arg	Thr	Val	Glu	Glu	Phe	Leu	Arg	Gln	Glu	His	
				35					40					45	
Phe	Gln	Gly	Lys	Arg	Gly	Leu	Asp	Gln	Asp	Val	Arg	Val	Leu	Lys	
				50					55					60	
Val	Val	Lys	Val	Gly	Ser	Phe	Gly	Asn	Gly	Thr	Val	Leu	Arg	Ser	
				65					70					75	
Thr	Arg	Glu	Val	Glu	Leu	Val	Ala	Phe	Leu	Ser	Cys	Phe	His	Ser	
				80					85					90	
Phe	Gln	Glu	Ala	Ala	Lys	His	His	Lys	Asp	Val	Leu	Arg	Leu	Ile	
				95					100					105	
Trp	Lys	Thr	Met	Trp	Gln	Ser	Gln	Asp	Leu	Leu	Asp	Leu	Gly	Leu	
				110					115					120	
Glu	Asp	Leu	Arg	Met	Glu	Gln	Arg	Val	Pro	Asp	Ala	Leu	Val	Phe	
				125					130					135	
Thr	Ile	Gln	Thr	Arg	Gly	Thr	Ala	Glu	Pro	Ile	Thr	Val	Thr	Ile	
				140					145					150	
Val	Pro	Ala	Tyr	Arg	Ala	Leu	Gly	Pro	Ser	Leu	Pro	Asn	Ser	Gln	
				155					160					165	
Pro	Pro	Pro	Glu	Val	Tyr	Val	Ser	Leu	Ile	Lys	Ala	Cys	Gly	Gly	
				170					175					180	
Pro	Gly	Asn	Phe	Cys	Pro	Phe	Phe	Ser	Glu	Leu	Gln	Arg	Asn	Phe	
				185					190					195	
Val	Lys	His	Arg	Pro	Thr	Lys	Leu	Lys	Ser	Leu	Leu	Arg	Leu	Val	
				200					205					210	
Lys	His	Trp	Tyr	Gln	Gln	Tyr	Val	Lys	Ala	Arg	Ser	Pro	Arg	Ala	
				215					220					225	
Asn	Leu	Pro	Pro	Leu	Tyr	Ala	Leu	Glu	Leu	Leu	Thr	Ile	Tyr	Ala	
				230					235					240	
Trp	Glu	Met	Gly	Thr	Glu	Glu	Asp	Glu	Asn	Phe	Met	Leu	Asp	Glu	
				245					250					255	
Gly	Phe	Thr	Thr	Val	Met	Asp	Leu	Leu	Leu	Glu	Tyr	Glu	Val	Ile	
				260					265					270	
Cys	Ile	Tyr	Trp	Thr	Lys	Tyr	Tyr	Thr	Leu	His	Asn	Ala	Ile	Ile	

Glu Asp Cys Val	275	Lys Glu Arg Pro Ile	285
Leu Asp Pro Ala	290	Val Ala Glu Gly Tyr	300
Trp Asp Ile Val	305	Gln Cys Leu Lys Gln	315
Cys Cys Tyr Asp	320	Ile Ser Ser Trp Asn	330
Lys Arg Ala Arg	335	Val Glu Gln Arg Gly	345
Pro Asp Phe Asn	350	Tyr Glu Pro Ile Arg	360
Val Lys Glu Lys	365	Gly Tyr Ser Gly Leu	375
Arg Leu Ser Phe	380	Glu Arg Gln Leu Leu	390
Ser Arg Cys Ser	395	Ile Phe Ser His Thr	405
Ile Tyr Leu Leu	410	Glu Ile Gln Val Phe	420
Lys Asn Pro Asp	425	Tyr Ala Ile Asn Pro	435
Ser Phe Ile Leu	440	Ile Glu Asp Gln Gln	450
	455		465
Leu Pro Lys Lys	470	Phe Gln Gly Gln Val	480
Gln Asp Trp Leu	485	Gly Ile Gln Asp Ser	495
Thr Leu Ile Leu	500	Glu Ala Leu Phe Pro	510
Ser			

<210> 25
 <211> 310
 <212> PRT
 <213> Homo sapiens

<220> -
 <223> Incyte ID No.: 3815083CD1

Met Arg Pro Leu	1	Gln Ile Val Pro Ser	10	Leu Ile Ser Gln	15
Tyr Cys Gly Leu	20	Lys Pro Pro Ala Ser	25	Thr Arg Asn Gln	30
Leu Lys Met Ala	35	Arg Pro Ser Ser Ser	40	Met Ala Asp Phe	45
Phe Phe Ala Lys	50	Ala Lys His Ile Val	55	Ile Ile Ser Gly	60
Val Ser Ala Glu	65	Ser Gly Val Pro Thr	70	Phe Arg Gly Ala	75
Tyr Trp Arg Lys	80	Trp Gln Ala Gln Asp	85	Leu Ala Thr Pro	90
Phe Ala His Asn	95	Pro Ser Arg Val Trp	100	Glu Phe Tyr His	105
Arg Glu Val Met	110	Gly Ser Lys Glu Pro	115	Asn Ala Gly His	120


```

Ile Ala Glu Cys Glu Thr Arg Leu Gly Lys Gln Gly Arg Arg Val
125 130 135
Val Val Ile Thr Gln Asn Ile Asp Glu Leu His Arg Lys Ala Gly
140 145 150
Thr Lys Asn Leu Leu Glu Ile His Gly Ser Leu Phe Lys Thr Arg
155 160 165
Cys Thr Ser Cys Gly Val Val Ala Glu Asn Tyr Lys Ser Pro Ile
170 175 180
Cys Pro Ala Leu Ser Gly Lys Gly Ala Pro Glu Pro Gly Thr Gln
185 190 195
Asp Ala Ser Ile Pro Val Glu Lys Leu Pro Arg Cys Glu Glu Ala
200 205 210
Gly Cys Gly Gly Leu Leu Arg Pro His Val Val Trp Phe Gly Glu
215 220 225
Asn Leu Asp Pro Ala Ile Leu Glu Glu Val Asp Arg Glu Leu Ala
230 235 240
His Cys Asp Leu Cys Leu Val Val Gly Thr Ser Ser Val Val Tyr
245 250 255
Pro Ala Ala Met Phe Ala Pro Gln Val Ala Ala Arg Gly Val Pro
260 265 270
Val Ala Glu Phe Asn Thr Glu Thr Thr Pro Ala Thr Asn Arg Phe
275 280 285
Arg Phe His Phe Gln Gly Pro Cys Gly Thr Thr Leu Pro Glu Ala
290 295 300
Leu Ala Cys His Glu Asn Glu Thr Val Ser
305 310

```

<210> 26
 <211> 331
 <212> PRT
 <213> Homo sapiens

<220> -
 <221> misc-feature
 <223> Incyte ID No.: 3988457CD1

```

<400> 26
Met Ala Ile Asn Arg Phe Arg Leu Glu Asn Asp Leu Glu Glu Leu
1 5 10 15
Ala Leu Tyr Gln Ile Gln Leu Leu Lys Asp Leu Arg His Thr Glu
20 25 30
Asn Glu Glu Asp Lys Val Ser Ser Ser Ser Phe Arg Gln Arg Met
35 40 45
Leu Gly Asn Leu Leu Arg Pro Pro Tyr Glu Arg Pro Glu Leu Pro
50 55 60
Thr Cys Leu Tyr Val Ile Gly Leu Thr Gly Ile Ser Gly Ser Gly
65 70 75
Lys Ser Ser Ile Ala Gln Arg Leu Lys Gly Leu Gly Ala Phe Val
80 85 90
Ile Asp Ser Asp His Leu Gly His Arg Ala Tyr Ala Pro Gly Gly
95 100 105
Pro Ala Tyr Gln Pro Val Val Glu Ala Phe Gly Thr Asp Ile Leu
110 115 120
His Lys Asp Gly Ile Ile Asn Arg Lys Val Leu Gly Ser Arg Val
125 130 135
Phe Gly Asn Lys Lys Gln Leu Lys Ile Leu Thr Asp Ile Met Trp
140 145 150
Pro Ile Ile Ala Lys Leu Ala Arg Glu Glu Met Asp Arg Ala Val
155 160 165
Ala Glu Gly Lys Arg Val Cys Val Ile Asp Ala Ala Val Leu Leu
170 175 180

```

Glu	Ala	Gly	Trp	Gln	Asn	Leu	Val	His	Glu	Val	Trp	Thr	Ala	Val	
				185					190						195
Ile	Pro	Glu	Thr	Glu	Ala	Val	Arg	Arg	Ile	Val	Glu	Arg	Asp	Gly	
				200					205						210
Leu	Ser	Glu	Ala	Ala	Ala	Gln	Ser	Arg	Leu	Gln	Ser	Gln	Met	Ser	
				215					220						225
Gly	Gln	Gln	Leu	Val	Glu	Gln	Ser	His	Val	Val	Leu	Ser	Ser	Pro	
				230					235						240
Cys	Gly	Ser	Arg	Ile	Ser	Pro	Asn	Ala	Arg	Trp	Arg	Lys	Pro	Gly	
				245					250						255
Pro	Ser	Cys	Arg	Ser	Ala	Phe	Pro	Arg	Leu	Ile	Arg	Pro	Ser	Thr	
				260					265						270
Glu	Lys	Phe	Ser	Val	Gly	Pro	Asp	Trp	Leu	Leu	Glu	Leu	Thr	Ser	
				275					280						285
Asp	Pro	Val	Val	Arg	Arg	Asn	Gly	Gly	Leu	Asp	Ala	His	Pro	Gly	
				290					295						300
Ser	Gly	Pro	Glu	Val	Gln	Ala	Ile	Leu	Cys	Arg	Thr	Trp	Pro	Gly	
				305					310						315
Leu	Val	Asp	Thr	Gly	Ser	Leu	Pro	Asn	Thr	Leu	Val	Phe	Gly	Gln	
				320					325						330

His

<210> 27
 <211> 200
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 131890CD1

<400> 27

Met	Met	Thr	Ala	Glu	Ser	Arg	Glu	Ala	Thr	Gly	Leu	Ser	Pro	Gln	
1				5					10					15	
Ala	Ala	Gln	Glu	Lys	Asp	Gly	Ile	Val	Ile	Val	Lys	Val	Glu	Glu	
				20					25					30	
Glu	Asp	Glu	Glu	Asp	His	Met	Trp	Gly	Gln	Asp	Ser	Thr	Leu	Gln	
				35					40					45	
Asp	Thr	Pro	Pro	Pro	Asp	Pro	Glu	Ile	Phe	Arg	Gln	Arg	Phe	Arg	
				50					55					60	
Arg	Phe	Cys	Tyr	Gln	Asn	Thr	Phe	Gly	Pro	Arg	Glu	Ala	Leu	Ser	
				65					70					75	
Arg	Leu	Lys	Glu	Leu	Cys	His	Gln	Trp	Leu	Arg	Pro	Glu	Ile	Asn	
				80					85					90	
Thr	Lys	Glu	Gln	Ile	Leu	Glu	Leu	Leu	Val	Leu	Glu	Gln	Phe	Leu	
				95					100					105	
Ser	Ile	Leu	Pro	Lys	Glu	Leu	Gln	Val	Trp	Leu	Gln	Glu	Tyr	Arg	
				110					115					120	
Pro	Asp	Ser	Gly	Glu	Glu	Ala	Val	Thr	Leu	Leu	Glu	Asp	Leu	Glu	
				125					130					135	
Leu	Asp	Leu	Ser	Gly	Gln	Gln	Val	Pro	Gly	Gln	Val	His	Gly	Pro	
				140					145					150	
Glu	Met	Leu	Ala	Arg	Gly	Met	Val	Pro	Leu	Asp	Pro	Val	Gln	Glu	
				155					160					165	
Ser	Ser	Ser	Phe	Asp	Leu	His	His	Glu	Ala	Thr	Gln	Ser	His	Phe	
				170					175					180	
Lys	His	Ser	Ser	Arg	Lys	Pro	Arg	Leu	Leu	Gln	Ser	Arg	Gly	Lys	
				185					190					195	

Lys Gln Gly Phe Ile

200

<210> 28
 <211> 100
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 238642CD1

<400> 28
 Met Gln Lys Pro Cys Lys Glu Asn Glu Gly Lys Pro Lys Cys Ser
 1 5 10 15
 Val Pro Lys Arg Glu Glu Lys Arg Pro Tyr Gly Glu Phe Glu Arg
 20 25 30
 Gln Gln Thr Glu Gly Asn Phe Arg Gln Arg Leu Leu Gln Ser Leu
 35 40 45
 Glu Glu Phe Lys Lys Asp Ile Asp Tyr Arg His Phe Lys Asp Glu
 50 55 60
 Glu Met Thr Arg Glu Gly Asp Glu Met Glu Arg Cys Leu Glu Glu
 65 70 75
 Ile Arg Gly Leu Arg Lys Lys Phe Arg Ala Leu His Ser Asn His
 80 85 90
 Arg His Ser Arg Asp Arg Pro Tyr Pro Ile
 95 100

<210> 29
 <211> 528
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 669862CD1

<400> 29
 Met Ser Ser Pro Tyr Pro Leu Leu Leu Glu Asn Ser Ile Cys Leu
 1 5 10 15
 Phe Phe His Phe Leu Pro Asp Phe Asn Phe Thr Thr Glu Ser Asn
 20 25 30
 Lys Leu Ser Ser Glu Lys Arg Asn Tyr Glu Val Asn Ala Tyr His
 35 40 45
 Gln Glu Thr Trp Lys Arg Asn Lys Thr Phe Asn Leu Met Arg Phe
 50 55 60
 Ile Phe Arg Thr Asp Pro Gln Tyr Thr Ile Glu Phe Gly Arg Gln
 65 70 75
 Gln Arg Pro Lys Val Gly Cys Phe Ser Gln Met Ile Phe Lys Lys
 80 85 90
 His Lys Ser Leu Pro Leu His Lys Arg Asn Asn Thr Arg Glu Lys
 95 100 105
 Ser Tyr Glu Cys Lys Glu Tyr Lys Lys Gly Phe Arg Lys Tyr Leu
 110 115 120
 His Leu Thr Glu His Leu Arg Asp His Thr Gly Val Ile Pro Tyr
 125 130 135
 Glu Cys Asn Glu Cys Gly Lys Ala Phe Val Val Phe Gln His Phe

31/91

Ile Arg His Arg	Lys	Ile His Thr Asp	Leu Lys Pro Tyr Glu Cys
Asn Gly Cys Glu	Lys	Ala Phe Arg Phe	Tyr Ser Gln Leu Ile Gln
His Gln Ile Ile	His	Thr Gly Met Lys	Pro Tyr Glu Cys Lys Gln
Cys Gly Lys Ala	Phe	Arg Arg His Ser	His Leu Thr Glu His Gln
Lys Ile His Val	Gly	Leu Lys Pro Phe	Glu Cys Lys Glu Cys Gly
Glu Thr Phe Arg	Leu	Tyr Arg His Met	Cys Leu His Gln Lys Ile
His His Gly Val	Lys	Pro Tyr Lys Cys	Lys Glu Cys Gly Lys Ala
Phe Gly His Arg	Ser	Ser Leu Tyr Gln	His Lys Lys Ile His Ser
Gly Glu Lys Pro	Tyr	Lys Cys Glu Gln	Cys Glu Lys Ala Phe Val
Arg Ser Tyr Leu	Leu	Val Glu His Gln	Arg Ser His Thr Gly Glu
Lys Pro His Glu	Cys	Met Glu Cys Gly	Lys Ala Phe Ser Lys Gly
Ser Ser Leu Leu	Lys	His Lys Arg Ile	His Ser Ser Glu Lys Leu
Tyr Asp Cys Lys	Asp	Cys Gly Lys Ala	Phe Cys Arg Gly Ser Gln
Leu Thr Gln His	Gln	Arg Ile His Thr	Gly Glu Lys Pro His Glu
Cys Lys Glu Cys	Gly	Lys Thr Phe Lys	Leu His Ser Tyr Leu Ile
Gln His Gln Ile	Ile	His Thr Asp Leu	Lys Pro Tyr Glu Cys Lys
Gln Cys Gly Lys	Ala	Phe Ser Arg Val	Gly Asp Leu Lys Thr His
Gln Ser Ile His	Ala	Gly Glu Lys Pro	Tyr Glu Cys Lys Glu Cys
Gly Lys Thr Phe	Arg	Leu Asn Ser Gln	Leu Ile Tyr His Gln Thr
Ile His Thr Gly	Leu	Lys Pro Tyr Val	Cys Lys Glu Cys Lys Lys
Ala Phe Arg Ser	Ile	Ser Gly Leu Ser	Gln His Lys Arg Ile His
Thr Gly Glu Lys	Pro	Tyr Glu Cys Lys	Glu Cys Asp Lys Ala Phe
Asn Arg Ser Asp	Arg	Leu Thr Gln His	Glu Thr Ile His Thr Gly
Val Lys Pro Gln	Lys	Cys Lys Glu Cys	Gly Lys Ala Phe Ser His
Cys Tyr Gln Leu	Ser	Gln His Gln Arg	Phe His His Gly Glu Arg
Leu Leu Met			

<210> 30
 <211> 350
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1003663CD1

```

<400> 30
Met His Pro Ala Ala Phe Pro Leu Pro Val Val Val Ala Ala Val
  1      5      10      15
Leu Trp Gly Ala Ala Pro Thr Arg Gly Leu Ile Arg Ala Thr Ser
  20      25      30
Asp His Asn Ala Ser Met Asp Phe Ala Asp Leu Pro Ala Leu Phe
  35      40      45
Gly Ala Thr Leu Ser Gln Glu Gly Leu Gln Gly Phe Leu Val Glu
  50      55      60
Ala His Pro Asp Asn Ala Cys Ser Pro Ile Ala Pro Pro Pro Pro
  65      70      75
Ala Pro Val Asn Gly Ser Val Phe Ile Ala Leu Leu Arg Arg Phe
  80      85      90
Asp Cys Asn Phe Asp Leu Lys Val Leu Asn Ala Gln Lys Ala Gly
  95      100     105
Tyr Gly Ala Ala Val Val His Asn Val Asn Ser Asn Glu Leu Leu
  110     115     120
Asn Met Val Trp Asn Ser Glu Glu Ile Gln Gln Gln Ile Trp Ile
  125     130     135
Pro Ser Val Phe Ile Gly Glu Arg Ser Ser Glu Tyr Leu Arg Ala
  140     145     150
Leu Phe Val Tyr Glu Lys Gly Ala Arg Val Leu Leu Val Pro Asp
  155     160     165
Asn Thr Phe Pro Leu Gly Tyr Tyr Leu Ile Pro Phe Thr Gly Ile
  170     175     180
Val Gly Leu Leu Val Leu Ala Met Gly Ala Val Met Ile Ala Arg
  185     190     195
Cys Ile Gln His Arg Lys Arg Leu Gln Arg Asn Arg Leu Thr Lys
  200     205     210
Glu Gln Leu Lys Gln Ile Pro Thr His Asp Tyr Gln Lys Gly Asp
  215     220     225
Gln Tyr Asp Val Cys Ala Ile Cys Leu Asp Glu Tyr Glu Asp Gly
  230     235     240
Asp Lys Leu Arg Val Leu Pro Cys Ala His Ala Tyr His Ser Arg
  245     250     255
Cys Val Asp Pro Trp Leu Thr Gln Thr Arg Lys Thr Cys Pro Ile
  260     265     270
Cys Lys Gln Pro Val His Arg Gly Pro Gly Asp Glu Asp Gln Glu
  275     280     285
Glu Glu Thr Gln Gly Gln Glu Glu Gly Asp Glu Gly Glu Pro Arg
  290     295     300
Asp His Pro Ala Ser Glu Arg Thr Pro Leu Leu Gly Ser Ser Pro
  305     310     315
Thr Leu Pro Thr Ser Phe Gly Ser Leu Ala Pro Ala Pro Leu Val
  320     325     330
Phe Pro Gly Pro Ser Thr Asp Pro Pro Leu Ser Pro Pro Ser Ser
  335     340     345
Pro Val Ile Leu Val
  350

```

```

<210> 31
<211> 315
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc-feature
<223> Incyte ID No.: 1432557CD1

```

<400> 31

```

Met Ala Ala Leu Gly Val Leu Glu Ser Asp Leu Pro Ser Ala Val
 1          5          10          15
Thr Leu Leu Lys Asn Leu Gln Glu Gln Val Met Ala Val Thr Ala
          20          25          30
Gln Val Lys Ser Leu Thr Gln Lys Val Gln Ala Gly Ala Tyr Pro
          35          40          45
Thr Glu Lys Gly Leu Ser Phe Leu Glu Val Lys Asp Gln Leu Leu
          50          55          60
Leu Met Tyr Leu Met Asp Leu Thr His Leu Ile Leu Asp Lys Ala
          65          70          75
Ser Gly Gly Ser Leu Gln Gly His Asp Ala Val Leu Arg Leu Val
          80          85          90
Glu Ile Arg Thr Val Leu Glu Lys Leu Arg Pro Leu Asp Gln Lys
          95          100          105
Leu Lys Tyr Gln Ile Asp Lys Leu Ile Lys Thr Ala Val Thr Gly
          110          115          120
Ser Leu Ser Glu Asn Asp Pro Leu Arg Phe Lys Pro His Pro Ser
          125          130          135
Asn Met Met Ser Lys Leu Ser Ser Glu Asp Glu Glu Glu Asp Glu
          140          145          150
Ala Glu Asp Asp Gln Ser Glu Ala Ser Gly Lys Lys Ser Val Lys
          155          160          165
Gly Val Ser Lys Lys Tyr Val Pro Pro Arg Leu Val Pro Val His
          170          175          180
Tyr Asp Glu Thr Glu Ala Glu Arg Glu Lys Lys Arg Leu Glu Arg
          185          190          195
Ala Lys Arg Arg Ala Leu Ser Ser Ser Val Ile Arg Glu Leu Lys
          200          205          210
Glu Gln Tyr Ser Asp Ala Pro Glu Glu Ile Arg Asp Ala Arg His
          215          220          225
Pro His Val Thr Arg Gln Ser Gln Glu Asp Gln His Arg Ile Asn
          230          235          240
Tyr Glu Glu Ser Met Met Val Arg Leu Ser Val Ser Lys Arg Glu
          245          250          255
Lys Gly Arg Arg Lys Arg Ala Asn Val Met Ser Ser Gln Leu His
          260          265          270
Ser Leu Thr His Phe Ser Asp Ile Ser Ala Leu Thr Gly Gly Thr
          275          280          285
Val His Leu Asp Glu Asp Gln Asn Pro Ile Lys Lys Arg Lys Lys
          290          295          300
Ile Pro Gln Lys Gly Arg Lys Lys Lys Gly Phe Arg Arg Arg Arg
          305          310          315

```

<210> 32

<211> 120

<212> PRT

<213> Homo sapiens

<220>

<221> misc-feature

<223> 1441770CD1

<400> 32

```

Met Asp Asp Ser Lys Val Val Gly Gly Lys Val Lys Lys Pro Gly
 1          5          10          15
Lys Arg Gly Arg Lys Pro Ala Lys Ile Asp Leu Lys Ala Lys Leu
          20          25          30
Glu Arg Ser Arg Gln Ser Ala Arg Glu Cys Arg Ala Arg Lys Lys
          35          40          45

```

Leu	Arg	Tyr	Gln	Tyr	Leu	Glu	Glu	Leu	Val	Ser	Ser	Arg	Glu	Arg
			50						55					60
Ala	Ile	Cys	Ala	Leu	Arg	Glu	Glu	Leu	Glu	Met	Tyr	Lys	Gln	Trp
			65						70					75
Cys	Met	Ala	Met	Asp	Gln	Gly	Lys	Ile	Pro	Ser	Glu	Ile	Lys	Ala
			80						85					90
Leu	Leu	Thr	Gly	Glu	Glu	Gln	Asn	Lys	Ser	Gln	Gln	Asn	Ser	Ser
			95						100					105
Arg	His	Thr	Lys	Ala	Gly	Lys	Thr	Asp	Ala	Asn	Ser	Asn	Ser	Trp
			110						115					120

<210> 33
 <211> 326
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1456684CD1

<400> 33

Met	Gln	Glu	Glu	Pro	Leu	Pro	Gln	Gly	Asn	Asp	Pro	Glu	Pro	Ser
1				5					10					15
Gly	Asp	Ser	Pro	Leu	Gly	Leu	Cys	Gln	Ser	Glu	Cys	Met	Glu	Met
			20						25					30
Ser	Glu	Val	Phe	Asp	Asp	Ala	Ser	Asp	Gln	Asp	Ser	Thr	Asp	Lys
			35						40					45
Pro	Trp	Arg	Pro	Tyr	Tyr	Asn	Tyr	Lys	Pro	Lys	Lys	Lys	Ser	Arg
			50						55					60
Gln	Leu	Lys	Lys	Met	Arg	Lys	Val	Asn	Trp	Arg	Lys	Glu	His	Gly
			65						70					75
Asn	Arg	Ser	Pro	Ser	His	Lys	Cys	Lys	Tyr	Pro	Ala	Glu	Leu	Asp
			80						85					90
Cys	Ala	Val	Gly	Lys	Ala	Pro	Gln	Asp	Lys	Pro	Phe	Glu	Glu	Glu
			95						100					105
Glu	Thr	Lys	Glu	Met	Pro	Lys	Leu	Gln	Cys	Glu	Leu	Cys	Asp	Gly
			110						115					120
Asp	Lys	Ala	Val	Gly	Ala	Gly	Asn	Gln	Gly	Arg	Pro	His	Arg	His
			125						130					135
Leu	Thr	Ser	Arg	Pro	Tyr	Ala	Cys	Glu	Leu	Cys	Ala	Lys	Gln	Phe
			140						145					150
Gln	Ser	Pro	Ser	Thr	Leu	Lys	Met	His	Met	Arg	Cys	His	Thr	Gly
			155						160					165
Glu	Lys	Pro	Tyr	Gln	Cys	Lys	Thr	Cys	Gly	Arg	Cys	Phe	Ser	Val
			170						175					180
Gln	Gly	Asn	Leu	Gln	Lys	His	Glu	Arg	Ile	His	Leu	Gly	Leu	Lys
			185						190					195
Glu	Phe	Val	Cys	Gln	Tyr	Cys	Asn	Lys	Ala	Phe	Thr	Leu	Asn	Glu
			200						205					210
Thr	Leu	Lys	Ile	His	Glu	Arg	Ile	His	Thr	Gly	Glu	Lys	Arg	Tyr
			215						220					225
His	Cys	Gln	Phe	Cys	Phe	Gln	Arg	Phe	Leu	Tyr	Leu	Ser	Thr	Lys
			230						235					240
Arg	Asn	His	Glu	Gln	Arg	His	Ile	Arg	Glu	His	Asn	Gly	Lys	Gly
			245						250					255
Tyr	Ala	Cys	Phe	Gln	Cys	Pro	Lys	Ile	Cys	Lys	Thr	Ala	Ala	Ala
			260						265					270
Leu	Gly	Met	His	Gln	Lys	Lys	His	Leu	Phe	Lys	Ser	Pro	Ser	Gln
			275						280					285

Gln Glu Lys Ile Gly Asp Val Cys His Glu Asn Ser Asn Pro Leu
 290 295 300
 Glu Asn Gln His Phe Ile Gly Ser Glu Asp Asn Asp Gln Lys Asp
 305 310 315
 Asn Ile Gln Thr Gly Val Glu Asn Val Val Leu
 320 325

<210> 34
 <211> 106
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1602916CD1

<400> 34
 Met Phe Pro Trp Met Arg Pro Gln Ala Ala Pro Gly Arg Arg Arg
 1 5 10 15
 Gly Arg Gln Thr Tyr Ser Arg Phe Gln Thr Leu Glu Leu Glu Lys
 20 25 30
 Glu Phe Leu Phe Asn Pro Tyr Leu Thr Arg Lys Arg Arg Ile Glu
 35 40 45
 Val Ser His Ala Leu Ala Leu Thr Glu Arg Gln Val Lys Ile Trp
 50 55 60
 Phe Gln Asn Arg Arg Met Lys Trp Lys Lys Glu Asn Asn Lys Asp
 65 70 75
 Lys Phe Pro Val Ser Arg Gln Glu Val Lys Asp Gly Glu Thr Lys
 80 85 90
 Lys Glu Ala Gln Glu Leu Glu Glu Asp Arg Ala Glu Arg Leu Thr
 95 100 105
 Asn

<210> 35
 <211> 209
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1692816CD1

<400> 35
 Met Asn Pro Ser Met Lys Gln Lys Gln Glu Glu Ile Lys Glu Asn
 1 5 10 15
 Ile Lys Asn Ser Ser Val Pro Arg Arg Thr Leu Lys Met Ile Gln
 20 25 30
 Pro Ser Ala Ser Gly Ser Leu Val Gly Arg Glu Asn Glu Leu Ser
 35 40 45
 Ala Gly Leu Ser Lys Arg Lys His Arg Asn Asp His Leu Thr Ser
 50 55 60
 Thr Thr Ser Ser Pro Gly Val Ile Val Pro Glu Ser Ser Glu Asn
 65 70 75
 Lys Asn Leu Gly Gly Val Thr Gln Glu Ser Phe Asp Leu Met Ile
 80 85 90
 Lys Glu Asn Pro Ser Ser Gln Tyr Trp Lys Glu Val Ala Glu Lys

	95		100		105
Arg Arg Lys Ala	Leu Tyr Glu Ala Leu	Lys Glu Asn Glu Lys	Leu		
	110		115		120
His Lys Glu Ile	Glu Gln Lys Asp Asn	Glu Ile Ala Arg Leu	Lys		
	125		130		135
Lys Glu Asn Lys	Glu Leu Ala Glu Val	Ala Glu His Val Gln	Tyr		
	140		145		150
Met Ala Glu Leu	Ile Glu Arg Leu Asn	Gly Glu Pro Leu Asp	Asn		
	155		160		165
Phe Glu Ser Leu	Asp Asn Gln Glu Phe	Asp Ser Glu Glu Glu	Thr		
	170		175		180
Val Glu Asp Ser	Leu Val Glu Asp Ser	Glu Ile Gly Thr Cys	Ala		
	185		190		195
Glu Gly Thr Val	Ser Ser Thr Asp	Ala Lys Pro Cys Ile			
	200		205		

<210> 36
 <211> 212
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1968191CD1

<400> 36

Met Leu Gly Asn Glu Trp Ser Lys Leu Pro	Pro Glu Glu Lys Gln
1	5 10 15
Arg Tyr Leu Asp Glu Ala Asp Arg Asp	Lys Glu Arg Tyr Met Lys
	20 25 30
Glu Leu Glu Gln Tyr Gln Lys Thr Glu Ala Tyr	Lys Val Phe Ser
	35 40 45
Arg Lys Thr Gln Asp Arg Gln Lys Gly Lys	Ser His Arg Gln Asp
	50 55 60
Ala Ala Arg Gln Ala Thr His Asp His Glu	Lys Glu Thr Glu Val
	65 70 75
Lys Glu Arg Ser Val Phe Asp Ile Pro Ile	Phe Thr Glu Glu Phe
	80 85 90
Leu Asn His Ser Lys Ala Arg Glu Ala Glu	Leu Arg Gln Leu Arg
	95 100 105
Lys Ser Asn Met Glu Phe Glu Glu Arg Asn	Ala Ala Leu Gln Lys
	110 115 120
His Val Glu Ser Met Arg Thr Ala Val Glu	Lys Leu Glu Val Asp
	125 130 135
Val Ile Gln Glu Arg Ser Arg Asn Thr Val	Leu Gln Gln His Leu
	140 145 150
Glu Thr Leu Arg Gln Val Leu Thr Ser Ser	Phe Ala Ser Met Pro
	155 160 165
Leu Pro Gly Ser Gly Glu Thr Pro Thr Val	Asp Thr Ile Asp Ser
	170 175 180
Tyr Met Asn Arg Leu His Ser Ile Ile Leu	Ala Asn Pro Gln Asp
	185 190 195
Asn Glu Asn Phe Ile Ala Thr Val Arg Glu	Val Val Asn Arg Leu
	200 205 210

Asp Arg

<210> 37
 <211> 359
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2052061CD1

<400> 37

Met	Val	Asp	Met	Asp	Lys	Leu	Ile	Asn	Asn	Leu	Glu	Val	Gln	Leu	
1				5					10					15	
Asn	Ser	Glu	Gly	Gly	Ser	Met	Gln	Val	Phe	Lys	Gln	Val	Thr	Ala	
				20					25					30	
Ser	Val	Arg	Asn	Arg	Asp	Pro	Pro	Glu	Ile	Glu	Tyr	Thr	Ser	Asn	
				35					40					45	
Met	Thr	Ser	Pro	Thr	Leu	Leu	Asp	Ala	Asn	Pro	Met	Glu	Asn	Pro	
				50					55					60	
Ala	Leu	Phe	Asn	Asp	Ile	Lys	Ile	Glu	Pro	Pro	Glu	Glu	Leu	Leu	
				65					70					75	
Ala	Ser	Asp	Phe	Ser	Leu	Pro	Gln	Val	Glu	Pro	Val	Asp	Leu	Ser	
				80					85					90	
Phe	His	Lys	Pro	Lys	Ala	Pro	Leu	Gln	Pro	Ala	Ser	Met	Leu	Gln	
				95					100					105	
Ala	Pro	Ile	Arg	Pro	Pro	Lys	Pro	Gln	Ser	Ser	Pro	Gln	Thr	Leu	
				110					115					120	
Val	Val	Ser	Thr	Ser	Thr	Ser	Asp	Met	Ser	Thr	Ser	Ala	Asn	Ile	
				125					130					135	
Pro	Thr	Val	Leu	Thr	Pro	Gly	Ser	Val	Leu	Thr	Ser	Ser	Gln	Ser	
				140					145					150	
Thr	Gly	Ser	Gln	Gln	Ile	Leu	His	Val	Ile	His	Thr	Ile	Pro	Ser	
				155					160					165	
Val	Ser	Leu	Pro	Asn	Lys	Met	Gly	Gly	Leu	Lys	Thr	Ile	Pro	Val	
				170					175					180	
Val	Val	Gln	Ser	Leu	Pro	Met	Val	Tyr	Thr	Thr	Leu	Pro	Ala	Asp	
				185					190					195	
Gly	Gly	Pro	Ala	Ala	Ile	Thr	Val	Pro	Leu	Ile	Gly	Gly	Asp	Gly	
				200					205					210	
Lys	Asn	Ala	Gly	Ser	Val	Lys	Val	Asp	Pro	Thr	Ser	Met	Ser	Pro	
				215					220					225	
Leu	Glu	Ile	Pro	Ser	Asp	Ser	Glu	Glu	Ser	Thr	Ile	Glu	Ser	Gly	
				230					235					240	
Ser	Ser	Ala	Leu	Gln	Ser	Leu	Gln	Gly	Leu	Gln	Gln	Glu	Pro	Ala	
				245					250					255	
Ala	Met	Ala	Gln	Met	Gln	Gly	Glu	Glu	Ser	Leu	Asp	Leu	Lys	Arg	
				260					265					270	
Arg	Arg	Ile	His	Gln	Cys	Asp	Phe	Ala	Gly	Cys	Ser	Lys	Val	Tyr	
				275					280					285	
Thr	Lys	Ser	Ser	His	Leu	Lys	Ala	His	Arg	Arg	Ile	His	Thr	Gly	
				290					295					300	
Glu	Lys	Pro	Tyr	Lys	Cys	Thr	Trp	Asp	Gly	Cys	Ser	Trp	Lys	Phe	
				305					310					315	
Ala	Arg	Ser	Asp	Glu	Leu	Thr	Arg	His	Phe	Arg	Lys	His	Thr	Gly	
				320					325					330	
Ile	Lys	Pro	Phe	Arg	Cys	Thr	Asp	Cys	Asn	Arg	Ser	Phe	Ser	Arg	
				335					340					345	
Ser	Asp	His	Leu	Ser	Leu	His	Arg	Arg	Arg	His	Asp	Thr	Met		
				350					355						

<210> 38
 <211> 445
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2056207CD1

<400> 38
 Met Val Glu Cys Ile Arg Glu Val Asn Glu Val Ile Gln Asn Pro
 1 5 10 15
 Ala Thr Ile Thr Arg Ile Leu Leu Ser His Phe Asn Trp Asp Lys
 20 25 30
 Glu Lys Leu Met Glu Arg Tyr Phe Asp Gly Asn Leu Glu Lys Leu
 35 40 45
 Phe Ala Glu Cys His Val Ile Asn Pro Ser Lys Lys Ser Arg Thr
 50 55 60
 Arg Gln Met Asn Thr Arg Ser Ser Ala Gln Asp Met Pro Cys Gln
 65 70 75
 Ile Cys Tyr Leu Asn Tyr Pro Asn Ser Tyr Phe Thr Gly Leu Glu
 80 85 90
 Cys Gly His Lys Phe Cys Met Gln Cys Trp Ser Glu Tyr Leu Thr
 95 100 105
 Thr Lys Ile Met Glu Glu Gly Met Gly Gln Thr Ile Ser Cys Pro
 110 115 120
 Ala His Gly Cys Asp Ile Leu Val Asp Asp Asn Thr Val Met Arg
 125 130 135
 Leu Ile Thr Asp Ser Lys Val Lys Leu Lys Tyr Gln His Leu Ile
 140 145 150
 Thr Asn Ser Phe Val Glu Cys Asn Arg Leu Leu Lys Trp Cys Pro
 155 160 165
 Ala Pro Asp Cys His His Val Val Lys Val Gln Tyr Pro Asp Ala
 170 175 180
 Lys Pro Val Arg Cys Lys Cys Gly Arg Gln Phe Cys Phe Asn Cys
 185 190 195
 Gly Glu Asn Trp His Asp Pro Val Lys Cys Lys Trp Leu Lys Lys
 200 205 210
 Trp Ile Lys Lys Cys Asp Asp Asp Ser Glu Thr Ser Asn Trp Ile
 215 220 225
 Ala Ala Asn Thr Lys Glu Cys Pro Lys Cys His Val Thr Ile Glu
 230 235 240
 Lys Asp Gly Gly Cys Asn His Met Val Cys Arg Asn Gln Asn Cys
 245 250 255
 Lys Ala Glu Phe Cys Trp Val Cys Leu Gly Pro Trp Glu Pro His
 260 265 270
 Gly Ser Ala Trp Tyr Asn Cys Asn Arg Tyr Asn Glu Asp Asp Ala
 275 280 285
 Lys Ala Ala Arg Asp Ala Gln Glu Arg Ser Arg Ala Ala Leu Gln
 290 295 300
 Arg Tyr Leu Phe Tyr Cys Asn Arg Tyr Met Asn His Met Gln Ser
 305 310 315
 Leu Arg Phe Glu His Lys Leu Tyr Ala Gln Val Lys Gln Lys Met
 320 325 330
 Glu Glu Met Gln Gln His Asn Met Ser Trp Ile Glu Val Gln Phe
 335 340 345
 Leu Lys Lys Ala Val Asp Val Leu Cys Gln Cys Arg Ala Thr Leu
 350 355 360
 Met Tyr Thr Tyr Val Phe Ala Phe Tyr Leu Lys Lys Asn Asn Gln
 365 370 375
 Ser Ile Ile Phe Glu Asn Asn Gln Ala Asp Leu Glu Asn Ala Thr
 380 385 390

```

Glu Val Leu Ser Gly Tyr Leu Glu Arg Asp Ile Ser Gln Asp Ser
      395              400              405
Leu Gln Asp Ile Lys Gln Lys Val Gln Asp Lys Tyr Arg Tyr Cys
      410              415              420
Glu Ser Arg Arg Arg Val Leu Leu Gln His Val His Glu Gly Tyr
      425              430              435
Glu Lys Asp Leu Trp Glu Tyr Ile Glu Asp
      440              445

```

```

<210> 39
<211> 433
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc-feature
<223> Incyte ID No.: 2101803CD1

```

```

<400> 39
Met Arg Ala Glu Gly Leu Gly Gly Leu Glu Arg Phe Cys Ser Pro
  1          5          10          15
Gly Lys Gly Arg Gly Leu Arg Ala Leu Gln Pro Phe Gln Val Gly
      20          25          30
Asp Leu Leu Phe Ser Cys Pro Ala Tyr Ala Tyr Val Leu Thr Val
      35          40          45
Asn Glu Arg Gly Asn His Cys Glu Tyr Cys Phe Thr Arg Lys Glu
      50          55          60
Gly Leu Ser Lys Cys Gly Arg Cys Lys Gln Ala Phe Tyr Cys Asn
      65          70          75
Val Glu Cys Gln Lys Glu Asp Trp Pro Met His Lys Leu Glu Cys
      80          85          90
Ser Pro Met Val Val Phe Gly Glu Asn Trp Asn Pro Ser Glu Thr
      95          100         105
Val Arg Leu Thr Ala Arg Ile Leu Ala Lys Gln Lys Ile His Pro
     110         115         120
Glu Arg Thr Pro Ser Glu Lys Leu Leu Ala Val Lys Glu Phe Glu
     125         130         135
Ser His Leu Asp Lys Leu Asp Asn Glu Lys Lys Asp Leu Ile Gln
     140         145         150
Ser Asp Ile Ala Ala Leu His His Phe Tyr Ser Lys His Leu Glu
     155         160         165
Phe Pro Asp Asn Asp Ser Leu Val Val Leu Phe Ala Gln Val Asn
     170         175         180
Cys Asn Gly Phe Thr Ile Glu Asp Glu Glu Leu Ser His Leu Gly
     185         190         195
Ser Ala Ile Phe Pro Asp Val Ala Leu Met Asn His Ser Cys Cys
     200         205         210
Pro Asn Val Ile Val Thr Tyr Lys Gly Thr Leu Ala Glu Val Arg
     215         220         225
Ala Val Gln Glu Ile Lys Pro Gly Glu Glu Val Phe Thr Ser Tyr
     230         235         240
Ile Asp Leu Leu Tyr Pro Thr Glu Asp Arg Asn Asp Arg Leu Arg
     245         250         255
Asp Ser Tyr Phe Phe Thr Cys Glu Cys Gln Glu Cys Thr Thr Lys
     260         265         270
Asp Lys Asp Lys Ala Lys Val Glu Ile Arg Lys Leu Ser Asp Pro
     275         280         285
Pro Lys Ala Glu Ala Ile Arg Asp Met Val Arg Tyr Ala Arg Asn
     290         295         300

```

Val	Ile	Glu	Glu	Phe	Arg	Arg	Ala	Lys	His	Tyr	Lys	Ser	Pro	Ser	
				305					310					315	
Glu	Leu	Leu	Glu	Ile	Cys	Glu	Leu	Ser	Gln	Glu	Lys	Met	Ser	Ser	
				320					325					330	
Val	Phe	Glu	Asp	Ser	Asn	Val	Tyr	Met	Leu	His	Met	Met	Tyr	Gln	
				335					340					345	
Ala	Met	Gly	Val	Cys	Leu	Tyr	Met	Gln	Asp	Trp	Glu	Gly	Ala	Leu	
				350					355					360	
Gln	Tyr	Gly	Gln	Lys	Ile	Ile	Lys	Pro	Tyr	Ser	Lys	His	Tyr	Pro	
				365					370					375	
Leu	Tyr	Ser	Leu	Asn	Val	Ala	Ser	Met	Trp	Leu	Lys	Leu	Gly	Arg	
				380					385					390	
Leu	Tyr	Met	Gly	Leu	Glu	His	Lys	Ala	Ala	Gly	Glu	Lys	Ala	Leu	
				395					400					405	
Lys	Lys	Ala	Ile	Ala	Ile	Met	Glu	Val	Ala	His	Gly	Lys	Asp	His	
				410					415					420	
Pro	Tyr	Ile	Ser	Glu	Ile	Lys	Gln	Glu	Ile	Glu	Ser	His			
				425					430						

<210> 40
 <211> 355
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2112362CD1

<400> 40

Met	Ser	Val	Asn	Tyr	Ala	Ala	Gly	Leu	Ser	Pro	Tyr	Ala	Asp	Lys	
1			5						10					15	
Gly	Lys	Cys	Gly	Leu	Pro	Glu	Ile	Phe	Asp	Pro	Pro	Glu	Glu	Leu	
			20						25					30	
Glu	Arg	Lys	Val	Trp	Glu	Leu	Ala	Arg	Leu	Val	Trp	Gln	Ser	Ser	
			35						40					45	
Asn	Val	Val	Phe	His	Thr	Gly	Ala	Gly	Ile	Ser	Thr	Ala	Ser	Gly	
			50						55					60	
Ile	Pro	Asp	Phe	Arg	Gly	Pro	His	Gly	Val	Trp	Thr	Met	Glu	Glu	
			65						70					75	
Arg	Gly	Leu	Ala	Pro	Lys	Phe	Asp	Thr	Thr	Phe	Glu	Ser	Ala	Arg	
			80						85					90	
Pro	Thr	Gln	Thr	His	Met	Ala	Leu	Val	Gln	Leu	Glu	Arg	Val	Gly	
			95						100					105	
Leu	Leu	Arg	Phe	Leu	Val	Ser	Gln	Asn	Val	Asp	Gly	Leu	His	Val	
			110						115					120	
Arg	Ser	Gly	Phe	Pro	Arg	Asp	Lys	Leu	Ala	Glu	Leu	His	Gly	Asn	
			125						130					135	
Met	Phe	Val	Glu	Glu	Cys	Ala	Lys	Cys	Lys	Thr	Gln	Tyr	Val	Arg	
			140						145					150	
Asp	Thr	Val	Val	Gly	Thr	Met	Gly	Leu	Lys	Ala	Thr	Gly	Arg	Leu	
			155						160					165	
Cys	Thr	Val	Ala	Lys	Ala	Arg	Gly	Leu	Arg	Ala	Cys	Arg	Gly	Glu	
			170						175					180	
Leu	Arg	Asp	Thr	Ile	Leu	Asp	Trp	Glu	Asp	Ser	Leu	Pro	Asp	Arg	
			185						190					195	
Asp	Leu	Ala	Leu	Ala	Asp	Glu	Ala	Ser	Arg	Asn	Ala	Asp	Leu	Ser	
			200						205					210	
Ile	Thr	Leu	Gly	Thr	Ser	Leu	Gln	Ile	Arg	Pro	Ser	Gly	Asn	Leu	
			215						220					225	

Pro	Leu	Ala	Thr	Lys	Arg	Arg	Gly	Gly	Arg	Leu	Val	Ile	Val	Asn	
				230					235					240	
Leu	Gln	Pro	Thr	Lys	His	Asp	Arg	His	Ala	Asp	Leu	Arg	Ile	His	
				245					250					255	
Gly	Tyr	Val	Asp	Glu	Val	Met	Thr	Arg	Leu	Met	Lys	His	Leu	Gly	
				260					265					270	
Leu	Glu	Ile	Pro	Ala	Trp	Asp	Gly	Pro	Arg	Val	Leu	Glu	Arg	Ala	
				275					280					285	
Leu	Pro	Pro	Leu	Pro	Arg	Pro	Pro	Thr	Pro	Lys	Leu	Glu	Pro	Lys	
				290					295					300	
Glu	Glu	Ser	Pro	Thr	Arg	Ile	Asn	Gly	Ser	Ile	Pro	Ala	Gly	Pro	
				305					310					315	
Lys	Gln	Glu	Pro	Cys	Ala	Gln	His	Asn	Gly	Ser	Glu	Pro	Ala	Ser	
				320					325					330	
Pro	Lys	Arg	Glu	Arg	Pro	Thr	Ser	Pro	Ala	Pro	His	Arg	Pro	Pro	
				335					340					345	
Lys	Arg	Val	Lys	Ala	Lys	Ala	Val	Pro	Ser						
				350					355						

<210> 41
 <211> 443
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2117346CD1

<400> 41

Met	Asp	Arg	Leu	Gly	Ser	Phe	Ser	Asn	Asp	Pro	Ser	Asp	Lys	Pro	
1				5					10					15	
Pro	Cys	Arg	Gly	Cys	Ser	Ser	Tyr	Leu	Met	Glu	Pro	Tyr	Ile	Lys	
				20					25					30	
Cys	Ala	Glu	Cys	Gly	Pro	Pro	Pro	Phe	Phe	Leu	Cys	Leu	Gln	Cys	
				35					40					45	
Phe	Thr	Arg	Gly	Phe	Glu	Tyr	Lys	Lys	His	Gln	Ser	Asp	His	Thr	
				50					55					60	
Tyr	Glu	Ile	Met	Thr	Ser	Asp	Phe	Pro	Val	Leu	Asp	Pro	Ser	Trp	
				65					70					75	
Thr	Ala	Gln	Glu	Glu	Met	Ala	Leu	Leu	Glu	Ala	Val	Met	Asp	Cys	
				80					85					90	
Gly	Phe	Gly	Asn	Trp	Gln	Asp	Val	Ala	Asn	Gln	Met	Cys	Thr	Lys	
				95					100					105	
Thr	Lys	Glu	Glu	Cys	Glu	Lys	His	Tyr	Met	Lys	His	Phe	Ile	Asn	
				110					115					120	
Asn	Pro	Leu	Phe	Ala	Ser	Thr	Leu	Leu	Asn	Leu	Lys	Gln	Ala	Glu	
				125					130					135	
Glu	Ala	Lys	Thr	Ala	Asp	Thr	Ala	Ile	Pro	Phe	His	Ser	Thr	Asp	
				140					145					150	
Asp	Pro	Pro	Arg	Pro	Thr	Phe	Asp	Ser	Leu	Leu	Ser	Arg	Asp	Met	
				155					160					165	
Ala	Gly	Tyr	Met	Pro	Ala	Arg	Ala	Asp	Phe	Ile	Glu	Glu	Phe	Asp	
				170					175					180	
Asn	Tyr	Ala	Glu	Trp	Asp	Leu	Arg	Asp	Ile	Asp	Phe	Val	Glu	Asp	
				185					190					195	
Asp	Ser	Asp	Ile	Leu	His	Ala	Leu	Lys	Met	Ala	Val	Val	Asp	Ile	
				200					205					210	
Tyr	His	Ser	Arg	Leu	Lys	Glu	Arg	Gln	Arg	Arg	Lys	Lys	Ile	Ile	
				215					220					225	

Arg Asp His Gly	Leu Ile Asn Leu Arg	Lys Phe Gln Leu Met Glu	
230		235	240
Arg Arg Tyr Pro	Lys Glu Val Gln Asp	Leu Tyr Glu Thr Met Arg	
245		250	255
Arg Phe Ala Arg	Ile Val Gly Pro Val	Glu His Asp Lys Phe Ile	
260		265	270
Glu Ser His Ala	Leu Glu Phe Glu Leu	Arg Arg Glu Ile Lys Arg	
275		280	285
Leu Gln Glu Tyr	Arg Thr Ala Gly Ile	Thr Asn Phe Cys Ser Ala	
290		295	300
Arg Thr Tyr Asp	His Leu Lys Lys Thr	Arg Glu Glu Glu Arg Leu	
305		310	315
Lys Arg Thr Met	Leu Ser Glu Val Leu	Gln Tyr Ile Gln Asp Ser	
320		325	330
Ser Ala Cys Gln	Gln Trp Leu Arg Arg	Gln Ala Asp Ile Asp Ser	
335		340	345
Gly Leu Ser Pro	Ser Ile Pro Met Ala	Ser Asn Ser Gly Arg Arg	
350		355	360
Ser Ala Pro Pro	Leu Asn Leu Thr Gly	Leu Pro Gly Thr Glu Lys	
365		370	375
Leu Asn Glu Lys	Glu Lys Glu Leu Cys	Gln Met Val Arg Leu Val	
380		385	390
Pro Gly Ala Tyr	Leu Glu Tyr Lys Ser	Ala Leu Leu Asn Glu Cys	
395		400	405
Asn Lys Gln Gly	Gly Leu Arg Leu Ala	Gln Ala Arg Ala Leu Ile	
410		415	420
Lys Ile Asp Val	Asn Lys Thr Arg Lys	Ile Tyr Asp Phe Leu Ile	
425		430	435
Arg Glu Gly Tyr	Ile Thr Lys Gly		
440			

<210> 42
 <211> 164
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2119917CD1

<400> 42	
Met Thr Ala Ser Ala Gln Pro Arg Gly Arg Arg Pro Gly Val Gly	
1 5 10 15	
Val Gly Val Val Val Thr Ser Cys Lys His Pro Arg Cys Val Leu	
20 25 30	
Leu Gly Lys Arg Lys Gly Ser Val Gly Ala Gly Ser Phe Gln Leu	
35 40 45	
Pro Gly Gly His Leu Glu Phe Gly Glu Thr Trp Glu Glu Cys Ala	
50 55 60	
Gln Arg Glu Thr Trp Glu Glu Ala Ala Leu His Leu Lys Asn Val	
65 70 75	
His Phe Ala Ser Val Val Asn Ser Phe Ile Glu Lys Glu Asn Tyr	
80 85 90	
His Tyr Val Thr Ile Leu Met Lys Gly Glu Val Asp Val Thr His	
95 100 105	
Asp Ser Glu Pro Lys Asn Val Glu Pro Glu Lys Asn Glu Ser Trp	
110 115 120	
Glu Trp Val Pro Trp Glu Glu Leu Pro Pro Leu Asp Gln Leu Phe	
125 130 135	

Trp	Gly	Leu	Arg	Cys	Leu	Lys	Glu	Gln	Gly	Tyr	Asp	Pro	Phe	Lys
				140					145					150
Glu	Asp	Leu	Asn	His	Leu	Val	Gly	Tyr	Lys	Gly	Asn	His	Leu	
				155					160					

<210> 43
 <211> 215
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2123456CD1

<400> 43

Met	Leu	Gly	Ala	Glu	Trp	Ser	Lys	Leu	Gln	Pro	Thr	Glu	Lys	Gln
1				5					10					15
Arg	Tyr	Leu	Asp	Glu	Ala	Glu	Arg	Glu	Lys	Gln	Gln	Tyr	Met	Lys
				20					25					30
Glu	Leu	Arg	Ala	Tyr	Gln	Gln	Ser	Glu	Ala	Tyr	Lys	Met	Cys	Thr
				35					40					45
Glu	Lys	Ile	Gln	Glu	Lys	Lys	Ile	Lys	Lys	Glu	Asp	Ser	Ser	Ser
				50					55					60
Gly	Leu	Met	Asn	Thr	Leu	Leu	Asn	Gly	His	Lys	Gly	Gly	Asp	Cys
				65					70					75
Asp	Gly	Phe	Ser	Thr	Phe	Asp	Val	Pro	Ile	Phe	Thr	Glu	Glu	Phe
				80					85					90
Leu	Asp	Gln	Asn	Lys	Ala	Arg	Glu	Ala	Glu	Leu	Arg	Arg	Leu	Arg
				95					100					105
Lys	Met	Asn	Val	Ala	Phe	Glu	Glu	Gln	Asn	Ala	Val	Leu	Gln	Arg
				110					115					120
His	Thr	Gln	Ser	Met	Ser	Ser	Ala	Arg	Glu	Arg	Leu	Glu	Gln	Glu
				125					130					135
Leu	Ala	Leu	Glu	Glu	Arg	Arg	Thr	Leu	Ala	Leu	Gln	Gln	Gln	Leu
				140					145					150
Gln	Ala	Val	Arg	Gln	Ala	Leu	Thr	Ala	Ser	Phe	Ala	Ser	Leu	Pro
				155					160					165
Val	Pro	Gly	Thr	Gly	Glu	Thr	Pro	Thr	Leu	Gly	Thr	Leu	Asp	Phe
				170					175					180
Tyr	Met	Ala	Arg	Leu	His	Gly	Ala	Ile	Glu	Arg	Asp	Pro	Ala	Gln
				185					190					195
His	Glu	Lys	Leu	Ile	Val	Arg	Ile	Lys	Glu	Ile	Leu	Ala	Gln	Val
				200					205					210
Ala	Ser	Glu	His	Leu										
				215										

<210> 44
 <211> 539
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2148792CD1

<400> 44

Met	Ala	Ala	Leu	Phe	Leu	Ser	Ala	Pro	Pro	Gln	Ala	Glu	Val	Thr
1				5					10					15
Phe	Glu	Asp	Val	Ala	Val	Tyr	Leu	Ser	Arg	Glu	Glu	Trp	Gly	Arg
			20						25					30
Leu	Gly	Pro	Ala	Gln	Arg	Gly	Leu	Tyr	Arg	Asp	Val	Met	Leu	Glu
			35						40					45
Thr	Tyr	Gly	Asn	Leu	Val	Ser	Leu	Gly	Val	Gly	Pro	Ala	Gly	Pro
			50						55					60
Lys	Pro	Gly	Val	Ile	Ser	Gln	Leu	Glu	Arg	Gly	Asp	Glu	Pro	Trp
			65						70					75
Val	Leu	Asp	Val	Gln	Gly	Thr	Ser	Gly	Lys	Glu	His	Leu	Arg	Val
			80						85					90
Asn	Ser	Pro	Ala	Leu	Gly	Thr	Arg	Thr	Glu	Tyr	Lys	Glu	Leu	Thr
			95						100					105
Ser	Gln	Glu	Thr	Phe	Gly	Glu	Glu	Asp	Pro	Gln	Gly	Ser	Glu	Pro
			110						115					120
Val	Glu	Ala	Cys	Asp	His	Ile	Ser	Lys	Ser	Glu	Gly	Ser	Leu	Glu
			125						130					135
Lys	Leu	Val	Glu	Gln	Arg	Gly	Pro	Arg	Ala	Val	Thr	Leu	Thr	Asn
			140						145					150
Gly	Glu	Ser	Ser	Arg	Glu	Ser	Gly	Gly	Asn	Leu	Arg	Leu	Leu	Ser
			155						160					165
Arg	Pro	Val	Pro	Asp	Gln	Arg	Pro	His	Lys	Cys	Asp	Ile	Cys	Glu
			170						175					180
Gln	Ser	Phe	Glu	Gln	Arg	Ser	Tyr	Leu	Asn	Asn	His	Lys	Arg	Val
			185						190					195
His	Arg	Ser	Lys	Lys	Thr	Asn	Thr	Val	Arg	Asn	Ser	Gly	Glu	Ile
			200						205					210
Phe	Ser	Ala	Asn	Leu	Val	Val	Lys	Glu	Asp	Gln	Lys	Ile	Pro	Thr
			215						220					225
Gly	Lys	Lys	Leu	His	Tyr	Cys	Ser	Tyr	Cys	Gly	Lys	Thr	Phe	Arg
			230						235					240
Tyr	Ser	Ala	Asn	Leu	Val	Lys	His	Gln	Arg	Leu	His	Thr	Glu	Glu
			245						250					255
Lys	Pro	Tyr	Lys	Cys	Asp	Glu	Cys	Gly	Lys	Ala	Phe	Ser	Gln	Ser
			260						265					270
Cys	Glu	Phe	Ile	Asn	His	Arg	Arg	Met	His	Ser	Gly	Glu	Ile	Pro
			275						280					285
Tyr	Arg	Cys	Asp	Glu	Cys	Gly	Lys	Thr	Phe	Thr	Arg	Arg	Pro	Asn
			290						295					300
Leu	Met	Lys	His	Gln	Arg	Ile	His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys
			305						310					315
Cys	Gly	Glu	Cys	Gly	Lys	His	Phe	Ser	Ala	Tyr	Ser	Ser	Leu	Ile
			320						325					330
Tyr	His	Gln	Arg	Ile	His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys	Cys	Asn
			335						340					345
Asp	Cys	Gly	Lys	Ala	Phe	Ser	Asp	Gly	Ser	Ile	Leu	Ile	Arg	His
			350						355					360
Arg	Arg	Thr	His	Thr	Gly	Glu	Lys	Pro	Phe	Glu	Cys	Lys	Glu	Cys
			365						370					375
Gly	Lys	Gly	Phe	Thr	Gln	Ser	Ser	Asn	Leu	Ile	Gln	His	Gln	Arg
			380						385					390
Ile	His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys	Cys	Asn	Glu	Cys	Glu	Lys
			395						400					405
Ala	Phe	Ile	Gln	Lys	Thr	Lys	Leu	Val	Glu	His	Gln	Arg	Ser	His
			410						415					420
Thr	Gly	Glu	Lys	Pro	Tyr	Glu	Cys	Asn	Asp	Cys	Gly	Lys	Val	Phe
			425						430					435
Ser	Gln	Ser	Thr	His	Leu	Ile	Gln	His	Gln	Arg	Ile	His	Thr	Gly
			440						445					450
Glu	Lys	Pro	Tyr	Lys	Cys	Ser	Glu	Cys	Gly	Lys	Ala	Phe	His	Asn
			455						460					465

```

Ser Ser Arg Leu Ile His His Gln Arg Leu His His Gly Glu Lys
470 475 480
Pro Tyr Arg Cys Ser Asp Cys Lys Lys Ala Phe Ser Gln Ser Thr
485 490 495
Tyr Leu Ile Gln His Arg Arg Ile His Thr Gly Glu Lys Pro Tyr
500 505 510
Lys Cys Ser Glu Cys Gly Lys Ala Phe Arg His Ser Ser Asn Met
515 520 525
Cys Gln His Gln Arg Ile His Leu Arg Glu Asp Phe Ser Met
530 535

```

```

<210> 45
<211> 182
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc-feature
<223> Incyte ID No.: 2751943CD1

```

```

<400> 45
Met Ala Arg Leu Leu Trp Leu Leu Arg Gly Leu Thr Leu Gly Thr
1 5 10 15
Ala Pro Arg Arg Ala Val Arg Gly Gln Ala Gly Gly Gly Gly Pro
20 25 30
Gly Thr Gly Pro Gly Leu Gly Glu Ala Gly Ser Leu Ala Thr Cys
35 40 45
Glu Leu Pro Leu Ala Lys Ser Glu Trp Gln Lys Lys Leu Thr Pro
50 55 60
Glu Gln Phe Tyr Val Thr Arg Glu Lys Gly Thr Glu Pro Pro Phe
65 70 75
Ser Gly Ile Tyr Leu Asn Asn Lys Glu Ala Gly Met Tyr His Cys
80 85 90
Val Cys Cys Asp Ser Pro Leu Phe Ser Ser Glu Lys Lys Tyr Cys
95 100 105
Ser Gly Thr Gly Trp Pro Ser Phe Ser Glu Ala His Gly Thr Ser
110 115 120
Gly Ser Asp Glu Ser His Thr Gly Ile Leu Arg Arg Leu Asp Thr
125 130 135
Ser Leu Gly Ser Ala Arg Thr Glu Val Val Cys Lys Gln Cys Glu
140 145 150
Ala His Leu Gly His Val Phe Pro Asp Gly Pro Gly Pro Asn Gly
155 160 165

Gln Arg Phe Cys Ile Asn Ser Val Ala Leu Lys Phe Lys Pro Arg
170 175 180
Lys His

```

```

<210> 46
<211> 534
<212> PRT
<213> Homo sapiens

```

```

<220>
<221> misc-feature
<223> Incyte ID No.: 3128913CD1

```

<400> 46

Met	Ala	Val	Glu	Ser	Gly	Val	Ile	Ser	Thr	Leu	Ile	Pro	Gln	Asp
1				5					10					15
Pro	Pro	Glu	Gln	Glu	Leu	Ile	Leu	Val	Lys	Val	Glu	Asp	Asn	Phe
				20					25					30
Ser	Trp	Asp	Glu	Lys	Phe	Lys	Gln	Asn	Gly	Ser	Thr	Gln	Ser	Cys
				35					40					45
Gln	Glu	Leu	Phe	Arg	Gln	Gln	Phe	Arg	Lys	Phe	Cys	Tyr	Gln	Glu
				50					55					60
Thr	Pro	Gly	Pro	Arg	Glu	Ala	Leu	Ser	Arg	Leu	Gln	Glu	Leu	Cys
				65					70					75
Tyr	Gln	Trp	Leu	Met	Pro	Glu	Leu	His	Thr	Lys	Glu	Gln	Ile	Leu
				80					85					90
Glu	Leu	Leu	Val	Leu	Glu	Gln	Phe	Leu	Ser	Ile	Leu	Pro	Glu	Glu
				95					100					105
Leu	Gln	Ile	Trp	Val	Gln	Gln	His	Asn	Pro	Glu	Ser	Gly	Glu	Glu
				110					115					120
Ala	Val	Thr	Leu	Leu	Glu	Asp	Leu	Glu	Arg	Glu	Phe	Asp	Asp	Pro
				125					130					135
Gly	Gln	Gln	Val	Pro	Ala	Ser	Pro	Gln	Gly	Pro	Ala	Val	Pro	Trp
				140					145					150
Lys	Asp	Leu	Thr	Cys	Leu	Arg	Ala	Ser	Gln	Glu	Ser	Thr	Asp	Ile
				155					160					165
His	Leu	Gln	Pro	Leu	Lys	Thr	Gln	Leu	Lys	Ser	Trp	Lys	Pro	Cys
				170					175					180
Leu	Ser	Pro	Lys	Ser	Asp	Cys	Glu	Asn	Ser	Glu	Thr	Ala	Thr	Lys
				185					190					195
Glu	Gly	Ile	Ser	Glu	Glu	Lys	Ser	Gln	Gly	Leu	Pro	Gln	Glu	Pro
				200					205					210
Ser	Phe	Arg	Gly	Ile	Ser	Glu	His	Glu	Ser	Asn	Leu	Val	Trp	Lys
				215					220					225
Gln	Gly	Ser	Ala	Thr	Gly	Glu	Lys	Leu	Arg	Ser	Pro	Ser	Gln	Gly
				230					235					240
Gly	Ser	Phe	Ser	Gln	Val	Ile	Phe	Thr	Asn	Lys	Ser	Leu	Gly	Lys
				245					250					255
Arg	Asp	Leu	Tyr	Asp	Glu	Ala	Glu	Arg	Cys	Leu	Ile	Leu	Thr	Thr
				260					265					270
Asp	Ser	Ile	Met	Cys	Gln	Lys	Val	Pro	Pro	Glu	Glu	Arg	Pro	Tyr
				275					280					285
Arg	Cys	Asp	Val	Cys	Gly	His	Ser	Phe	Lys	Gln	His	Ser	Ser	Leu
				290					295					300
Thr	Gln	His	Gln	Arg	Ile	His	Thr	Gly	Glu	Lys	Pro	Tyr	Lys	Cys
				305					310					315
Asn	Gln	Cys	Gly	Lys	Ala	Phe	Ser	Leu	Arg	Ser	Tyr	Leu	Ile	Ile
				320					325					330
His	Gln	Arg	Ile	His	Ser	Gly	Glu	Lys	Ala	Tyr	Glu	Cys	Ser	Glu
				335					340					345
Cys	Gly	Lys	Ala	Phe	Asn	Gln	Ser	Ser	Ala	Leu	Ile	Arg	His	Arg
				350					355					360
Lys	Ile	His	Thr	Gly	Glu	Lys	Ala	Cys	Lys	Cys	Asn	Glu	Cys	Gly
				365					370					375
Lys	Ala	Phe	Ser	Gln	Ser	Ser	Tyr	Leu	Ile	Ile	His	Gln	Arg	Ile
				380					385					390
His	Thr	Gly	Glu	Lys	Pro	Tyr	Glu	Cys	Asn	Glu	Cys	Gly	Lys	Thr
				395					400					405
Phe	Ser	Gln	Ser	Ser	Lys	Leu	Ile	Arg	His	Gln	Arg	Ile	His	Thr
				410					415					420
Gly	Glu	Arg	Pro	Tyr	Glu	Cys	Asn	Glu	Cys	Gly	Lys	Ala	Phe	Arg
				425					430					435
Gln	Ser	Ser	Glu	Leu	Ile	Thr	His	Gln	Arg	Ile	His	Ser	Gly	Glu
				440					445					450
Lys	Pro	Tyr	Glu	Cys	Ser	Glu	Cys	Gly	Lys	Ala	Phe	Ser	Leu	Ser

	455		460		465
Ser Asn Leu Ile	Arg His Gln Arg Ile	His Ser Gly Glu Glu	Pro		
	470		475		480
Tyr Gln Cys Asn	Glu Cys Gly Lys Thr	Phe Lys Arg Ser Ser	Ala		
	485		490		495
Leu Val Gln His	Gln Arg Ile His Ser	Gly Asp Glu Ala Tyr	Ile		
	500		505		510
Cys Asn Glu Cys	Gly Lys Ala Phe Arg	His Arg Ser Val Leu	Met		
	515		520		525
Arg His Gln Arg	Val His Thr Ile Lys				
	530				

<210> 47
 <211> 206
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3282941CD1

<400> 47

Met Ser Thr Gly Ser Val Ser Asp Pro Glu Glu Met Glu Leu Arg	
1 5 10 15	
Gly Leu Gln Arg Glu Tyr Pro Val Pro Ala Ser Lys Arg Pro Pro	
20 25 30	
Leu Arg Gly Val Glu Arg Ser Tyr Ala Ser Pro Ser Asp Asn Ser	
35 40 45	
Ser Ala Glu Glu Glu Asp Pro Asp Gly Glu Glu Arg Cys Ala	
50 55 60	
Leu Gly Thr Ala Gly Ser Ala Glu Gly Cys Lys Arg Lys Arg Pro	
65 70 75	
Arg Val Ala Gly Gly Gly Ala Gly Gly Ser Ala Gly Gly Gly	
80 85 90	
Gly Lys Lys Pro Leu Pro Ala Lys Gly Ser Ala Ala Glu Cys Lys	
95 100 105	
Gln Ser Gln Arg Asn Ala Ala Asn Ala Arg Glu Arg Ala Arg Met	
110 115 120	
Arg Val Leu Ser Lys Ala Phe Ser Arg Leu Lys Thr Ser Leu Pro	
125 130 135	
Trp Val Pro Pro Asp Thr Lys Leu Ser Lys Leu Asp Thr Leu Arg	
140 145 150	
Leu Ala Ser Ser Tyr Ile Ala His Leu Arg Gln Leu Leu Gln Glu	
155 160 165	
Asp Arg Tyr Glu Asn Gly Tyr Val His Pro Val Asn Leu Thr Trp	
170 175 180	
Pro Phe Val Val Ser Gly Arg Pro Asp Ser Asp Thr Lys Glu Val	
185 190 195	
Ser Ala Ala Asn Arg Leu Cys Gly Thr Thr Ala	
200 205	

<210> 48
 <211> 172
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3286656CD1

<400> 48
 Met Glu Ser Val Thr Phe Glu Asp Val Ala Val Glu Phe Ile Gln
 1 5 10 15
 Glu Trp Ala Leu Leu Asp Ser Ala Arg Arg Ser Leu Cys Lys Tyr
 20 25 30
 Arg Met Leu Asp Gln Cys Arg Thr Leu Ala Ser Arg Gly Thr Pro
 35 40 45
 Pro Cys Lys Pro Ser Cys Val Ser Gln Leu Gly Gln Arg Ala Glu
 50 55 60
 Pro Lys Ala Thr Glu Arg Gly Ile Leu Arg Ala Thr Gly Val Ala
 65 70 75
 Trp Glu Ser Gln Leu Lys Pro Glu Glu Leu Pro Ser Met Gln Asp
 80 85 90
 Leu Leu Glu Glu Ala Ser Ser Arg Asp Met Gln Met Gly Pro Gly
 95 100 105
 Leu Phe Leu Arg Met Gln Leu Val Pro Ser Ile Glu Glu Arg Glu
 110 115 120
 Thr Pro Leu Thr Arg Glu Asp Arg Pro Ala Leu Gln Glu Pro Pro
 125 130 135
 Trp Ser Leu Gly Cys Thr Gly Leu Lys Ala Ala Met Gln Ile Gln
 140 145 150
 Arg Val Val Ile Pro Val Pro Thr Leu Gly His Arg Asn Pro Trp
 155 160 165
 Val Ala Arg Asp Ser Ala Met
 170

<210> 49
 <211> 275
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3490802CD1

<400> 49
 Met Gly Pro Leu Gln Phe Arg Asp Val Ala Ile Glu Phe Ser Leu
 1 5 10 15
 Glu Glu Trp His Cys Leu Asp Thr Ala Gln Arg Asn Leu Tyr Arg
 20 25 30
 Asp Val Met Leu Glu Asn Tyr Arg Asn Leu Val Phe Leu Gly Ile
 35 40 45
 Val Val Ser Lys Pro Asp Leu Val Thr Cys Leu Glu Gln Gly Lys
 50 55 60
 Lys Pro Leu Thr Met Glu Arg His Glu Met Ile Ala Lys Pro Pro
 65 70 75
 Val Met Ser Ser His Phe Ala Gln Asp Leu Trp Pro Glu Asn Ile
 80 85 90
 Gln Asn Ser Phe Gln Ile Gly Met Leu Arg Arg Tyr Glu Glu Cys
 95 100 105
 Arg His Asp Asn Leu Gln Leu Lys Lys Gly Cys Lys Ser Val Gly
 110 115 120
 Glu His Lys Val His Lys Gly Gly Tyr Asn Gly Leu Asn Gln Cys
 125 130 135
 Leu Thr Thr Thr Gln Lys Glu Ile Phe Gln Cys Asp Lys Tyr Gly

Lys Val Phe His	140	Lys Phe Ser Asn Ser	145	Asn Thr Tyr Lys Thr	150
	155		160		165
His Thr Gly Ile	Asn Leu Phe Lys Cys	Ile Ile Cys Gly Lys	Ala		
	170		175		180
Phe Lys Arg Ser	Ser Thr Leu Thr Thr	His Lys Lys Ile His	Thr		
	185		190		195
Gly Glu Lys Pro	Tyr Lys Cys Glu Glu	Cys Gly Lys Ala Phe	Asn		
	200		205		210
Gln Ser Ser Asn	Leu Thr Thr His Lys	Arg Ile His Thr Gly	Glu		
	215		220		225
Lys Pro Tyr Lys	Cys Glu Glu Cys Gly	Lys Ala Phe Asn Trp	Ser		
	230		235		240
Ser Asp Leu Asn	Lys His Lys Lys Ile	His Ile Glu Arg Lys	Pro		
	245		250		255
Tyr Ile Val Lys	Asn Val Thr Asp Leu	Leu Asn Val Pro Pro	Leu		
	260		265		270
Leu Ile Ser Ile	Arg				
	275				

<210> 50
 <211> 236
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3507366CD1

<400> 50

Met Asp Lys Arg Tyr	Leu Gln Phe Asp	Ile Lys Ala Phe Val	Glu
1	5	10	15
Asn Asn Pro Ala Ile	Lys Trp Cys Pro	Thr Pro Gly Cys Asp	Arg
	20	25	30
Ala Val Arg Leu Thr	Lys Gln Gly Ser	Asn Thr Ser Gly Ser	Asp
	35	40	45
Thr Leu Ser Phe Pro	Leu Leu Arg Ala	Pro Ala Val Asp Cys	Gly
	50	55	60
Lys Gly His Leu Phe	Cys Trp Glu Cys	Leu Gly Glu Ala His	Glu
	65	70	75
Pro Cys Asp Cys Gln	Thr Trp Lys Asn	Trp Leu Gln Lys Ile	Thr
	80	85	90
Glu Met Lys Pro Glu	Glu Leu Val Gly	Val Ser Glu Ala Tyr	Glu
	95	100	105
Asp Ala Ala Asn Cys	Leu Trp Leu Leu	Thr Asn Ser Lys Pro	Cys
	110	115	120
Ala Asn Cys Lys Ser	Pro Ile Gln Lys	Asn Glu Gly Cys Asn	His
	125	130	135
Met Gln Cys Ala Lys	Cys Lys Tyr Asp	Phe Cys Trp Ile Cys	Leu
	140	145	150
Glu Glu Trp Lys Lys	His Ser Ser Ser	Thr Gly Gly Tyr Tyr	Arg
	155	160	165
Cys Thr Arg Tyr Glu	Val Ile Gln His	Val Glu Glu Gln Ser	Lys
	170	175	180
Glu Met Thr Val Glu	Ala Glu Lys Lys	His Lys Arg Phe Gln	Glu
	185	190	195
Leu Asp Arg Phe Met	His Tyr Tyr Thr	Arg Phe Lys Asn His	Glu
	200	205	210
His Ser Tyr Gln Leu	Glu Gln Arg Leu	Leu Lys Thr Ala Lys	Glu

	215		220	225
Lys Met Glu Gln Met Ser Arg Val Ser Lys Asn				
	230		235	

<210> 51
 <211> 214
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3573060CD1

<400> 51

Met Asn Leu Ser Ser Ala Ser Ser Thr Glu Glu Lys Ala Val Thr	
1 5 10 15	
Thr Val Leu Trp Gly Cys Glu Leu Ser Gln Glu Arg Arg Thr Trp	
20 25 30	
Thr Phe Arg Pro Gln Leu Glu Gly Lys Gln Ser Cys Arg Leu Leu	
35 40 45	
Leu His Thr Ile Cys Leu Gly Glu Lys Ala Lys Glu Glu Met His	
50 55 60	
Arg Val Glu Ile Leu Pro Pro Ala Asn Gln Glu Asp Lys Lys Met	
65 70 75	
Gln Pro Val Thr Ile Ala Ser Leu Gln Ala Ser Val Leu Pro Met	
80 85 90	
Val Ser Met Val Gly Val Gln Leu Ser Pro Pro Val Thr Phe Gln	
95 100 105	
Leu Arg Ala Gly Ser Gly Pro Val Phe Leu Ser Gly Gln Glu Arg	
110 115 120	
Tyr Glu Ala Ser Asp Leu Thr Trp Glu Glu Glu Glu Glu Glu	
125 130 135	
Gly Glu Glu Glu Glu Glu Glu Glu Asp Asp Glu Asp Glu Asp	
140 145 150	
Ala Asp Ile Ser Leu Glu Glu Gln Ser Pro Val Lys Gln Val Lys	
155 160 165	
Arg Leu Val Pro Gln Lys Gln Ala Ser Val Ala Lys Lys Lys Lys	
170 175 180	
Leu Glu Lys Glu Glu Glu Glu Ile Arg Ala Ser Val Arg Asp Lys	
185 190 195	
Ser Pro Val Lys Lys Ala Lys Ala Thr Ala Arg Ala Lys Lys Pro	
200 205 210	
Gly Phe Lys Lys	

<210> 52
 <211> 396
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3573661CD1

<400> 52
 Met Asn Phe Thr Val Gly Phe Lys Pro Leu Leu Gly Asp Ala His

1	5	10	15
Ser Met Asp Asn Leu	Glu Lys Gln Leu	Ile Cys Pro Ile Cys Leu	
	20	25	30
Glu Met Phe Ser Lys	Pro Val Val Ile	Leu Pro Cys Gln His Asn	
	35	40	45
Leu Cys Arg Lys Cys	Ala Asn Asp Val	Phe Gln Ala Ser Asn Pro	
	50	55	60
Leu Trp Gln Ser Arg	Gly Ser Thr Thr	Val Ser Ser Gly Gly Arg	
	65	70	75
Phe Arg Cys Pro Ser	Cys Arg His Glu	Val Val Leu Asp Arg His	
	80	85	90
Gly Val Tyr Gly Leu	Gln Arg Asn Val	Leu Val Glu Asn Ile Ile	
	95	100	105
Asp Ile Tyr Lys Gln	Glu Ser Ser Lys	Pro Leu His Ser Lys Ala	
	110	115	120
Glu Gln His Leu Met	Cys Glu Glu His	Glu Glu Glu Lys Ile Asn	
	125	130	135
Ile Tyr Cys Leu Ser	Cys Glu Val Pro	Thr Cys Ser Leu Cys Lys	
	140	145	150
Val Phe Gly Ala His	Lys Asp Cys Glu	Val Ala Pro Leu Pro Thr	
	155	160	165
Ile Tyr Lys Arg Gln	Lys Ser Glu Leu	Ser Asp Gly Ile Ala Met	
	170	175	180
Leu Val Ala Gly Asn	Asp Arg Val Gln	Ala Val Ile Thr Gln Met	
	185	190	195
Glu Glu Val Cys Gln	Thr Ile Glu Asp	Asn Ser Arg Arg Gln Lys	
	200	205	210
Gln Leu Leu Asn Gln	Arg Phe Glu Ser	Leu Cys Ala Val Leu Glu	
	215	220	225
Glu Arg Lys Gly Glu	Leu Leu Gln Ala	Leu Ala Arg Glu Gln Glu	
	230	235	240
Glu Lys Leu Gln Arg	Val Arg Gly Leu	Ile Arg Gln Tyr Gly Asp	
	245	250	255
His Leu Glu Ala Ser	Ser Lys Leu Val	Glu Ser Ala Ile Gln Ser	
	260	265	270
Met Glu Glu Pro Gln	Met Ala Leu Tyr	Leu Gln Gln Ala Lys Glu	
	275	280	285
Leu Ile Asn Lys Val	Gly Ala Met Ser	Lys Val Glu Leu Ala Gly	
	290	295	300
Arg Pro Glu Pro Gly	Tyr Glu Ser Met	Glu Gln Phe Thr Val Arg	
	305	310	315
Val Glu His Val Ala	Glu Met Leu Arg	Thr Ile Asp Phe Gln Pro	
	320	325	330
Gly Ala Ser Gly Gly	Gly Arg Gly Gly	Gly Pro Arg Arg Lys Lys	
	335	340	345
Arg Ala Thr Arg Gly	Pro Glu Glu Lys	Thr Ala Arg Met Gly Pro	
	350	355	360
Tyr Arg Pro Leu Arg	Pro Asn Pro Asp	Pro Leu Leu Arg Lys Ser	
	365	370	375
Pro Arg Arg Leu Arg	Ile Ser Gly Gly	Arg Asn Ser Cys Arg Lys	
	380	385	390
Lys Thr Pro Ala Ser	Phe		
	395		

<210> 53
 <211> 486
 <212> PRT
 <213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 3633422CD1

<400> 53

```

Met Arg Arg Leu Val His Asp Leu Leu Pro Pro Glu Val Cys Ser
 1      5      10      15
Leu Leu Asn Pro Ala Ala Ile Tyr Ala Asn Asn Glu Ile Ser Leu
 20      25      30
Arg Asp Val Glu Val Tyr Gly Phe Asp Tyr Asp Tyr Thr Leu Ala
 35      40      45
Gln Tyr Ala Asp Ala Leu His Pro Glu Ile Phe Ser Thr Ala Arg
 50      55      60
Asp Ile Leu Ile Glu His Tyr Lys Tyr Pro Glu Gly Ile Arg Lys
 65      70      75
Tyr Asp Tyr Asn Pro Ser Phe Ala Ile Arg Gly Leu His Tyr Asp
 80      85      90
Ile Gln Lys Ser Leu Leu Met Lys Ile Asp Ala Phe His Tyr Val
 95      100     105
Gln Leu Gly Thr Ala Tyr Arg Gly Leu Gln Pro Val Pro Asp Glu
 110     115     120
Glu Val Ile Glu Leu Tyr Gly Gly Thr Gln His Ile Pro Leu Tyr
 125     130     135
Gln Met Ser Gly Phe Tyr Gly Lys Gly Pro Ser Ile Lys Gln Phe
 140     145     150
Met Asp Ile Phe Ser Leu Pro Glu Met Ala Leu Leu Ser Cys Val
 155     160     165
Val Asp Tyr Phe Leu Gly His Ser Leu Glu Phe Asp Gln Ala His
 170     175     180
Leu Tyr Lys Asp Val Thr Asp Ala Ile Arg Asp Val His Val Lys
 185     190     195
Gly Leu Met Tyr Gln Trp Ile Glu Gln Asp Met Glu Lys Tyr Ile
 200     205     210
Leu Arg Gly Asp Glu Thr Phe Ala Val Leu Ser Arg Leu Val Ala
 215     220     225
His Gly Lys Gln Leu Phe Leu Ile Thr Asn Ser Pro Phe Ser Phe
 230     235     240
Val Asp Lys Gly Met Arg His Met Val Gly Pro Asp Trp Arg Gln
 245     250     255
Leu Phe Asp Val Val Ile Val Gln Ala Asp Lys Pro Ser Phe Phe
 260     265     270
Thr Asp Arg Arg Lys Pro Phe Arg Lys Leu Asp Glu Lys Gly Ser
 275     280     285
Leu Gln Trp Asp Arg Ile Thr Arg Leu Glu Lys Gly Lys Ile Tyr
 290     295     300
Arg Gln Gly Asn Leu Phe Asp Phe Leu Arg Leu Thr Glu Trp Arg
 305     310     315
Gly Pro Arg Val Leu Tyr Phe Gly Asp His Leu Tyr Ser Asp Leu
 320     325     330
Ala Asp Leu Met Leu Arg His Gly Trp Arg Thr Gly Ala Ile Ile
 335     340     345
Pro Glu Leu Glu Arg Glu Ile Arg Ile Ile Asn Thr Glu Gln Tyr
 350     355     360
Met His Ser Leu Thr Trp Gln Gln Ala Leu Thr Gly Leu Leu Glu
 365     370     375
Arg Met Gln Thr Tyr Gln Asp Ala Glu Ser Arg Gln Val Leu Ala
 380     385     390
Ala Trp Met Lys Glu Arg Gln Glu Leu Arg Cys Ile Thr Lys Ala
 395     400     405
Leu Phe Asn Ala Gln Phe Gly Ser Ile Phe Arg Thr Phe His Asn
 410     415     420
Pro Thr Tyr Phe Ser Arg Arg Leu Val Arg Phe Ser Asp Leu Tyr

```

Met	Ala	Ser	Leu	425				430				435		
			Ser		Cys	Leu	Leu	Asn	Tyr	Arg	Val	Asp	Phe	Thr
				440					445					450
Phe	Tyr	Pro	Arg	Arg	Thr	Pro	Leu	Gln	His	Glu	Ala	Pro	Leu	Trp
				455					460					465
Met	Asp	Gln	Leu	Cys	Thr	Gly	Cys	Met	Lys	Thr	Pro	Phe	Leu	Gly
				470					475					480
Asp	Met	Ala	His	Ile	Arg									
				485										

<210> 54
 <211> 555
 <212> PRT
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3993377CD1

<400> 54

Met	Gly	Ala	Glu	Asp	Lys	Leu	Pro	Leu	Glu	Asp	Ser	Pro	Val	Ile
1				5					10					15
Ala	Ala	Leu	Asp	Cys	Pro	Ser	Leu	Asn	Asn	Ala	Thr	Ala	Phe	Ser
				20					25					30
Leu	Leu	Ala	Asp	Asp	Ser	Gln	Thr	Ser	Thr	Ser	Ile	Phe	Ala	Ser
				35					40					45
Pro	Thr	Ser	Pro	Pro	Val	Leu	Gly	Glu	Ser	Val	Leu	Gln	Asp	Asn
				50					55					60
Ser	Phe	Asp	Leu	Asn	Asn	Gly	Ser	Asp	Ala	Glu	Gln	Glu	Glu	Met
				65					70					75
Glu	Thr	Gln	Ser	Ser	Asp	Phe	Pro	Pro	Ser	Leu	Thr	Gln	Pro	Ala
				80					85					90
Pro	Asp	Gln	Ser	Ser	Thr	Ile	Gln	Leu	His	Pro	Ala	Thr	Ser	Pro
				95					100					105
Ala	Val	Ser	Pro	Thr	Thr	Ser	Pro	Ala	Val	Ser	Leu	Val	Val	Ser
				110					115					120
Pro	Ala	Ala	Ser	Pro	Glu	Ile	Ser	Pro	Glu	Val	Cys	Pro	Ala	Ala
				125					130					135
Ser	Thr	Val	Val	Ser	Pro	Ala	Val	Phe	Ser	Val	Val	Ser	Pro	Ala
				140					145					150
Ser	Ser	Ala	Val	Leu	Pro	Ala	Val	Ser	Leu	Glu	Val	Pro	Leu	Thr
				155					160					165
Ala	Ser	Val	Thr	Ser	Pro	Lys	Ala	Ser	Pro	Val	Thr	Ser	Pro	Ala
				170					175					180
Ala	Ala	Phe	Pro	Thr	Ala	Ser	Pro	Ala	Asn	Lys	Asp	Val	Ser	Ser
				185					190					195
Phe	Leu	Glu	Thr	Thr	Ala	Asp	Val	Glu	Glu	Ile	Thr	Gly	Glu	Gly
				200					205					210
Leu	Thr	Ala	Ser	Gly	Ser	Gly	Asp	Val	Met	Arg	Arg	Arg	Ile	Ala
				215					220					225
Thr	Pro	Glu	Glu	Val	Arg	Leu	Pro	Leu	Gln	His	Gly	Trp	Arg	Arg
				230					235					240
Glu	Val	Arg	Ile	Lys	Asn	Ser	Ser	His	Arg	Trp	Gln	Gly	Glu	Thr
				245					250					255
Trp	Tyr	Tyr	Gly	Pro	Cys	Gly	Lys	Arg	Met	Lys	Gln	Phe	Pro	Glu
				260					265					270
Val	Ile	Lys	Tyr	Leu	Ser	Arg	Asn	Val	Val	His	Ser	Val	Arg	Arg
				275					280					285
Glu	His	Phe	Ser	Phe	Ser	Pro	Arg	Met	Pro	Val	Gly	Asp	Phe	Phe

	290		295		300
Glu Glu Arg Asp	Thr Pro Glu Gly Leu	Gln Trp Val Gln Leu Ser			
	305		310		315
Ala Glu Glu Ile	Pro Ser Arg Ile Gln	Ala Ile Thr Gly Lys Arg			
	320		325		330
Gly Arg Pro Arg	Asn Thr Glu Lys Ala	Lys Thr Lys Glu Val Pro			
	335		340		345
Lys Val Lys Arg	Gly Arg Gly Arg Pro	Pro Lys Val Lys Ile Thr			
	350		355		360
Glu Leu Leu Asn	Lys Thr Asp Asn Arg	Pro Leu Lys Lys Leu Glu			
	365		370		375
Ala Gln Glu Thr	Leu Asn Glu Glu Asp	Lys Ala Lys Ile Ala Lys			
	380		385		390
Ser Lys Lys Lys	Met Arg Gln Lys Val	Gln Arg Gly Glu Cys Gln			
	395		400		405
Thr Thr Ile Gln	Gly Gln Ala Arg Asn	Lys Arg Lys Gln Glu Thr			
	410		415		420
Lys Ser Leu Lys	Gln Lys Glu Ala Lys	Lys Lys Ser Lys Ala Glu			
	425		430		435
Lys Glu Lys Gly	Lys Thr Lys Gln Glu	Lys Leu Lys Glu Lys Val			
	440		445		450
Lys Arg Glu Lys	Lys Glu Lys Val Lys	Met Lys Glu Lys Glu Glu			
	455		460		465
Val Thr Lys Ala	Lys Pro Ala Cys Lys	Ala Asp Lys Thr Leu Ala			
	470		475		480
Thr Gln Arg Arg	Leu Glu Glu Arg Gln	Arg Gln Gln Met Ile Leu			
	485		490		495
Glu Asp Met Lys	Lys Pro Thr Glu Asp	Met Cys Leu Thr Asp His			
	500		505		510
Gln Pro Leu Pro	Asp Phe Ser Arg Val	Pro Gly Leu Thr Leu Pro			
	515		520		525
Ser Gly Ala Phe	Ser Asp Cys Leu Thr	Ile Val Glu Phe Leu His			
	530		535		540
Ser Phe Gly Lys	Val Leu Gly Leu Asp	Pro Ala Gln Gly Cys Ala			
	545		550		555

<210> 55

<211> 61

<212> PRT

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 4717936CD1

<400> 55

Met Gln Pro Arg Thr Gln Pro Leu Ala Gln Thr Leu Pro Phe Phe		
1	5	10
Leu Gly Gly Ala Pro Arg Asp Thr Gly Leu Arg Val Pro Val Ile		
20	25	30
Lys Met Gly Thr Gly Trp Glu Gly Phe Gln Arg Thr Leu Lys Glu		
35	40	45
Val Ala Tyr Ile Leu Leu Cys Cys Trp Cys Ile Lys Glu Leu Leu		
50	55	60
Asp		

<210> 56
<211> 2781
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 025733CB1

<400> 56
gaagtggggt gcacgcttcg ggttgggtgc atggcagctg cggggagccg caagaggcgc 60
ctggcgagc tgacgggtgga cgagttccta gcttcgggct ttgactccga gtccgaatcc 120
gagtcgcaaa attctccaca agcggagaca cgggaagcac gcgaggctgc ccggagtccg 180
gataagccgg gcgggagccc ctcgccagc cggcgtaaag gccgtgcctc tgagcacaaa 240
gaccagctct ctcggtgaa ggacagagac cccgagttct acaagttcct gcaggagaa 300
gaccagagcc tgctaaactt cagcgactcg gacagctctg aggaggaaga ggggcccgtc 360
cactccctgc catagtgtgt ggaggaaagc agtgaggagg aggatggagc ggaggaagga 420
gaagatgggg acagagtccc cagagggtcg aagggggaaga agaattctgt tcctgtgacc 480
gtcgccatgg ttgagagatg gaagcaggca gcaaagcaac gcctcactcc aaagctgttc 540
catgaagtgg tacaggcggt cccagcagct gtggccacca cccgagggga ccaggaaagt 600
gctgaggcca acaaattcca ggtcacggac agtgctgcat tcaatgctct ggttaccttc 660
tgcatcagag acctcattgg ctgtctccag aagctgctgt ttggaaaggt ggcaaaaggt 720
agcagcagga tgctgcagcc gtccagcagc ccgctctggg ggaagcttcg tgtggacatc 780
aaggcttacc tgggctcggc catacagctg gtgtcctgtc tgtcggagac gacggtgttg 840
gcgcccgctg tgccggacat cagcgtgtct gtgcctgtct tcctgacctt ccccaagcag 900
tgccgcatgc tectcaagag aatggtgggt gtagggagca ctggggagga gtctctgcgg 960
gtgctgggtt tcctgtctct cagcagagtc tgccggcaca agaaggacac ttctctggc 1020
cccgctctca agcaaatgta catcacgtat gtgaggaaact gcaagttcac ctgcctgggt 1080
gccctccctt tcatcagttt catgcagttg accttgacgg agctgctggc cctggagccg 1140
ggtgtggcct accagcagc ctctctctac atccgccagc tcgccataca cctgcgcaac 1200
gccatgacca cccgcaagaa ggaacatac cagtctgtgt acaactggca gtatgtgcac 1260
tgctctctcc tgtgtgctcg ggtcctgagc actgcgggccc ccagcgaagc cctccagccc 1320
ttggtctacc ccttgcccca agtcacatt ggctgtatca agctcatccc cactgcccgc 1380
ttctaccgct tgcaatgca ctgcacogt gcctgacgc tgctctcggt gagctcggtg 1440
gccttcaccc tgaagtctgt cctccgggag tgcaaggtgg ccaactactg ccggcaggtg 1500
aagccagggc gcatgagctc caagcccatc aacttctccg tgatcctgaa gctgtccaat 1560
gtcaacctgc aggagaaggg gtaccgggac ggctgtgtgg agcagctgta cgacctcacc 1620
ctggagtacc tgcaagacca ggcacactgc atcggtcttc cggagctggt gctgctgtg 1680
gtcctgcagc tgaagtctgt cctccgggag tgcaaggtgg ccaactactg ccggcaggtg 1740
cagcagctgc ttgggaaggt tcaggagaac tcggcataca tctgcagccg ccgccagagg 1800
gtttcctctg gctctcttga gcagcaggca gtggaagcct gggagaagct gaccgggaa 1860
gaggggacac ccttgacctt gtactacagc cactggcgca agctgcgtga ccgggagatc 1920
cagctggaga tcagtggcaa agagcggctg gaagacctga acttccctga gatcaaacga 1980
aggaagatgg ctgacaggaa ggatgaggac aggaagcaat ttaaagacct ctttgacctg 2040
aacagctctg aagaggacga caccgaggga ttctcggaga gagggatact gaggcccctg 2100
agcactcggc atgggggtgga agacgatgaa gaggacgagg aggagggcga ggaggacagc 2160
agcaactcgg aggtggaatg gtcttgggat ggagaccag acgcagaggg ggggctggcc 2220
cctggggagc tgccagagct ggcccagggg cgggaggacg agctggagga tctgcagctc 2280
tcaggaggac actgaggcag cccatctggg gggcctgtag gggctgcccg gctggtggcc 2340
agtgtttcca cctccctggc agtcaggcct agaggctggc gtctgtgcag ttgggggagg 2400
cagtagacac gggacaggct ttattattta tttttcagca tgaaagacca aacgtatcga 2460
gagctgggct gggctgggct ggtgtggctg ctgaagcccc acagctgtgg gctgctgaag 2520
tcagctccgc gggggagctg accctgacgt cagcagaccg agaccagtcc cagttccagg 2580
gggaggcctg cagggccctg gccctttcca ccacctctgc cctccgtctg cagacctctg 2640
ccatctgcac caggtctctg cttcactccc ccaagtcttt gaaaatttgt tcctttcctt 2700
tgaagtccca tttcttttta aaattttttg ttttgcaccc gaaaccgaaa gaaataaagc 2760
ggtgggaggg aaaaaaaaaa a 2781

<210> 57
<211> 2544
<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 079702CB1

<400> 57

```
cgggaaacca aaatggcgag gggctgtatt gaagtgggct gtgtttgagg ccgggtgtaag 60
aacgctcatt ctacccccaa ccttctgtct caaggacctc ggtttgtgcg tgcataatgtg 120
ccgggtaccc ggtggggcgg gtgcccagta agtgctcgga ctgcagggg aagcgccac 180
ggggacggat tgggtgtttt ttctgtatg aagcggttgg caccactgaa gtgaccgaat 240
gagagactct acaggggcag gtaattcact ggtccacaag cggctctcct tacgtcgaaa 300
ccaaaagacc ccaacatcct tgaccaagct gtctttacag gatggacata aagccaaaaa 360
gccagcatgt aaatttgaag agggtcagga tgtcctagct agatggtcag atggcttggt 420
ttatcttggc actatcaaaa agataaacat attgaaacag agctgcttca tcatatttga 480
agacagttct aaatcctggg ttctctggaa ggacattcaa acaggagcca ctggaagtgg 540
ggaaatggtc tgtacaatat gtcaagaaga gtattcagaa gctcccaatg aaatggttat 600
atgtgacaag tgtggccaag gatcatca gttgtgtcac acacctcata ttgattccag 660
tgtgattgat tcagatgaaa aatggctctg tcggcagtggt gtttttgcaa caacaacaaa 720
gaggggtggt gcacttaaga aaggacaaa tgccaaagca ttgcaagtca tgaagcagac 780
attaccctat agtgtggcag accttgaatg ggaatgcagg cataaaacca atgtccagca 840
gtgttactgc tattgtggag gccctggaga ctggtatttg aagatgctac agtgctgcaa 900
atgtaagcag tggtttcatg aggccttggt gcaatgcctt caaaagccaa tgctatttgg 960
agacagattt tatacgttta tatgctctgt ctgcagttct ggaccagaat acctcaaacg 1020
totaccatta cagtgggtag atatagcaca cctatgcctt tacaacctaa gtgttattca 1080
taagaagaaa tactttgatt ctgaacttga gcttatgaca tacattaatg aaaactggga 1140
tagattgcac cctggagagc tggcagacac accaaaatct gaaagatatg agcatgttct 1200
ggaggcatta aatgattaca agaccatgtt tatgtctggg aaagaaataa agaagaagaa 1260
gcatttgttt gggttgcgaa ttctgtgtcc tctgttgcca ccaaatgtgg ctttcaaacg 1320
agagaaagaa cctgaaggaa catctcatga atttaaaatt aaaggcagaa aggcattcaa 1380
acctatatct gattcaaggg aagtaagcaa tggcatagaa aaaaaaaaaa aaaaaaaatc 1440
tgtaggtcgt ccacctggcc catatacaag aaaaatgatt caaaaaactg ctgagccact 1500
tttgataaag gaatcaattt cagagaatcc tactttggat ttaccttggt ctatagggag 1560
aactgagga actgcacatt catccaatac ctccagatgt gatttcacgg gtgcttccag 1620
tgcaaaaagaa actacctcgt ctagcatttc caggcattat ggattatctg actccagaaa 1680
aagaacgcgt acaggaagat cttggcctgc tgcaatacca catttgcgga gaagaagagg 1740
tcgtcttcca agaagagcac tccagactca gaactcagaa attgtaaaag atgatgaagg 1800
caaagaagat tatcagtttg atgaactcaa cacagagatt ctgaataact tagcagatca 1860
ggagttacaa ctcaatcac taaagaactc cattaccagt tatttttggt ctgcaggtag 1920
aatagcatgt ggcgaaaaat accgagtttt ggcacgtcgg gtgacacttg atggaaaggt 1980
gcagtatcct gtggaatggg aaggagcaac tgcacactga ctgtaggact gaacattatg 2040
ttcactgcac tctgattttc ttaggttaca gttcaaagcc cttaaaggag ctggctttta 2100
ctatctttct taaaaaaaaa aaaaagtcaa aaaaattcaa aaaaggggat gatactagcc 2160
ttaacatgta cctgtcaatg ttatggatat tgtcataaaa aggtatcttt taaaaatcag 2220
aacagagact taatttttta aatcttaaga ttgttagaat gtttctagga taggatatta 2280
aaaatgattg aaacccatgc atgggtgttag acaatttttc taattattcc attgagtcag 2340
ttttttgtga ttagtgatta tcagagcaaa catcatgtag atagcacaag tatttgaga 2400
aacgttgttt gttttgttac caaaatgttg gaaaaattta tttcaatacc ttttagattt 2460
cataaagtgc agtgtatata atgcctactg aaagactgta aaatattgaa attttctttc 2520
aagcaaatgt taaataaata tatic 2544
```

<210> 58

<211> 1627

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 116208CB1

<400> 58

```

ctgatgaggg cgctgcattt attgaagagc ggctgcagcc ctgcggttca gattaaaatc 60
cgagaattgt atagacgcgc atatccacga actcttgaag gactttctga tttatccaca 120
atcaaatcat cggttttcag tttggatggg ggctcatcac ctgtagaacc tgacttggcc 180
gtggctggaa tccactcggt gccttccact tcagttacac ctactcacc atcctctcct 240
gttggttctg tgctgcttca agatactaag cccacatttg agatgcagca gccatctccc 300
ccaattcttc ctgtccatcc tgatgtgcag ttaaaaaatc tgccctttta tgatgtcctt 360
gatgttttca tcaagcccac gagtttagtt caaagcagta ttcagcgatt tcaagagaag 420
ttttttattt ttgctttgac acctcaacaa gttagagaga tatgcatac cagggatttt 480
ttgccaggtg gtaggagaga ttatacagtc caagttcagt tgagactttg cctggcagag 540
acaagttgcc ctcaagaaga taactatcca aatagtctat gtataaaagt aaatgggaag 600
ctatttctct tgcttggtca tgcaccaccg cctaaaaatg ggattgaaca gaagcgccct 660
ggacgcccct tgaattattac atcttttagtt aggttatctt cagctgtgcc aaaccaaatt 720
tccatttctt gggcatcaga aattgggaag aattactcta tgtctgtata tcttgtacgg 780
cagctttacat cagccatggt attacagaga ttaaaaaatga aaggtattag aaacctgat 840
cattccagag cactaagtaa agaaaaactt actgcagatc ctgatagtga aattgctaca 900
actagccttc gggatatcct gatgtgccct ttaggaaaaa tgaggctgac aatcccatgc 960
cgtgcagtga cttgtacaca tctgcagtgt tttgatgctg cctctatctt acaaatgaat 1020
gagaaaaagc ccacctggat ttgtctctgt tgtgacaaaa aagctgccta tgaaagtcta 1080
atattagatg ggctttttat ggaaattctc aatgactggt ctgatgtaga tgagatcaaa 1140
ttccaagaag atggttcttg gtgtccaatg agaccgaaga aagaagctat gaaagtatcc 1200
agccaaccgt gtacaaaaat agaaagtcca agcgtcctca gtaagccttg ttcagtgaat 1260
gtgccaagtg agccaagcaa gaagaaagta gatgttattg atcttacaat agaaagctct 1320
tctgacgaag aggaagaccc tctgtccaaa aggaatgca tctttatgtc agaaacacaa 1380
agcagcccaa ccaaaggggt tctcatgtat cagccatctt ctgtaagggt gccagtggtg 1440
acttcggttg atcctgctgc tattccgcct tcattaacag actactcagt accattccac 1500
catacgcaaa gtccaagcat gtcatcagat ttgccaggag aacaaagaag aaatgatatt 1560
aataatgaac tgaagcttgg aacatcttct gatactgtgc aacagtgaat acaaaataaa 1620
accgata 1627

```

<210> 59
<211> 1043
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 179261CB1

```

<400> 59
gtgggatatt tcagtcaaat gaaaatcatc tctgaaaatg tgcccagtta caaaactcat 60
gaatctctta ctttacctcg gagaactcat gacagtgaga agccctatga atacaaggaa 120
tatgagaagg tcttcagttg tgacttagag tttgatgaat atcagaaaat acatactggt 180
ggaaaaaact atgaatgtaa tcaatgttgg aaaacctttg ggatagataa ctccagtatg 240
ttacaactga atattcatac tgggtgtgaaa ccttgtaaat atatggaata tgggaatata 300
tgtagttttt ataaagactt taatgtatac cagaaaaattc ataatgagaa gttctataaa 360
tgtaaggaat acagaaggac ctttgaaaag gttggaaaag ttactccact tcaaagagtt 420
catgatgggt agaaacactt tgaatgctca ttctgtggga aatcctttag agtgcattgca 480
caacttactc gacatcagaa aatccatact gatgagaaaa cttacaaatg tatggaatgt 540
ggcaaggact tcagatttca ttcacagctt accgaacatc agagaattca tactggtgag 600
aaaccctaca aatgtatgca ctgtgagaag gtttttagaa ttagttcaca gtcattgaa 660
catcagagaa ttcacactgg tgagaaacct tatgcatgta aggaatgtgg gaaggctttt 720
ggagtatgta gagaacttgc tgcgtcatcag agaattcata ctggaaaata ctgtggatgg 780
atttaatatg taatcaaggc aattcagtat ctccctctct taaagtctgt ttttagactt 840
catggtcatt ctgtatgtag acgtagaatt gcttagtcat agctgatata tacttatagc 900
tttggttagat gttgccaaat attttctctt tttatgttaa aaagttttt tcatgagttt 960
ctcatcctgg catgttttgt ttacaatagc ttttgatggt tgtattattg ctcttttgac 1020
taatcagttt ttttaattctg gat 1043

```

<210> 60
<211> 2448

<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 259161CB1

<400> 60
ctggcgaggaa gatttttactc ccgagtagcgg gaaagatctg ctcgaggcct ggggtgctttg 60
gtgtcgggaga tccgagagtc ggagatcgga gagtccggaca caggacagtc ggacaccgga 120
cagtcaaaca ccggagagtt agactgggct tctcggtgagg gagaggctct gggataacta 180
ctgttacagc tttgaagggt caagggagga ttggccacca aagcctgttt attagcagct 240
gccattttgtt gaaagaaatt tggattattt tagaaacaaa tttgaaaga aaaagaatgg 300
cgtccgttttc agctctaact gaggaactgg attctataac cagtgaagta catgcagtag 360
aaattcaaat tcaagaactt acggaaaggc aacaagagct tattcagaaa aaaaaagtcc 420
tgacaaagaa aataaagcag tgttttagagg attctgatgc cggggcaagc aatgaatatg 480
attcttccacc tgcgcttgg aataaagaag attttccatg gtctggtaaa gttaaagata 540
ttctgcaaaa tgtctttaaa ctggaagaatg tcagaccact tcagcttgaa actattaacg 600
taacaatggc tggaaaggag gtatttcttg ttatgcctac aggaggtgga aagagcttat 660
gttaccagtt accagcatta tgttcagatg gttttacact cgtcatttgc ccattgatct 720
ctcttatgga agaccaatta atgggttttaa aacaattagg aatttcagca accatgttaa 780
atgcttctag ttctaaggag catgttaaat ggggtccatgc tgaaatggta aataaaaaact 840
ccgagttaaa gctgatttat gtgactccag agaaaattgc aaaaagcaaa atgtttatgt 900
caagactaga gaaagcctat gaagcaagga gatttactcg aattgctgtg gatgaagttc 960
actgctgtag tcagtgggga catgatttca gacctgatta taaggcactt ggtatcttaa 1020
agcggcagtt ccctaacgca tcactaattg ggctgactgc aactgcaaca aatcacgttt 1080
tgacgatggc tcagaaaatt ttgtgcattg aaaagtgttt tacttttaca gcttctttta 1140
ataagccaga tgtgaggttt gttatccatc attcaatgag taaatccatg gaaaattatt 1200
accaagagag tggacgtgca ggtcagatg acatgaaagc agactgtatt ttgtactacg 1260
gctttggaga tatattcaga ataagttcaa tgggtggatg ggaaaatgtg ggacagcaga 1320
agctttatga gatggatca tactgtcaaa acataagcaa atgtcgtcgt gtgttgatgg 1380
ctcaacattt tgatgaagta tggaaactcag aagcatgtaa caaatgtgc gataactgct 1440
gtaaaagacag tgcatttgaa agaaagaaca taacagagta ctgcagagat ctaatcaaga 1500
tcctgaagca ggcagaggaa ctgaatgaaa aactcactcc attgaaactg attgattctt 1560
ggatgggaaa ggtgcaagca aaactgagag tagcaggtgt tgtggctccc acacttcctc 1620
gtgaagatct ggagaagatt attgcacact ttctaataca gcagtatctt aaagaagact 1680
acagttttac agcttatgct accatttctgt atttgaaaat aggacctaaa gctaattctt 1740
tgaacaatga ggcacatgct attactatgc aagtgcacaa gtccacgcag aactctttca 1800
gggtggaact gtctcaaaact tgtcattctg aacaaggtga taaaaagatt ggaggaaaaa 1860
aattccaggc aacttccaga agaaggctgc aaacatgctt cagcaatctg gttctaagaa 1920
tacaggagct aagaaaagaa aaatcgatga tgcctgatat gactgttact aaattttcta 1980
attaaagatg gtttatgcat gtatatgcca ttatttttgt agttagacaa tagtttttaa 2040
aagaatttca tagatatatt atatgtatgg atctatattt tcagagctta tctctgaaga 2100
tctaaaactt tgagaatggt tgaaaattag agatcatgaa ttatataatt ttccagtata 2160
aaacaaggga aaaattttta tgtaaaaccc tttaaatgta aaatatttga gaataagttc 2220
atacaatcgt cttaagtttt ttatgccttt atatacttag ctatattttt tcttttgaca 2280
taactatctt tttgaaagca atattatact gacagaggct cactgagtga tactttaagt 2340
taaatatgta gatcaaggat gtccaatctt ttggcttccc tgagccacac tggaagaaga 2400
attgtcttgg gccgcacata aaatatgcta acactgatga tagctgat 2448

<210> 61
<211> 2255
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 320087CB1

```

<400> 61
ttccgggttct gtacccccat cctttctctc gcccttcta cccgcagctc ctggcgctcg 60
gcggggctaa ctgcagcggg gagatctcgg ccgccaagct ccgcctccc ccccgggctg 120
tgccccgggg ctgcctgag gccgaccacc cgcaaccac ctctagcggc ttgtctcgag 180
gccacacctt tcccaacccc cggcaaaact cagtaggctc gccctcgctg actccccgc 240
cccgctcaa ctgcaagggg cccgcccata gccagtccg gggcggttgc tcacatcgac 300
cggaactccc cgcctccctc cgcggccctt ggggcccgtag gaggccgcag cgaggaggta 360
gagggggcgg gggctgcact aggggtgtccc tagagaacga ggactctgaa ggcgggacat 420
ttgggcgacc cccgggcggg gccagccatt aaacagtccc acttctgtgc cagacactga 480
actgggctct tgacgggcat catctcttaa tcctcagaac atcccaggga gctccacagg 540
atccccatat cctggggccat gaggtaggtt aaagactgcc ccttgagtt ccacgacttc 600
aagtctgtgg atcacctgaa ggtctgtccc cgctacacgg cagtgcctgg acgctctgag 660
gatgatggca tcggcctgac ggagctggac accctgcagc tggagctgga gacctgctg 720
tcttctgcca gccggcgctt gcgtgtgctt gaggccgaaa cccagatcct caccgactgg 780
caggataaga aaggtgacag acgattcctg aagctgggtc gagaccatga acttggagct 840
cccccaaac atgggaagcc caagaagcag aaactggaag ggaaggcagg acatggggccg 900
ggccttgccc caggacggcc caaatccaaa aaccttcagc ccaagatcca ggaatatgaa 960
ttcactgatg accctatcga cgtgccacgg atccccaaaa atgatgcccc caacaggttc 1020
tgggcttcag tggagcccta ctgtgtgac atccaccagg aggaggtccg cacacttgag 1080
gagttactga agccccaga agatgaggct gaggattaca agatcccacc cctggggaag 1140
cactactccc agcgtggggc ccaggaggac ctgctggagg agcagaagga tggggcccg 1200
gcagcggctg tggctgacaa gaagaaaggc ctcatggggc cactgaccga actggacact 1260
aaagatgtgg atgcccgtct gaagaagtct gaggcccagc atgaacagcc ggaagatgga 1320
tgcccccttg gtgccctgac gcagcgctc ctgcaggccc tgggtggagg aaatattatt 1380
tcccctatgg aggattctcc tattcttgac atgtctggga aagaatcagg ggctgacggg 1440
gcaagcacct cccctcgcaa tcagaacaag cccttcagtg tgcgcatac taagtccctg 1500
gagagccgca tcaaggagga gctaattgcc cagggccttt tggagtctga ggaccgccc 1560
gcagaggact ccgaggatga ggtccttgcg gactctgca aacggcaggc tgagctgaag 1620
gcacttagtg cccacaaccg caccaagaag cagcacctgc tgaggctggc aaaggaggag 1680
gtgagccgct gcagctgag gcagcgggtg gcgatggctg acaacgaggt catggacg 1740
tttcgcaaga tcatggctgc ccggcagaag aagcggactc ccaccaagaa agaaaaggac 1800
caggcctgga agactctgaa ggagcgtgag agcatcctga agctgctgga tgggtagccc 1860
tcacccttgc ctgaggtgaa ttatctggcc taggggaggg gaaggagggc ccacttcctt 1920
ctttgggca cccaaacatt ggcctgtggc tgtccctcaa atggcggcag tctctagagg 1980
gccgtggccc ttcctctgag gtcttttggc ctgactctgt acaaccagga cacaggaagc 2040
cctgctgggc tagcctgagg cctagtctct gcttgggtccc cgagatgggg ttggaggggga 2100
cttcgtttct gggctcttct cttccctctt ttaccatccc ccactcccta atcccctacc 2160
cctgtctccc cttcaaggac ttctcccttg tggttttgta aagtgc aaac ttaagaataa 2220
agtgactgct gtggtttttt aaaaaaaaaa aaaaaa 2255

```

```

<210> 62
<211> 2982
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc-feature
<223> Incyte ID No.: 491271CB1

```

```

<400> 62
gcgtggggat gtttaccgcc gtttatccgg gatagagact ccacgtgct gacagcatcc 60
ttttattcac cgctccgaa tttgcaaaga ggaggaagga gggacttctt ggcttctccc 120
agcatagccc cagttatgcc atctcagaac tatgacctc cccagaagaa gcaggagaaa 180
atgaccaagt ttcaggaggc tgtgacattc aaggatgtgg ctgtggtctt ctccaggagg 240
gaactgcgac tgctcgatct taccagagg aagctgtacc gagatgtcat ggtggagaa 300
ttcaagaacc tggttgcagt ggggcattct cccttccaac cagatatggt atcccaattg 360
gaagcagaag aaaagctttg gatgatggaa acagaaaccc aaagaagcag caagcatcaa 420
aataagatgg aaacactcca aaaatttgca ttaaaatacc tttcaaatca agagctgtcc 480
tgctggcaaa tctggaaaca ggttgcaagt gaattaaaca ggtgtcttca ggggaagagt 540
tcccagttat tacaaggatg ctctattcag gtttctgaaa atgagaacaa tataatgaac 600
cctaaaggag atagccctat ttatattgaa aatcaagagt ttccattttg gagaaccacg 660

```



```

cattcttgcg ggaatacata tctgagtgcg tcacagattc agagtagagg taagcaaatt 720
gatgtgaaaa ataacctgca aatacgtgaa gacttcgtga agaaatcacc atttcatgag 780
catattaaaa ctgacacaga accaaaaccc tgcaaaggta atgaatatgg caaaatcatt 840
agtgatgggt ccaaccagaa attaccctta ggagagaaac cccatccatg tggtagtggt 900
ggaagggggt tcagttatag cccaaggctt ccccttcac cgaatgttca cacaggagaa 960
aaatgtctca gtcaaagctc acatctgcga actcatcaga gaattcacc aggagagaaa 1020
ctcaatagat gtcataaacc ttggtgattgc ttcaataaga gctcttttca ttcttatcaa 1080
tctaatacata caggagagaa gtcttataga tgcgacagtt gcggcaaggg attcagtagc 1140
agcacgggtc ttatcattca ttacagaact catactggag agaaacccta taaatgagc 1200
gaatgtggta aatgcttttag tcaaagttca aattttcagt gccatcagag agtccacact 1260
gaagaaaaac catacaaatg cgaagagtgt ggtaagggtc tcggttggag tgttaatctc 1320
cgtgttcacc agaggggtcca caggggtgag aagccctata aatgtgagga atgtggtaag 1380
ggcttcactc aggtctgcaca ttttcacatc catcagagag tccacactgg agagaaaccc 1440
tacaagtgtg atgtgtgtgg taagggttc agccacaatt caccattaat atgccactcg 1500
agagtccaca caggagagaa gccatacaag tgtgaggcgt gtgggaaagg ctttaccctg 1560
aatacagatc tgcataattca tttcagagtt cacacgggag agaaacccta taaatgtaag 1620
gagtgtggta agggcttcag tcaggcttca aatcttcaag tccatcagaa tgtccacact 1680
ggggagaaac gattcaagtgt tgaacgtgtt ggaaggggtc tcagtcagtc ctcaaagctt 1740
caaaccatc agcgaggtcca cactggagag aaaccatata gatgtgatgt gtgtggtaag 1800
gacttcagtt atagttcaaa tcttaacta caccaagtaa ttcacactgg agaaaaacca 1860
tataaatgtg aggaatgtgg gaagggtctt agttggagat caaatcttca tgcacatcaa 1920
agagttcact caggagaaaa accctataaa tgtgagcagt gtgataagag cttcagtcag 1980
gcatagatt ttcggttaca tcagagagtc catactggag agaagccata caaatgttgt 2040
gtctgtggta agggcttcag tcagtcctct ggtcttcaat cccatcagag agtccacacg 2100
ggggaaaagc catacaaatg tgatgtgtgt ggaaagggtc ttagatacag ttcgcagttt 2160
atataccatc agagaggcca cactggagaa aaaccttaca aatgtgaaga gtgtgggaaa 2220
ggctttggta agagcttcaa tcttcgccat catcagaggg tccacacggg agagaaaccc 2280
catatatgtg aggagtgtgg taaggccttc agtctccctt caaatcttct agtccacctg 2340
ggtgttcaca ccagggaaaa actcttttaa tgtgaagagt gtggtaaagg cttcagtcag 2400
agtgcacgtc ttgaagccca tcagagagtc cacactggag aaaaaccata caaatgtgac 2460
atatgtgata agcggttccg tcaccgttca cgtcttacct atcatcagaa agtccatact 2520
ggtaaaaagc tttagaaatg agaaatgtgt taccaacttt tgtctgaatg cacatcttca 2580
agtttttggc tagtccatgc tgggtggtaaa ccctgtaaaa ctactgagag tggagggggg 2640
tttgttcaca cttggaatct ttctaacaaa tccatcaaga tgataacaca gaacctgaa 2700
caggatagata agcgttattt aggggagaaa tagggctggg ggctctcttg gtaagatcta 2760
gttaataata atgatacact ttcatgtgta atatatgcct gaagataatg tgtggaagga 2820
tatttgccat atgctaactg gttttttggc cagggagagt tttgggttat tatccctttt 2880
ctttaatttt cattttatac ttacagtgtt cattattttc ataaaagctg taaagctatg 2940
aaaaatgaat aaaattacta aaaattttct gtaaaaaaaa aa 2982

```

<210> 63
 <211> 1185
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 585172CB1

```

<400> 63
ggagcgtcca gagtccctggc cctgagcggg aatcgagtg gccgaggctg agcggcaggt 60
agaagggggc tctccggggc ttcacaggga acacaggggc ttcggcccaa ccacaagtac 120
gcagttgcac atgccctact tttgcccatt ctttgcaaat cccctaagga acgaacgcgc 180
ctcgcggtgc ggcctttcta atcttcgctt gtcccttcac ttcacagct ggaggtcgaa 240
ttcaccagct cacttagaga aaaggggcgg tgtctcgggt ctcccgtgt ggcatcaagg 300
cgggctccct ataggagctg gtgtggacgt ccgggcgtgg ggttagggcg gatgcgggga 360
tcgcgggggc ggtgttgaca ttgtgtcttc accaggcgga tcgccccgac cctcactcct 420
ggcgtctgag tctctggcgt agccatgctg agtgggcggc tggctcctgg tctgggtctc 480
atggctggcc gcgtttgttt gtgccagggc agcgcgggat ccggggccat cggctccggtg 540
gagggcgcca ttcgcacgaa gttggaggag gccctgagcc ccgaggtgct agagcttcgc 600
aacgagagcg gtggccacgc ggtcccgcct ggcagtgaga ctcacttccg cgtggctgtg 660

```

```

gtgagctctc gtttcgaggg actgagcccc ctacaacgac accgggtggt ccacgcagcg 720
ctggccgagg agctgggagg tccgggtccat gcgctggcca tccaggcacg gacccccgcc 780
cagtgaggag agaactctca gctggacact agccccccat gcctgggtgg gaacaagaaa 840
actctaggaa cccctgaac cccaagagag ggaggaccag gatccgaatg ggctgggtga 900
gcacgaatta cggaggcctt ccccttgata cagtcagga tttgtaaggg atgaagacc 960
ctgggccccca ttctgttggg gtccatacat actctccgaa gatagcaact tgcttcaggt 1020
caaagtgaac ccgagaaaag agaagaatca ctactactg ctcttgccct ggactattca 1080
ggaagggcag cccggatgtt ccatgttaaa tcgtgacaga attgcaccag acctgatgag 1140
ttggaacaaa tcctatacat taaaagaaat tacactaaaa aaaaa 1185

```

```

<210> 64
<211> 1191
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc-feature
<223> Incyte ID No.: 615200CB1

```

```

<400> 64
tggacggacg cgtgggcttg atttctgatt tatgactgct ttttgttgta cccaataggt 60
cgtctaagaa aggtgattat tttgagaggc ctggggagac acacatgctc attctcgagg 120
gtggcggtgg tgcagagggc agagccatgc tcgttcttgc tatcctgaga ttggttgcta 180
tctgtttcct ttgctgctgt gtttttttct gtcagtatta aaggtggaag aaggtccata 240
tctttttctg tgggtgcttc aagtgttggt ggaagtggag gcagcagtga caaggggaag 300
ctttccctgc aggatgtagc tgagctgatt cgggccagag cctgccagag ggtggtggtc 360
atggtggggg cgggcatcag cacaccaggt ggcattccag acttcagatc tccggggagt 420
ggcctgtaca gcaacctcca gcagtaacat ctcccgtaac ccgaggccat ttttgaactc 480
ccattcttct ttcacaaccc caagcccttt ttcacttttg ccaaggagct gtaccctgga 540
aactacaagc ccaacatcac tcaactactt ctccggctgc ttcattgaaa ggggctgctt 600
ctgcccgtct acacgcagaa catcgatggg cttgagagag tgtcgggcat ccctgcctca 660
aagctgggtg aagctcatgg aacctttgcc tctgccacct gcacagtctg ccaaagaccc 720
tcccaggggg aggacattcg gctgacgtg atggcagaca gggttccccg ctgcccggtc 780
tgcaccggcg ttgtgaagcc cgacattgtg ttctttgggg agccgctgcc ccagagggtc 840
ttgctgcatg tggttgattt ccccatggca gatctgctgc tcatccttgg gacctccctg 900
gaggtggagc cttttgccag cttgaccgag gccgtgcgga ctcaagtccc cgactgctca 960
tcaaccggga cttggtgggg cccttggtt ggcattctcg cagcagggac gtggcccagc 1020
tgggggacgt ggttcacggc gtggaaagcc tagtggagct tctgggctgg acagaagaga 1080
tgcgggacct tgtgcagcgg gaaactggga agcttgatgg accagacaaa taggatgatg 1140
gctgccccca cacaataaat ggtaacatag gagacatcca catcccaatt c 1191

```

```

<210> 65
<211> 2596
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc-feature
<223> Incyte ID No.: 997067CB1

```

```

<400> 65
ccggtgtgtg ggtctgtgac aggggtccaac agggcctggt ccgtgtccgg tcccccaaat 60
ctgtcgtccc tgccccagg cattggcacc aacaaaagtc agaattcccg ggaacttgaa 120
cagaggctgc taaattccca gtaattgctc ctttggcctt ctagggactg acttcaaaga 180
aggaaggaaa gaatcaggca gtgcttcctc attctctttt aaaaccgct tcccgctgag 240
tctgcaccca ggagaccaga gagcaccttg ccttccatg gaaactcagg ctgatctcgt 300
atctcaggaa cctcaggccc tgcttgacag tgctcttctc tcaaaagttc tgccttttc 360
cgacaaggac agcctggggg atgagatgtt ggccgctgcg ctccctaaagg ccaagtcca 420
ggagctggta accttgagg atgtagctgt gtacttcac cggaaggagt ggaagcgttt 480
ggaacctgct cagagggacc tctatagaga tgtgatgctg gagaattacg ggaatgtgtt 540

```

```

ctcactggat cgtgagacca ggactgaaaa tgatcaagaa atttctgaag acacaagatc 600
acatgggggtc ctactgggaa gatttcaaaa ggatatttct caggggtctca agtttaaaga 660
agcctatgaa cgagaagtca gtctgaaaag gccgctgggg aactccccctg gagaaagact 720
gaacaggaaa atgccagatt ttgggtcaagt gacagttgag gagaagctaa cccccagggg 780
agagagaagc gagaaatata atgatttttg gaacagcttc actgtgaatt ccaaccttat 840
ctcacatcag agactccccg tgggagacag accccataag tgtgatgaat gtagcaagag 900
ctttaatcga acttcagacc ttattcaaca tcagagaatc cacactgggg aaaagcccta 960
tgaatgtaat gagtgtggga aggccttcag ccagagctca caccttattc agcatcagag 1020
aatccacact ggggaaaaac cttatgaatg tagtgattgt gggaaaacct tcagctgtag 1080
ctctgccctc attctgcac cggaggatcca caccgggggag aaaccttatg aatgtaatga 1140
gtgtgggaag accttcagct ggagctccac cctcaccac catcagagaa tccacactgg 1200
tgagaaacct tagcctgca atgaatgtgg gaaggccttc agcaggagct caaccttat 1260
tcaccatcag agctcacacc tctatcagca ccagagaatc cacactggag agaagcccta 1320
cttcagccag agctcacacc tctatcagca ccagagaatc cacactggag agaagcccta 1380
cgaatgtatg gaatgtggag gaaagtctac ctacagttca ggccttattc agcatcaaag 1440
aatccacacc ggggagaaac cctatgaatg tagtgattgt gggaaagcct tcaggtagag 1500
ctcggtctct gttcgccatc agagaattca cactggagag aagcctttga atgggacgg 1560
catgacaaa agctccctca gagttacgac cgagttaa atcagagagt ccacgtgaaa 1620
gagccacaca cccattttcc tcaactttcc tgagtctcaa gagctcttgc cttaccctat 1680
aaatctcaac agcttaggat gtgtcccttt caactcagac ttttcatttt agagaatggg 1740
gcagatgggg caaatcggtt aattttccca gaaatcacac cagccttaga aagcgtcaag 1800
gcaagtggat ggcgtgctgg gaatagaaag cagctctggg accagttaac ccatttagga 1860
aaggagtgtt cactaaactg tttttctcac agcagaggaa cctttccaag gtggggatgg 1920
aagcacagtc gggacagaat tcgatggatt cctttagttg gagtccgct agtcagcaca 1980
gacagcagtg gaaaggacgc tctgggtcct gttattttgt agggagggt aaggagagact 2040
atttcaaaag tactgttctc agtccagctt taagtttcgg taagaaacat gctgttttgt 2100
ttcatgattt cgttaattat ggaaatttgg cattgagggg ttattttatt gagggtagaa 2160
gagattccag aatcatcatc tgtgatgatg gtgtccttta gggctcttgg agcagccaga 2220
ccatgtttcc aagagaaacc tggatgatatt gccagcagac ccctgccc atccccccagt 2280
tgtcagtgagg ttgatgggg aaatctgtcc aaacagctag taaccggctg tgaggagag 2340
ggtcagaagc acttagcggt ggccctctgat tgcgtgcttc tcttgcttc tteccactcc 2400
aatgatgaaa atgattttct ctaaaagcct gggtaaggat gctttcaagg agctcacttg 2460
gctgtcttgc cctgcccctc cacctctgac acccagcccc aggagccaga ccactcctgc 2520
ctccacctct gactcttcag cagctgaaga ttaatgcaga gaaagagcaa agcccaaaag 2580
ggagaaaaaa aaaaaa 2596

```

<210> 66
<211> 1574
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 1443262CB1

```

<400> 66
cagcccgtag gacgcccgtc gctggacgcg ggtagccgtc tgaggtgccg gagctgcccc 60
aggatggagc cgctgaaggt ggaaaagtcc gcaaccgcc aacaggggaaa cgggctgcgc 120
gccgtgaccc cgctgcgccc cggagagcta ctcttcgct cggatccctt ggcgtacacg 180
gtgtgcaagg ggagtcgtgg cgctgctctg gaccgctgcc ttctcgggaa ggaaaagctg 240
atgcgatgct ctcagtgcgc cgctgcctaa tactgtagt ctaagtgtca gaaaaagct 300
tggccagacc acaagcggga atgcaaatgc cttaaaagct gcaaaccag atatcctcca 360
gactccgttc gacttcttgg cagagtgttc ttcaaactta tggatggagc accttcagaa 420
tcagagaagc ttactctatt ttatgatctg gagtcaaata ttaacaaact gactgaagat 480
aagaaagagg gcctcaggca actcgtaatg acatttcaac atttcatgag agaagaaata 540
caggatgcct ctcagctgcc acctgccttt gacctttttg aagcctttgc aaaagtgatc 600
tgcaactctt tcaccatctg taatgcggag atgcaggag ttggtgttgg cctatatccc 660
agtatctctt tgcctaatca cagctgtgac cccaactgtt cgattgtgtt caatgggccc 720
cacctcttac tgccagcagt ccgagacatc gaggtgggag aggagctcac catctgctac 780
ctggatatgc tgatgaccag tgaggagcgc cggaagcagc tgagggacca gtactgcttt 840
gaatgtgact gtttccgttg ccaaaccag gacaaggatg ctgatatgct aactggtgat 900

```

```

gagcaagtat ggaaggaagt tcaagaatcc ctgaaaaaaa ttgaagaact gaaggcacac 960
tggaagtggg agcaggttct ggccatgtgc caggcgatca taagcagcaa ttctgaacgg 1020
cttcccgcata tcaacatcta ccagctgaag gtgctcgact gcgccatgga tgcctgcata 1080
aacctcggcc tgttgaggga agccttggtc tatggtactc ggaccatgga gccatacagg 1140
atttttttcc caggaagcca tcccgtcaga ggggttcaag tgatgaaagt tggcaaactg 1200
cagctacatc aaggcatgtt tcccgaagca atgaagaatc tgagactggc ttttgatatt 1260
atgagagtga cacatggcag agaacacagc ctgattgaag atttgattct acttttagaa 1320
gaatgcgacg ccaacatcag agcatcctaa gggaacgcag tcagagggaa atacggcggtg 1380
tgtctttgtt gaatgcctta ttgaggtcac acactctatg ctttgtagc tgtgtgaacc 1440
tctcctattg gaaattctgt tccgtgtttg tgtaggtaaa taaaggcaga catggtttgc 1500
aaaccacaag aatcattagt tgtagagaag cacgattata ataaattcaa aacatttggg 1560
tgaaaaaaa aaaa                                     1574

```

```

<210> 67
<211> 2197
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 1521648CB1

```

```

<400> 67
cccgacgaac gcgaggaggg cgaggcgagg ggtgcaggag ggaccctcgc catgggtcca 60
cgggccctaga gtggcggaag ataccggcct ggtgccaaac tggctactgc tgcttctgt 120
ggcctccatg cgtgaggact ggctggactg cccggccctg ggccctggct ggaagcgccg 180
cgaagtcttt cgcaagtcag gggccacctg tggacgctca gacacctatt accagagccc 240
cacaggagac aggatccgaa gcaaagtga gctgactcga tacctgggccc ctgcgtgtga 300
tctcaccctc ttcgacttca aacaaggcat cttgtgctat ccagcccccaggcccatcc 360
cgtggcggtt gccagcaaga agcgaaagaa gccctcaagg ccagccaaga ctcggaacg 420
tcaggttgga cccagagtg gtgaggtcag gaaggaggcc ccgagggatg agaccaaggc 480
tgacactgac acagccccag cttcattccc tgcctctggg tgcgtgtgaga actgtggaat 540
cagcttctca ggggatggca cccaaaggca gcggtcctaaa acgttgtgca aagactgtcg 600
agcacagaga attgccttca accgggaaca gagaatgttt aagcgtgtgg gctgtgggga 660
gtgtgcagcc tgccaggtaa cagaagactg tggggcctgc tccacctgcc tctgcagct 720
gccccatgat gtggcatcgg ggctgttctg caagtgtgaa cggagacgct gcctccggat 780
tgtggaaagg agccgagggt gtggagtatg cgggggctgt cagacccaag aggattgtgg 840
ccattgcctc acttgccttc gccctccccg cccctgtctc aggcgccagt ggaaatgtgt 900
ccagcgacgt tgcctacggg gtaaacatgc ccgcccgaag ggaggctgtg actccaagat 960
ggctgccagg cggcgccccg gagcccagcc actgcctcca ccacccccat cacagtcccc 1020
agagcccaca gagccgcacc ccagagccct ggccccctcg ccacctgcgg agttcatcta 1080
ttactgtgta gacgaggacg agctacagcc ctacacgaac cgccggcaga accgcaagt 1140
cggggcctgt gcagcctgcc tacggcggat ggactgtggc cgtgcgact tctgctgcga 1200
caagcccaaa ttcgggggca gcaaccagaa gcgcagaag tgtcgttggc gccaatgcct 1260
gcagtttgcc atgaagcggc tgcctgccag tctctgtgca gactctgagg atggggcagg 1320
atcgccccca ccttaccgtc gtcgaaagag gccagctct gcccgacggc accatcttgg 1380
ccctaccttg aagccacct tggctacagc cacagcccaa ccagaccata cccaggctcc 1440
aacgaagcag gaagcagggt gtggcttctg gctgccccg cctggcactg acctgtgtgt 1500
tttacgggaa ggcgcaagca gtcctgtgca ggtgccgggc cctgttgcag ctccacaga 1560
agccctgttg caggtgaagc aagagaaggc ggatacccag gacgagtgga caccaggcac 1620
agetgtcctg acttctccc tattgtgccc tggctgccct agcaaggcag tagaccagg 1680
cctgccttct gtgaagcaag agccacctga ccagaggag gacaaggagg agaacaagga 1740
tgattctgcc tccaaattgg ccccagagga agaggcagga ggggctggca caccgtgat 1800
cacggagatt ttcagcctgg gtggaacccg cttccgagat acagcagctc ggttgccaag 1860
gtccaaagac cttaaaaaac ctggagctag aaagcagtag actggaggct tctacagact 1920
gtaggattca agtctgcagg gcaggcactc ggaaggga gatggatgta aagtgtggga 1980
gaccgaggac acagtggagc ccacgagcac gagctggaac ccacgaggat ggcctggaac 2040
ccatgtcagt ctctcaccac ctccagcttc gatgatgtgg tgtcctgca gaagaagctg 2100
gtgccttccc tcacagaggtt aaatatgcat ctggcccagg aattagagaa gctgaaagga 2160
tgatcctggg gaaggtggaa cagctgcagg cctggct 2197

```

<210> 68
 <211> 2081
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1685494CB1

<400> 68
 gcatacgcag tccgtgccta agcggcacga ccatcgagcc aaggtcctcc agcgttccta 60
 gagcggagaa gaaagcgctc cgaagagcta gagctgacac tcggcgatga gctaagacgc 120
 tggttcagag cgtttgggtc ctctgaggcc ccttgaccag gagtgtctct gaagatacag 180
 tccaaagaag gttctccaaa acaaggagag cagtctgaag ctggggatgg caacagcatt 240
 ggtgagtgcc cattccctgg ctcccctgag tctgaagaag gaggggcttc gggtagtgag 300
 ggaggatcac tactctactt gggaaacaggg attcaagctg caaggaaaca gtaaaggcct 360
 tggacaggag ccattgtgca aacaattcag gcagtgcgt tatgaagaga ccacaggacc 420
 tcgagaagag aagatcggtc tccgggagct ctgtcaacag tggctacagc ccgagaccca 480
 taccaaggag cagatcctgg agctgctggt gctggagcag tttctgatca tctgcctaa 540
 ggagctccag gccgggtgctc agggagcatca cccagagagc agggaggacg tggttgttgt 600
 tctggaggat ttgcagctgg atcttggaga aacaggacaa caggtggacc cagaccagcc 660
 aaagaacaa aaaatacttg tggaggagat ggccctctg aaaggagtac aggaacagca 720
 ggttcggcat gagtgtgaag ttacaaagcc tgagaagag aaggttgagg agacaaggat 780
 tgagaatggg aagcttattg tagtaacaga ctcttgtgga agagttagag catctgggaa 840
 aatatctgaa cccatggagg ctcataatga gggctctaac ttggaaggc atcaggccaa 900
 gcccaaagag aagattgagt ataaatgctc agaacgtgag cagagattca tccagcactt 960
 ggacctgatt gaacatgcga gtacacacac gggaaagaaa ctctgcgagt ctgatgtgtg 1020
 tcagagtcc agtcttacag gacataagaa agtccctctc agagagaaag gtcattcagt 1080
 tcatgagtgt gggaaagcct ttcagaggag ttcacacctc gtcagacatc agaaaatcca 1140
 tcttggtag aagccttctc agtgcaatga gtgtggcaaa gtctttagcc agaatgcagg 1200
 ccttttggaa catctcagaa ttcatactgg agagaaacct tatctatgta tccattgttg 1260
 aaaaaatttt aggcgcagct ctacacctaa tcgacatcag agaattcaca gtcaggagga 1320
 gccctgtgag tgcaaggagt gtggaaaaac ctttagtcag gccttactcc tcaccacca 1380
 tcgagaatc catagtcact ccaaagcca tcaatgtaac gagtgtggaa aagctttcag 1440
 tttgacctca gaccttatcc gacaccacag aattcatact ggagaaaaac ctttcaagt 1500
 taacatatgc cagaaagcct tccgactaaa ctacacacct gctcagcatg taagaatcca 1560
 caatgaagaa aaacctatc agtgtagtga atgtggagaa gccttcaggc aaaggctcagg 1620
 tcttttcaa catcagagat atcaccacaa agacaaactg gcttgatgag gtgttctctc 1680
 cttgtagaac atcagagaag gcacattgac tagcaaacag cacttttaga aaagtcaccg 1740
 tagccactg tggcatcaga aaattcttgg gggctgagtt ggaggctccc tgcctctatt 1800
 ctctctcctt tgccttctct gaagtcagct ttggaccaca ataatttcac tgtagatgat 1860
 atgctaggat caaagttaaa cagcattctt cactgcagga catctcagag catgtaacat 1920
 aactgcata ttatatactc taagcaatag agagcttcat gactgagtaa gagttttgaa 1980
 gtcagcagtg aatcaagtgc ccacagattt gcaggcttaa gcagaacaag ggaagattga 2040
 tatttttggg tatgctatag cagctttctc ctatgaaata a 2081

<210> 69
 <211> 2785
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1730829CB1

<400> 69
 gaccgtgcgg cgcccagcgg agtaggggct gcgcttgggg tttgctgaag ctggctgcct 60
 ctcccactcc ccttttgggt gcaaagcgcc gctagcggga agacgggggc cgggcgggga 120
 caggggcacc tgcgtactgg actgagagcc tgcgcccagc ttacatcgac cccaccggc 180
 cccggcccgga cccgacgcga cccgatccga tccgatccca ttccatccgt tctcgtctc 240
 ctcccggtct gaccggttgc ccggccgtgg ttgcgcacac caggcatcca aagctgaggt 300

```

cgctcctacg gcctgggctc gccttcgctt tagagatggt tggcctcttc cctcccaaac 360
agctcatctt caaaacctgg actcttggac tggcacctgg ccacctttcc ctctaccaag 420
actccacttc cgtcttacct acttcttctt cagattcttg gtaccccttg ggttggagac 480
tgctcatttt ccttccaaat taatcccaga cccctaaaaa tattgacaac cttgacaacc 540
ccccaaccca ggagccagac tttgttttgg actaacttcc atagccctat catggaggca 600
gtgtacctgg tagtgaatgg gttgggcttg gtgctggacg tgetgacctt ggtgttggac 660
ctcaacttcc tgctgggtgtc ctccctcctg gcttcccttg cctggctcct ggccttcgtc 720
tacaacctgc cgcacacggt actgactagt cttctgcact tgggcgcgag agtcttgctt 780
tcattgcttg ccttgatcga agccgtggtc cgggttcacat gtgggggctt gcaggccttg 840
tgtactctgc tgtatagctg ctgctctggc ctagagagcc taaagctcct ggggcacctg 900
gcctctcatg gggcactgcg gagcaggagg atactgcacc ggggcgtcct caatgtggtc 960
tccagtggcc atgctttgct ggcgcaggcc tgtgacatct gtgccattgc catgagcctg 1020
gtgctgtatg cctgtcaac cctggctcaac atctgcctca tcggcactca gaacctcttt 1080
tccctggtgc tggccctgtg ggatgcagtg accgggcttc tgtggaggat gacagacgta 1140
gtggctgcct tcctagccca catttcagc agtgctgtgg ccatggccat cctcctttgg 1200
acacctgccc aactagccct ggagctgctg gcctcagctg cccgcctcct ggcagctttt 1260
gtgcttgtca atctcactgg cttggtgttg ctagcttgtg tgetggcagt gacggtgact 1320
gtgttgctac cggactcac cctgaggctg gctaccagag cactcagcca gctccatgcc 1380
cggccatcct accaccgtct tcgagaggat gtcattgcggc tctctcgctt agcactgggc 1440
tcagaggcct ggcgcagagt ctggagccgc agtctgcagc tggcgagtgt gccaaaccgg 1500
ggagggggcac ctggagctcc ccagggtgac cctatgaggg tattctcagt taggaccggg 1560
agacaggaca ctcttcctga agcggggcgc agatcagagg cagaagagga ggaggccagg 1620
accatcagag tgacacctgt cagggggcca gagaggctca atgaggagga gcctccagggt 1680
gggcaagacc cttggaaatt gctgaaggag caagaggagc ggaagaagtg tgtcatctgc 1740
caggaccaga gcaagacagt gttgctcctg ccttgccggc atctgtgctt gtgccaggcc 1800
tgactgaaaa tcctgatcgc ccaccccgtc taccaccgca attgccgctt ctgccggcgg 1860
ggcatcctgc agacctcaa tgtctacctc tgaagcctcc ttccttgcct gccacccctt 1920
ccatgctcca cgcaggcact cacgctagga cagcattaac acctcatctc cgggtccttg 1980
tctgaatccc ctctacccc tgtggccatc ctgccataca tccaggacat tgagttggaa 2040
gacttgatc tgggtggggg caggataaca tggcttctct ttaccagtg ggtcccttcg 2100
atgctgaggg tgggtagtat gtcactatgc aagggccctg agactatttg ctgtgggctc 2160
tcctccagcc tgcccagggc ccaccagat gcctctgggg ttaccctgtt ctgcctcttg 2220
ttttctgtt ggagatctat aggtcctttt cctgcctcct tcacatttcc tccccagctt 2280
ttgcggccac aacacatcag tgtcatttgg gtgttttggc aactcagggg ccttcggatg 2340
atcttaaac tttgtgttca gccagagccc ctgtgccttg gtaggcgttg gggtagtat 2400
ctctcgggtg cctcagagc cacctctgcc tgtgatctgc tgatgaggct ccctcccaac 2460
ctgatccaaa agccagtctc aggagtttac ccttgggatg ggggatgeat ctgcacctga 2520
cttttggggc acgtgcctg tggcacccca agtctcagga gggataaccg 2580
gatttctgct ctttcccttg tcactccac atcacacaga aaaatggcat tctctctgt 2640
ctctccctgg catggagagg gcagactgtg cacatttcac tagggttcaa atacagaagg 2700
cccaggggcc aggggttgca gcttcgtgag gggctctctg ccagtttcc aatgaataaa 2760
gttctcttga cagctaaaaa aaaaaa 2785

```

<210> 70
 <211> 1231
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1864641CB1

```

<400> 70
tcccgtccta tgacgtcagc cgtaaggcgc tgctgtcgta aaaggacgtc cggctccgtct 60
cctagtgtcc ggaatcggct gtcagcctcc ctggctgtta gtaccttctt tcccggagtc 120
ctggctccac agttggattt actgctgtcg cgggtggggc tcacgccatt ccctgtccct 180
cggccccctg agtgagtcgg gtctcccggc gaaagtgagc gaggtttgce cggagcgcgc 240
acgaggggaa aatgcctaaa aaaaagactg gtgcgaggaa gaaggctgag aaccgcccag 300
aacgtgaaaa acaactaaga gcataagaa gcactataga tttagctaaa catccatgta 360
atgcctcaat ggaatgtgac aagtgtcaga ggcggcagaa gaatagagca ttttgctact 420

```

```

tttgtaatc tgtacagaag ttaccaatct gtgcacagtg tgggaaaaca aagtgcata 480
tgaagtcttc agactgtgtc ataaagcatg ctgggtgtata cagtactggc ctgcaatgg 540
tgggtgcaat atgtgacttc tgtgaagcct ggggttgcca tggtaggaaa tgtctcagta 600
cacatgcttg tgccctgccct cttaccgatg ctgagtggtg tgaatgtgaa cgaggcgtgt 660
gggaccatgg aggcagaata ttcagttgtt ctttttgcca taactttctc tgtgaagatg 720
atcaatttga gcatcaagcc agctgccagg ttttagaggc agaaacattt aaatgtgttt 780
catgcaatcg gcttgggtcag cactcatgtc tccgttgtaa ggcttgtttc tgtgatgatc 840
atacaaggag caaagtgttt aagcaagaaa aaggaaaaca gcctccttgt cctaaatgtg 900
ggcatgaaac tcaggagact aaggacctta gcatgtcaac acgctccctg aaatttggca 960
ggcagactgg aggtgaagag ggagatggag cttctgggta tgatgcttat tggagaacc 1020
tttcatctga taagtatggt gataccagct accacgatga ggaggaggat gagtatgaag 1080
cagaggatga tgaagaggaa gaagatgaag ccagaaagga ttcagatact gagtcatcag 1140
atttgtttac taatttgaat ttaggaagga cctatgctag tggctatgct cactatgagg 1200
aacaagagaa ctaggggagc tgctctggtg g 1231

```

```

<210> 71
<211> 700
<212> DNA
<213> Homo sapiens

<220> -
<221> misc-feature
<223> Incyte ID No.: 2444604CB1

```

```

<400> 71
gccccagggtg acacaatggc cgcagtcctat ggccggctggc ttcttccagc ccttcattgtc 60
accgcgcttc ccagggggcc cccggccac cctgcggatg ccgagtcagc ctcccgcatg 120
ctccctggc tcccagcccc tctccctgg cgccatggag cctcccccac gagcccaggg 180
gcatccgagc atgggcccgc caatgcagag ggtgacgcct cctcgtggca tggccagcgt 240
ggggcccccag agctatggag gtggcatgag acccccaccc aactccctcg ccggcccagg 300
cctgcctgcc atgaacatgg gcccaggagt tcgtggcccg tgggcccagc ccagtggaaa 360
ctcgatcccc tactctctct catcccccg cagctacacc ggacccccag gaggagggtg 420
gccccctgga acacccatca tgcctagccc tggagattcc accaactcca gcgaaaacat 480
gtacactatc atgaaccca tcgggcaggg cgccggcagg gctaatttcc cgctcggccc 540
tggcccgag ggcctatggc cgccatgagc gcatggagc ctcaccacgt gaacggatcc 600
ctgggctcgg gcgacatgga cgggttgccg aagagttccc ccggcgccgt ggccggcctg 660
agcaacgccc cggggcaccc cgccggagca cggcgagatg 700

```

```

<210> 72
<211> 2332
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 2445008CB1

```

```

<400> 72
aggcggacgg ggaacgaggc cgtcggccat tttgtgtctg ctctctgtgg gacgtggtgg 60
tagccgttgg gttgggaaag tgagggtatt ttggcctcgt ttctcctgct tcttttctcc 120
tcccttttac tttgccggtg gaacacagtt atgggtcgca agaagaagaa gcagctgaag 180
ccgtggtgct ggtattgtta tagagatttt gatgatgaga agatccttat tcagcaccaa 240
aaagcaaaagc attttaaatg ccataatgt cacaagaaat tgtatacagg acctggctta 300
gctattcatt gcatgcaggt acataaagaa acaatagatg ccgtaccaa tgcaatacct 360
ggaagaacag acatagagtt ggaatatat ggtatggaag gtattccaga aaaagacatg 420
gatgaaagac gacgacttct tgaacagaaa acacaagaaa gtcaaaaaaa gaagcaacaa 480
gatgattctg atgaatatga tgatgacgac tctgcagcct caacttcatt tcagccacag 540
cctgttcaac ctcagcaagg ttatatctct ccaatggcac agccaggact gccaccagta 600
ccaggagcac caggaatgcc tccaggcata cctccattaa tgccagggtg tctcctctctg 660
atgccaggaa tgccaccagt tatgccaggc atgccacctg gattgcatca tcagagaaaa 720

```

```

tacacccagt catttttgcgg tgaanaacata atgatgccaa tgggtggaat gatgccacct 780
ggaccaggaa taccacctct gatgcttggg atgccaccag gtatgcccc accgtgtcca 840
cgtcctggaa ttctctcaat gactcaagca caggctgttt cagcgccagg tattcttaat 900
agaccacctg caccaacagc aactgtacct gccccacagc ctccagttac taagcctctt 960
ttccccagtg ctggacagat ggggacacct gtcacaagct caagtacagc ttcatccaat 1020
tcagaaagtc tgtctgcac ttctaaagct ctgtttccta gcacagcaca agctcaggca 1080
gctgtccaag gacctgttgg tacagatttc aaacctttaa atagtacccc tgcaacaact 1140
acagaacccc caaagcctac attccttgc tatacacagt ctacagcttc aacaactagt 1200
acaacaaata gtactgcagc taaaccagcg gcttcaataa caagtaagcc tgctacactt 1260
acaacaacta gtgcaaccag taagttgatc catccagatg aggtatatac cctggaagag 1320
agaaggcac agttacctaa gtatcaacgt aatcttctc ggccaggaca ggccccatc 1380
ggtaatccac cagttggacc aattggaggt atgatgccac cacagccagg catcccacag 1440
caacaaggaa tgagacccc aatgccacct catggtcagt atggtggtca tcatcaagge 1500
atgccaggat accttctcgg tgctatgccc ccgtatgggc agggaccgcc aatggtgccc 1560
ccttaccagg gtgggcctcc tcgacctccg atgggaatga gacctcctgt aatgtcgca 1620
ggtggccgtt actgatctta ctcatccag tctaataagt ttggagatta aaccttttct 1680
caacttgtgc tgtttatata gccaaagcttc cgtcaataag gcttcattgt gactttaaca 1740
aacattatct tcccacatac caggaaactat tggacattta tttacatgg gaaaaattat 1800
ttggaataat aaagcaggaa cttttcctga agttgcaatt tatactgtat ggcttctttt 1860
tcatgtttca tctaggtttt tagaagtga gtatagtaaa ttggttcgt taaattgtga 1920
aggcgctgga attacatgaa cataccaccc tagtaaaggc aagttctgta agcttacatt 1980
gctattttga aagtttgcct tcacagcatt tcagatgctg ttggacttca tgtcccaac 2040
ctagcttggt gagggctgta actgtttcca agtacttgta cattggaagt ctgaatgtgt 2100
aacaatattt aatgtattta gagttcctca tgttgccagg ttaagaaat ctgaccacc 2160
aaggctatgt gactttcttg tactgttaaa ctctattgta ataaaatgag agaaaaattt 2220
atgctttttt atcataacc cagctgtgga ccactgcctg aaaggtttgt acagatgcat 2280
gccacagtag atgtccacat aataaaattc atagttacca aaaaaaaaaa aa 2332

```

<210> 73
<211> 1936
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature
<223> Incyte ID No.: 2572462CB1

```

<400> 73
ttggttctat attactgttt tatagctgag gagaaagact cacatcattt actaagatca 60
tatagctagc tagtaaatgt ttgagtga aaacaaacaa aggttttctg actttaagag 120
cttgagtttt ttccactata ccataattga tctgttgtaa ttgttaacta atgtgcattt 180
taaaatttct atttgtctta tgtactgagc ccttatacca gtgctaattt atgtgactcc 240
tttctcctgc agctaagaga aaaatacctt ttaattcat ttatagtacc cagtttttaa 300
agaagattta ttttgtaaaa ttttgcttat ggtacatgct atcttagcct gtaaatataa 360
taaagcatta atttttatcc ctccctgggc ttttctcct tctgacttta tacgtctttc 420
tagagagctt atcttctata ataacaattc tttgttttaa agtgagaaag atcagtctaa 480
agaaaaggag aagaaagtga aaaaaacaat tccttctcgg gctacccttt ctgccagcca 540
gctagccagg gccagaaac aaacaccgat ggcttcttcc ccactgccca agatggatgc 600
aatcttaact gaggccatta aggcattgct ccagaagagt ggtgcatcag tggttgctat 660
tcgaaaatac atcatccata agtatcctt tctggagctg gagagaagg gttatctcct 720
taaacaagca ctgaaaagag aattaaatag aggagtcac aaacaggta aaggaaaagg 780
tgcttctgga agttttgttg tggttcagaa atcaagaaaa acacctcaga aatccagaaa 840
cagaaagaat aggagctctg cagtggtatc agaaccacaa gtaaaattgg aggatgtcct 900
cccactggcc tttagtcgac tttgtgaacc taaagaagct tcctacagtc tcatcaggaa 960
atatgtgtct cagtattatc ctaagcttag agtggacatc aggcctcagc tgttgaagaa 1020
cgctctgcag agagcagtag agaggggcca gttagaacag ataactggca aaggtgcttc 1080
ggggacattc cagctgaaga aatcagggga gaaacccctg cttggtggaa gcctgatgga 1140
atatgcaatc ttgtctgcca ttgctgccat gaatgagccg aagacctgct ctaccactgc 1200
tctgaagaag ttgtcctag agaatacccc aggaaccaat tctaactatc aaatgcattt 1260
gctgaaaaaa accctgcaga aatgcgaaaa gaatgggtgg atggaacaga tctctgggaa 1320
aggggtcagt ggcaccttc agctctgttt tcctattat ccagcccag gagttctgtt 1380

```



```

tccgaagaaa gagccagatg attctagaga tgaggatgaa gatgaagatg agtcatcaga 1440
agaagactct gaggatgaag agccgccacc taagagaagg ttgcagaaga aaaccccagc 1500
caagtcccca gggaaggccg catctgtgaa gcagagaggg tccaaacctg cacctaaagt 1560
ctcagctgcc cagcggggga aagctaggcc cttgcctaag aaagcacctc ctaaggccaa 1620
aacgcctgcc aagaagacca gaccctcatc cacagtcac aagaaacctc gtggtggctc 1680
ctcaaagaag cctggcaacca gtgcaagaaa ggaagtaaaa ttgccgggca agggc aaatc 1740
caccatgaag aagtctttca gagtgaaaaa gttaaatttta taggaaaaaa ggggtatcatg 1800
atgaaattca aaatcttatt ttctaaggtc agtgtgcatt tgtttagttt tgatgctttt 1860
caaattacat tattttcttc ccctatgaac attgtgggga gggactctaa ataaaccagt 1920
ttaggcaaaa aaaaaa
1936

```

<210> 74
 <211> 1667
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2572892CB1

```

<400> 74
cgcgaaatcgg cgaccccagt gcctcgacca ctatgccgcg ctctttcttc gtcaggaagc 60
cctccgaccc caatcggaag cctaactaca gcgagctgca ggactctaata ccagagttaa 120
ccttccagca gccctacgac caggcccacc tgctggcagc catcccacct ccggagatcc 180
tcaaccccac cgctcgctg ccaatgctca tctgggactc tgtctctggcg ccccaagccc 240
agccaattgc cttgggctcc cttcggctcc aggagagtcc cagggtggca gagctgacct 300
ccctgtcaga cgaggacagt gggaaaggct cccagccccc cagcccacc tcaccggctc 360
cttgcctctt ctctctact tcagcctctt ccttggaggc cgaggcctat gctgccttcc 420
caggcttggg ccaagtgcgc aagcagctgg cccagctctc tgaggccaag gatctccagg 480
ctcgaaaggc cttcaactgc aaatactgca acaaggaata cctcagcctg ggtgccctca 540
agatgcacat ccgaagccac acgctgcctt gcgtctgcgg aacctgcggg aaggccttct 600
ctaggccctg gctgtacaaa ggccatgtcc ggaccacac tggcgagaag ccttctctct 660
gtccccactg cagccgtgcc ttcgctgacc gctccaaact gcggggccac ctccagaccc 720
actcagatgt caagaagtac cagtgcagg cgtgtgctcg gaccttctc cgaatgtccc 780
tgctccacaa gcaccaagag tccggctgct caggatgtcc ccgctgacct tcgaggctcc 840
ctcttctctt ccataactgc cctgcctga cagccttccc cagctccagc aggaaggacc 900
ccacatcctt ctactgcca tggaaattccc tcttgagtgc cccacttctg gccacatcag 960
cccacagga ctttgatgaa gaccttttc tggttctgtg tctctgctt gggctctgga 1020
agaggccttc ccgtggccat tctgtggag ggagggcagc tggccccag cctgggggga 1080
ttcctgagct ggcctgtctg cgtgggtttt tgtatccaga gctgtttgga tacagctgct 1140
ttgagctaca ggacaaaggc tgacagactc actgggaagc tcccacccca ctcaggggac 1200
cccactccc tcacacacac cccccacaa ggaaccctca ggccaccctc cagaggtgt 1260
gactaactat gcaataatcc acccccaggc cagccccag ggcctgcgga ggcggtggca 1320
gactagagtc tgagatgccc cgagcccagg cagctatttc agcctcctgt ttggtggggt 1380
ggcacctgtt tcccgggcaa ttaacaatg tctgaaaagg gactgtgagt aatggctgtc 1440
acttgcctgg ggcacaagtg ggtgtctctg gctgaccga tgtgtctccc agaactatc 1500
tgggggcccg acaggtgggc ctgggaggaa gatgtttaca tttttaagg tacactggt 1560
tttatatttc aaacattttg tatcaaggaa acgttttcta tagttatatg tacagtttat 1620
tgatattcaa taaagcagtt aatttatata ttaaaaaaaa aaaaaa
1667

```

<210> 75
 <211> 759
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2785674CB1

<400> 75

```

tgggctcgc tccctgggac taggtttcag cggccgctgc gatgaccaa ataaaggcag 60
atccccgacg gcccgaggct caggcggagg cgtgttccgg ggagcgcacc taccaggagc 120
tgctgggtcaa ccagaacccc atcgcgcagc ccctggcttc tcgcccgcctc acgcggaagc 180
tctacaaatg catcaagaaa gcggtgaagc agaagcagat tcggcgcggg gtgaaagagg 240
ttcagaaatt tgtcaacaaa ggagaaaaag ggatcatggg tttggcagga gacacactgc 300
ccattgaggt atactgccat ctcccagtc tgtgtgagga ccgaaatttg ccctatgtct 360
atatccccctc taagacggac ctgggtgcag ccgcaggctc caagcgcgcc acctgtgtga 420
taatgggtcaa gccccatgag gagtaccagg aggttacga tgagtgcctg gaggagggtg 480
agtccttgcc cctacccta tgaggggctc cggtagcacc tgggcacctg ccgtggaag 540
ctattgggct ggcagcagga cgactggctg tctctctgcc caccacact gacggcatct 600
tcccagttcc ccaaggcacg cttcttccc aggcagctct aacagccctt tcatgaagg 660
aatgctagtc ttctgtccat cagtgccatt tctgtagaa ctaaaggctg ttccaagaat 720
gtgggggtggg gaaagtaaat gctaagacta aaaaaaaaa 759

```

<210> 76

<211> 1421

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2797479CB1

<400> 76

```

ccccagtcac tgttctgcc cgttgccgtg cagcctcact gcctttaatc cgcttcagg 60
gctcattcac ctgaatatat tgggtgccat gtcgagatat ggtgaggata ctgcgcaagg 120
catcccggat ccgaaccttc tcaagtagct gttgatagtg ctggagctcc acggtgacat 180
gagctaacag gcgctgatca tcatgggtga tgacatatcg acgcacatag ccccccaga 240
acttagacac aaacatccca gctctgttga tgaagtggcc cagggtgtta agcagctcag 300
aattattctt cagcagcagg tccgtccagg agaaagcact gtctggccc tcaggccgaa 360
tgtacagcag atagaagcgc cagatgtcag cagggatccc cgtgtcctgg gctttgattt 420
tcttggttca gcacgaattc atgaaggaac tcacactcta gagaaacct atgaatgtaa 480
gcaatgtggg aaattgttat ctcatcgctc aagctttcga agacacatga tggcacacac 540
tggagatggc cctcataaat gcacagtatg tgggaaagcc tttgactctc ctagtgtatt 600
tcaaagacat gaaaggactc aactggaga gaaacctat gaatgcaagc aatgtgggaa 660
agccttcctg acttccagtt cccttcgaaa acatgaaaca acacacactg gagagcaacc 720
ctataaatgt aaatgtggaa aagcttttag tgatttattt tctttcaca gtcattgaaac 780
aacacacagt gaagaggagc cttatgaatg taaggagtgt gggaaagcat ttagttcttt 840
taaatacttt tctgcctatg aacggactca cagtgaagaa aaatcttatg agtgtcaaat 900
ttgtggcaaa ctttcagtcg tttcagttac ttaaaaaactc atgaaaggac tcacacggca 960
gagaagccat atgaatgtaa gcaatgcagg aaagcattct tttggccctc tttccttcta 1020
agacatgaaa ggactcacac tggagaaaag ccctatgaat gtaaacactg tggtaaagcc 1080
ttcagtcgtt ccagtttctg tcgagaacat gaaagaactc acgctggaga gaagccctat 1140
gaatgtaagg aatgtgggaa agccttcagt tctctcagtt cctttaatag acataaaagg 1200
acacactgga aggatattct ataagtgtat ggaatgtggg aaagcattca ttggttttat 1260
cacattcaga tacttgaaag aaataaatcc ttgtaatgta aacgtggtaa agccttaaga 1320
agtttcagg ctgggcgcag cggctcacac ctgtaatccc agcactttga gaggccgagg 1380
agggcatatc acgaggccag gagatcgaga ccagcctggg g 1421

```

<210> 77

<211> 2386

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2960640CB1

<400> 77

```

gacttttaag ggatcacaga gctcacacca aagaccaggg gaacagtcag aagcctggct 60

```

```

tgctctcag gctcccagga acctgcctca aaacacaggt ctccacgacc aggagacagg 120
tgctgtggtc tggacagctg ggtcccaggg accagccatg cgtgacaaca gagctgtatc 180
cctctgtcag caagaatgga tgtgcccagg ccctgcacaa agggccctct acaggggtgc 240
caccagagg aaggacagtc acgtctcgtt ggcaacaggt gtgccctggg gctatgaaga 300
gaccaagacg ctccctggcta ttcttagtag ttctcaattt tatggaaaac tccagacctg 360
tcagcagaac agccagatct acagggccat ggcggaagga ctctgggagc aggggtttct 420
gcggaaccca gaacagtgtc gcaccaagtt caaaagccta cagttgagtt accgcaaagt 480
gaggagaggc cgtgtgcctg agccttgtat cttttatgag gaaatgaatg ctctttcagg 540
ctcctgggccc tctgcacctc ctatggcaag cgatgctgtt cctggccaag aaggaaagtga 600
tattgaggct ggagagctga atcaccagaa tggggaaccc acggaggtag aagatggcac 660
tgtggatggt gcagacaggg atgaaaagga cttcaggaat cctggccagg aagtcaggaa 720
actagacctg ccagtgtctg tcccaaacag acttggtttt gaggttcaaga acgagattaa 780
aaaagaaaaat ctaaaatggg atgattcaga ggaagtagaa ataaacaagg ctttacagag 840
aaagtccaga ggagtttatt ggcactctga gctacaaaaa ggcttgagga gtgagccaac 900
atcaagaagg caatgtagaa attctccagg ggagagttag gagaaaaccc catcccagga 960
gaagatgagt caccagagtt tttgtgccag ggacaaagcc tgtacacata tcctctgtgg 1020
gaaaaaactgc tctcagagtg tgcaactctc ccacaagcca gcgctcaaac tggaaaaagt 1080
atctcaatgt cctgaatgtg ggaaaacctt tagcgaagt tcttatcttg ttcggcatca 1140
aagaatccac acaggcgaga agcctcacia gtgcagttag tgcgggaagg gctttagtga 1200
gcgctccaac ctactgtccc acctacgaac tcacacaggg gagaggccct atcagtgtgg 1260
gcaatgtggg aaaagcttca accagagttc cagcctcatt gtccaccaga ggaccatac 1320
cggggaaaaag ccttaccagt gcattgtctg tggaaagaga ttcaacaaca gttcccagtt 1380
cagtgtcac cgcgcatcc acactgggga gagcccatc aagtgtgcag tgtgtgggaa 1440
aatcttcaac aatagctccc acttcagtgc ccaccgaaaa acccacactg gtgaaaagcc 1500
ttacaggtgt tctcactgtg agagaggctt cactaagaac tctgccctca cccgtcatca 1560
gacagtcacac atgaaagcag tactctcacc acaggaagga agagatgcgt tatgagtgtg 1620
tcggtaaaact gtcagattaa gttcctcagg tcagcatgta tgagctttct tctgctgtgg 1680
agagatctag ccagtccctg actttgcaac agacctactg actactggat cttaagacct 1740
atgtctagga ccaggagtca gcataaggac gctgacctct cctggctgtg cctgtgactc 1800
cagagtctcta tcttactgtg acttaaagtt tgatggagag aaagctgtag gatccataaa 1860
ttctaccagg aatccagggt ctctctgttc cttagcactga gaatgggcac ccagtggtec 1920
aagaacactt tctgggctaa catagtcttc acacagggct gagaaagaaa gtgtctcctt 1980
tcttgaaag cacatgtaaa agttaagggc ctgaatctct tctaaaccaa taattgacct 2040
ctaggcaccc accgtaatac tgctaccttc agcagagaag acactgctct ttgtaaccca 2100
gccaccacca aaaagcaaac agaaagaagg aaggatggtt aagccattgg ataatactga 2160
atctgtttcc ctaagtgact taaccttaga gcaagcacia catccaggtt aattaattgt 2220
aagattttct ctctcattat tgccctcacc atagtctctg attgtctctt aaagtaagtg 2280
gtttatagac attactattt ctgataataa tttacaaact actataaaca aatttataaa 2340
cattactaat ttctgatgaa aataaagttg tttctccctc caccac 2386

```

<210> 78

<211> 1432

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 3454051CB1

<400> 78

```

agtatgagct ggacctgtcc gcgttgccag caacctgttt tcttcgctga gaaggtgagc 60
tccttgggca agaactggca ccgcttctgc ctgaaatgtg agcgctgcca cagcatcctg 120
tcccctggcg ggcattgcaga gcacaatggg aggcatact gccacaagcc atgctatggg 180
gctctctttg gacccagggg cctcccccac atgaagacat tcaactggga gacctcgtg 240
tgccctggct gtggggagcc cgtctatttt gctgagaagg tgatgtcatt aggcagaaat 300
tggcaccgac cgtgtcttag gtgccagcgt tgccacaaga ccctgactgc tgggagtcac 360
gctgagcatg atggagtccc ctactgccac gtccctgtct acggctacct gtttggcccc 420
aaaggtgtga atctggcga tgtgggctgc tacatctatg acccagtga gataaaattc 480
aaatgagacg ctacacaaaa aggtcacctc aactcagcc tcccatcatg cccctcatgg 540
tccaatggaa gctacaaaaa tctccagtc catgggggtt ggggaagggt ggatcttggg 600
ggcctgggccc taggctccat ggtaggccag agagtctaga ctttctctgc caattttttc 660

```

```

cctttcccat ttctatctgg ttagggaaca gcccgttttg aagggtatcc tcctcctggc 720
catcacaacc cttttcccca gcacattctg gagcttcaag gtactcataa aacttggtgt 780
tattgaattt cagcctctgt ggcttcttca ataaaatgtt ggctcccatg ctttcaactc 840
ttctttgggc atgagaccag tgggttggag gatggggagt gtgggggttg ggatgacatg 900
cattgccttg caggggtgctt cggaggttagc agggccagcc atgagaacaa aaagctctgt 960
tccttttgtc ccttggggcct ggcatgtgca gtccctagcac cacacagtgg acagcatgcc 1020
caccagcccc attggtacca gaagtctcat atgctagtcc tttcttttagc accatctcta 1080
gaagaagcag aagcacctta ttcagtaact cttttgagca tggcaacaga tcctatggta 1140
gggcctccca agagccttca ttatccattt gggagatgaa cagactgagg cccagagagg 1200
gaaagccaca tgcccaaggt cacacaacaa gttaatgggt aagggtttat cagagcccag 1260
ggcagactca gtggcttctt atccagggct cttctcacag ctctgcacca ctgccccaac 1320
ccaaggggca cctttattta cagaatctcc ccaaccgtga gacgggtgcc agcagaccac 1380
tgcattctgg gaagtcaact gttcctagaa gcaaataatc aaggatggat ga 1432

```

<210> 79

<211> 1816

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 3510640CB1

<400> 79

```

cactgatgca ggaactgtat agcacaccag cctccaggct ggactccttc gtggctcagt 60
ggctgcagcc ccaccgggag tggaagggaag aggtgctaga cgctgtgcgg accgtggagg 120
agtttctgag gcaggagcat ttccagggga agcgtgggct ggaccaggat gtgcgggtgc 180
tgaaggtagt caaggtgggc tccttcggga atggcacggt tctcaggagc accagagagg 240
tggagctggt ggcgtttctg agctgtttcc acagcttcca ggaggcagcc aagcatcaca 300
aagatgttct gaggcgtata tggaaaacca tgtggcaaaag ccaggacctg ctggacctcg 360
ggctcgagga cctgaggatg gagcagagag tccccgatgc tctcgtcttc accatccaga 420
ccagggggac tgcggagccc atcacggtca ccattgtgcc tgcctacaga gccctggggc 480
cttctcttcc caactcccag ccaccccctg aggtctatgt gagectgatc aaggcctgcg 540
gtggtcctgg aaatttctgc ccattcttca gcagactgca gagaaatttc gtgaaacatc 600
ggccaactaa gctgaagagc ctctctgcgc tgggtgaaaca ctggtaccag cagtatgtga 660
aagccaggtc ccccagagcc aatctgcccc ctctctatgc tcttgaactt ctaaccatct 720
atgcctggga aatgggtact gaagaagacg agaatttcat gttggacgaa ggcttcacca 780
ctgtgatgga cctgctcctg gagtatgaag tcactctgtat ctactggacc aagtactaca 840
cactccacaa tgcaatcatt gaggattgtg tcagaaaaca gctcaaaaaa gagaggccca 900
tcactcctga tccggccgac cccaccctca acgtggcaga aggttacaga tgggacatcg 960
ttgctcagag ggctctccag tgcttgaac aggactgttg ctatgacaac agggagaacc 1020
ccatctccag ctggaacgtg aagagggcac gagacatcca cttgacagtg gagcagaggg 1080
gttaccacga tttcaacctc atcgtgaacc cttatgagcc cataaggaag gttaaagaga 1140
aaatccggag gaccaggggc tactctggcc tgcagcgtct gtccctccag gttcctggca 1200
gtgagaggca gcttctcagc agcaggtgct ccttagccaa atatggggtc ttctcccaca 1260
ctcacatcta tctgtggag accatccctt ccgagatcca ggtcttctgt aagaatcctg 1320
atgggtgggag ctagccttat gccatcaacc ccaacagctt catcctgggt ctgaagcagc 1380
agattgaaga ccagcagggg cttcctaaaa agcagcagca gctggaattc caaggccaag 1440
tcctgcagga ctggttgggt ctgggggatct atggcatcca agacagtgc actctcatcc 1500
tctcgaagaa gaaaggagag gctctgtttc cagccagtta gttttctctg ggagacttct 1560
ctgtacattt ctgccatgta ctccagaact catcctgtca atcactctgt cccattgtct 1620
actgggaagg tcccaggtct tcaccagttt tacaatgagt tatcccaggc cagacgtggg 1680
agctcacacc tgtaatccca gaactttggg aggccagagt gggaggagcg cttgagccga 1740
ggagttcaag accagcctgg gtatcatagg gagacccgt ctctacaaaa taaaaaata 1800
attcactggg aaaaaa 1816

```

<210> 80

<211> 1556

<212> DNA

<213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3815083CB1

<400> 80
 ctccaggtccg gagcgcggtc gggacacagc gcctctagga gaaagcctgg aaggcgctcc 60
 ggggggtatcc agagctctta gcgggccggc agcatgtgcg gggccccagt aaatggaaat 120
 gtttttctaac atataaaaaac ctacagaaga agaaaataat tttctggatc aaattagaag 180
 tctgtattat attgatgtct ccagattcaa atatattaga aagcagccgt ggagacaacc 240
 atcttcattt tgggagaaat aactaaagcc cgctcaagc attagaacta cagacaaacc 300
 ctgatgagac ctctccagat tgtcccaagt cgattgattt ccagctata ttgtggcctg 360
 aagcctccag cgtccacacg aaaccagatt tgctgaaaa tggctcggcc aagttcaagt 420
 atggcagatt ttcgaaagtt ttttgcaaaa gcaaagcaca tagtcatcat ctcaggagct 480
 ggtgttagtg cagaaagtgg tgttccgacc ttcagaggag ctggagggtta ttggagaaaa 540
 tggcaagccc aggcctggc gactccccctg gcctttgccc acaaccctgc cggggtgtgg 600
 gagttctacc actaccggcg ggaggtcatg gggagcaagg agcccaacgc cgggcaccgc 660
 gccatagccg agtgtgagac cggctggggc aagcagggcc ggcgagtcgt ggtcatcacc 720
 cagaacatcg atgagctgca ccgcaaggct ggcaccaaga accttctgga gatccatggg 780
 agcttattta aaactcgatg tacctcttgt ggagttgtgg ctgagaatta caagagtcca 840
 atttgtccag ctttatcagg aaaaggtgct ccagaacctg gaactcaaga tgccagcatc 900
 ccagttgaga aacttccccg gtgtgaagag gcaggctgcg ggggcttgcg gcgacctcac 960
 gtctgttggt ttggagaaaa cctggatcct gccattctgg aggaggttga cagagagctc 1020
 gccactgtg atttatgtct agtgggtggc acttctctctg tgggtgtacc agcagccatg 1080
 tttgcccccc aggtggctgc caggggcgtg ccagtggtg aatttaacac ggagaccacc 1140
 ccagctacga acagattcag gtttcatttc cagggaccct gtggaacgac tcttcctgaa 1200
 gcccttgccg gtcattgaaa tgaaactgtt tcttaagtgt cctggggaag aaagaaatta 1260
 cagtatatct aagaactagg ccacacgcag aggagaaatg gtcttatggg tggtagctg 1320
 agtactgaac aatctaaaaa tagcctctga ttcctctgct ggaatccaac ctgttgataa 1380
 gtgatggggg tttagaagta gcaaagagca ccacattca aaagtacag aactggaaag 1440
 ttaattcata ttatttggtt tgaactgaaa cctgaggtat ctttgatgtg tatggttggt 1500
 tattgggagg gaaaaatttt gtaaattaga ttgtctaaaa aaaataaaaa aaaaaa 1556

<210> 81
 <211> 1951
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3988457CB1

<400> 81
 ctccgattat gggataggag aagtgccgt ggagcccctg gatgtcccct taccctccac 60
 gatcaggcca gcttcccccg tggccgggtc tccaaagcag ccggtgctgt gctactaccg 120
 tggcgctgtc ggtggcacgt ttgaccgcct gcacaacgcc cacaagggtg tgctcagtgt 180
 cgcgtgcatc ctggcccagg agcagcttgt ggtgggagta gcagacaaag atctgttgaa 240
 gagcaagttg ctccctgagc tgctccaacc ttatacagaa cgtgtggaac atctgagtga 300
 attcctggtg gacatcaagc cctccttgac ttttgatgtc atccccctgc tggaccctta 360
 tgggcccgtc ggcctctgacc cctccctgga gttcctggtg gtcagcgagg agacctatcg 420
 tgggggggatg gccatcaacc gcttccgctc tgagaatgac ctggaggaaac ttgctttgta 480
 ccagatccag ctgctgaagg acctcagaca tacagagaat gaagaggaca aagtcagctc 540
 ctccagcttc cgcagcgaa tgttggggaa cctgcttcgg cctccatatt aaaggccaga 600
 gctccccaca tgtctctatg taattgggct gactggcctc agtggctctg ggaagagctc 660
 aatagctcag gcactgaagg gcctgggggc gtttgcatt gacagtgaac acctgggtca 720
 tggggcctat gccccagtg gccctgccta ccagcctgtg gtggaggcct ttggaacaga 780
 tattctccat aaagatggca tcatcaacag gaaggtccta ggcagccggg tggttgggaa 840
 taagaagcag ctgaagatgc tcacggacat tatgtggcca attatcgcaa agctggcccc 900
 agaggagatg gactgggctg tggctgaggg aaagcgtgtg tgtgtgattg atgcccgtgt 960
 gttgcttgaa gcgggctggc agaacctggg ccatgaggtg tggactgctg tcatccaga 1020
 gactgaggct gtaagacgca ttgtggagag ggaatggcctc agtgaagccg cggctcaaag 1080
 cgggctgcag agccagatga gcgggcagca gcttgtggaa cagagccacg tgggtgctcag 1140

```

cagcccttgt gggagccgca tatcacccaa cgccagggtgg agaaagcctg ggcctcttgg 1200
cagaagcgca ttcccaagac tcatcaggcc ctcgactgaa aagttctcag tggggccaga 1260
ctggctcctg gagctgacaa ggcaccccggt ggtgaggaga aatggggggc ttgatgctca 1320
ccctgggttca ggcccagagg tccaagctat actgtgcagg acatggccag gcctgggtgga 1380
cacaggaagc ctacccaaca cgctggtatt tggccaacac tgaggatgtg gttcatgggg 1440
gagcagtcct ctccccactc ttgcccattg gtgactctta cccacagctg actagggcca 1500
gcgcaaatac tggaaacctgt aacagaatta aagggtgaat ttctgaaaaa aaaatagaat 1560
tttgacatc tacaactaac tcgatttaca cttacgaaca taatggactc ttaaaaaatg 1620
gaaagggata acagggaccc cccgggttct gcatccttcc tcccgggat tttttccgg 1680
ccgggtcctt gcggtgaac tgattttct tacactgcgc ctatttaaac gttggggtaa 1740
ccagggtcgg accttttccc ttggaacttt tttaccctg cacgatttcc actgcaactt 1800
agactcggga aacttagggg ggaatccctg gggggcctaa ggcggaact cctcccttt 1860
atctgcttgg ggtcaaaagag cccgtttctc catctggcaa ctctgtccat ccaaggggtt 1920
ttcgggttct cgggccaagg cccgggttgg g 1951

```

<210> 82
 <211> 1313
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 131890CB1

```

<400> 82
cgctcgggagg gcctaagtcc gtgtgcggtg cccttcggcc ggcttgagcc ccagagtcag 60
ctcccccttcc tcgcccagcg cccccaggcc gctcccgggg ctcacggaat agtaaagaaa 120
cacatcataa aacctcccag gacataaagg tgagcacaga ccctgttttg atcaagtcag 180
ttcctggagc ctgaatgatg actgctgaat cacgggaagc cacgggtctg tccccacagg 240
ctgcacagga gaaggatggt atcgttatag tgaagggtgga agaggaagat gaggaagacc 300
acatgttggg gcaggattcc acctacagg acacgcctcc tccagacca gagatattcc 360
gccaaagcct caggcgcttc tgttaccaga acacttttgg gccccgagag gctctcagtc 420
ggctgaagga accttgtcat cagtggctgc ggccagaaat aaacaccaag gaacagatcc 480
tggagcttct ggtgcttagag cagtttcttt ccactctgcc caaggagctc caggctctggc 540
tgaggaata cgcctccgat agtggagagg aggccgtgac ccttctagaa gacttgagc 600
ttgatttatc aggacaacag gtcccaggtc aagttcatgg acctgagatg ctcgcaaggg 660
ggatggtgcc tctggatcca gttcaggagt cctcgagctt tgaccttcat cacgaggcca 720
cccagtccta cttcaaacat tcgtctcgga aacccgcct cttacagtca cgaggtaaga 780
agcaaggttt catttagggg aagggaatg attcaggacg agagtctttg tgctgctgag 840
tgcctgtgat gaagaagcat gttagtcttg ggcaacgtag cgagacccca tctctacaaa 900
aaatagaaaa attagccagg tatagtggcg cacacctgtg attccagcta cgcaggaggc 960
tgagggtggg ggaattgctt agcccaggag gttgaggctg cagtgaagtc taatcatgcc 1020
actactccaa cctgggcaac acagcaagga cctgtctca aaagctactt acagaaaaga 1080
attaggtctg gcacggtagc tcacacctgt aatcccagca ctttgggagg ctgaggcggg 1140
cagatcactt gaggtcagga gtttgagacc agcctggcca acatggtgaa acctgtctc 1200
tactaaaaat atgaaaatta gccaggcatg gtggcacatt cctgtaatcc cagctactcg 1260
ggaggctgag gcaggagaat cacttgaacc caggaggtgg aggttgagtc aag 1313

```

<210> 83
 <211> 1197
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 238642CB1

```

<400> 83
cggtctgagc gtgaaccgaa gtccttatat tcccgggctt ctttctcttc tgggtaccag 60
ctccttactg ccttcgagac aagcgtgccg tgcgtgcttg tggccaaggg aaggaagagc 120
tggttgatcc acagatagct ccttctctcc cgccccttcc tttttgtttg gaggtcccag 180

```

```

gatctgtgtt cacagacatc tgggggaaga aaaggagcag gaaactaccc cgcacagagt 240
taagcaggaa acaacaacaa catcatgcaa aaaccctgca aagaaaacga aggaaagcca 300
aagtgcagcg tgccaaagag ggaggaaaaa cgcccgtatg gagaatttga acgccagcaa 360
acagaaggga attttagaca gaggctgctt cagtctctcg aagaatttaa agaggacata 420
gactataggc attttaaaga tgaagaaatg acaaggaggg gagatgagat ggaaagggtg 480
ttggaagaga taaggggtct gagaaagaaa tttagggtct tgcattctaa ccataggcat 540
tctcgggacc gtctttatcc catttaatta atttctctga caattcaatt atttctgtt 600
attaatgttg ccactgcttt ctgtttgtct gcactttctt gataaatatt tgctatcgtt 660
ttactccagt cattcgatgt tgctgagatt tacatatgac tcttgtcaac atctcatctt 720
ttgaccaaat cttattcatt taataagagg tctcattcat ttgcatggaa aaatgctcat 780
tgtatattgc aaagtgaaaa taacgagttg caaacacagt tatacatata tgttgtata 840
tatgtacact ttatttgtac atttctatgt gacataatgc aaaggaaagt gtctgatttt 900
attatcacac aaaggtaaac agtgaatctc cttttttttt ctttttgctt 960
atctgcatct tctcacttgc caaaaaatga atatatgttt atgtgtgtat attactgtg 1020
tcacaaaaaa ccctaaagta gacagtaaaa gaacttgtca atcgctttg gaaggcaatg 1080
aaacacttaa taaactctca ataacagaag cgtaaaaatg aaatgtaaac ctccaattac 1140
ctctggatct cttagccaga gtaataaact ggtaattatt acagataaaa aaaaaaa 1197

```

```

<210> 84
<211> 2170
<212> DNA
<213> Homo sapiens

```

```

<220>
<221> misc-feature
<223> Incyte ID No.: 669862CB1

```

```

<400> 84
agcgtgcgcg ataggacagg gccttaaatt catggttatg tgtttggttt tcacttetaag 60
ataaagccga accactgaca aaattatata gagagcaatg tgataatgtt aggaatagag 120
atattcagga cactactcctg tgcattctga tccactcagt ctgagtggat atctaaaaga 180
tagtgtatgt aaaatgctcc tatggtaatt ctgatttgca actaaatatg agaactagtg 240
actttgtctt tgcctactca taaatcttac atcctcctgt gcactttcat aattttctgt 300
gcccttcttg ctttttataat attgcttggc agtactttag aattttctaa actctttgat 360
tactaacaga tacacttttt agttgatcac tatttttaat atagatgtcc tctccctacc 420
cgctattatt ggaaaatagc atttgtttgt ttttccattt tcttccagac ttttaattca 480
caactgaaag caacaagtta tcttcagaaa aaagaaatta tgaagttaa gcgtaccatc 540
aggagacatg gaaaagaaat aaaaccttca accttatgag gtttattttc agaactgacc 600
cacagtacac aattgaaatt gggagacaac agagacctaa agtgggatgt ttagtcaaaa 660
tgatattcaa aaaacataaa tcccttcctc tacataagag aaataacaca agagagaaat 720
catatgagtg taaggaaatc aagaagggtc ttgagaaaata tttgcacctt actgaacatc 780
tgagagacca tactggtgtg ataccctatg aatgtaatga atgtggaaaa gcatttgtag 840
ttttccagca ttttattaga catcgaaaaa tccacactga tttgaaacc tatgaatgca 900
atggatgtga gaaggccttt aggttttatt cacagcttat tcagcatcag ataattcata 960
ctggtatgaa accctatgaa tgaagcaat gcgggaaggc ttttagacgt cattctcacc 1020
ttacagaaca tcagaaaatt catgttggct tgaaccctt tgaatgtaag gaatgtgggg 1080
aaacgtttag atttatatga catatgtgtc tgcatacaga aattcatcat ggtgtgaaac 1140
cctacaaatg taaagaatgt ggaaaggctt ttggtcatcg ttcaagtctt taccaacata 1200
agaaaattca ttctggtgag aaaccatata aatgtgaaca atgtgaaaag gcctttgttc 1260
gcagctatct acttgttgaa catcaaagaa gtcatactgg tgagaaacct catgaatgca 1320
tggaatgtgg aaaggctttt agtaagggtc caagccttct taaacataag agaattcata 1380
gtagttagaa actctatgat tgtaaggatt gtggaaaggc cttttgtaga ggctctcaac 1440
ttacacagca tcagagaatt cactactggtg agaagccaca tgaatgtaaa gaatgtggga 1500
agacttttaa tcttacttca tatcttattc aacatcagat aattcactat gatttgaagc 1560
catatgaatg taagcaatgt gggaaagcct tcagtcgtgt tggagacctt aagacacatc 1620
aatcaattca tgctggggag aaaccctatg aatgtaagga atgtggaaaa accttttagac 1680
ttaattctca actaatttat catcagacaa ttcatactgg tttgaaacc tatgtatgta 1740
aagatgtaaa gaaggccttt cgctctatct caggctcttc tcaacataag agaattcata 1800
ctggtgaaaa accctatgaa tgtaagaaat gtgataaggc ctttaatcgc agtgatcgac 1860
ttactcaaca tgagacaatt cactactggtg tgaaccaca gaaatgcaaa gaatgtggta 1920
aggcctttag tcattgctat caacttagtc aacatcaaag atttcacat ggtgagagac 1980

```

```

tcttaatgta atgagagggga aagcctttag ccatggcaca tttttactgt tgtcactatt 2040
atgatgctat agtgaagatt aaactagtta atatagaata taaataacttg gaaaggcatc 2100
tggcacatcg tatttgctta ctaaatacat tatttttatg ataattgtta gaattactaa 2160
gaataaatga                                     2170

```

<210> 85

<211> 1904

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 1003663CB1

<400> 85

```

cagctctcag gtccgacacc cgctggaagc cggcgcgggc gcaggcgcgc acgcaaaggc 60
ggccgggagt aaggcgggact gaaggaggag ctgatggaa gctgcgaga aggggcgtaa 120
ctgatttgga aaccagagga aaggcgctgt ttccaccgaa ttagaatcgc gggaaaatag 180
agaagagttt gtttgaaggt ctgcgcgat cgagaccgga agtccttcat ctcaagcatc 240
caatgctgaa agcggcctga ttttctctac cggaagccct tttccagagg ctgggaacac 300
ggcccaccta gcaggaagtc ccacctcctt gagctccgcc acccttcccg aagtttttct 360
gtcacctgtg ttaggtctcg tcccccttcc gcgttttatc cccgtaccag aaaaggatac 420
atttagtgcc tcccaccag ctccactaaa cgggttggat atctcattct ttgagttggt 480
gttcttcccc cggcgcctccc atgtagctgg gaagtgggac ctgggggttg ttggaccctt 540
gggatccctaa agggggggca gggagggcgc agaactccgc ttctgctcct tgctaccagg 600
acgcgcggcc tcttcagcct ctttctccc gctgccatgc accctgcagc cttcccgctt 660
cctgtggttg tggcgcgtgt gctgtgggga gcggcccccga cccgggggct cattcgagcg 720
acctcgggacc acaatgccag catggacttt gcagaccttc cagctctgtt tggggctacc 780
ttgagccagg agggcctcca ggggttccct gtggaggctc accagacaa tgcctgcagc 840
cccattgccc caccaccccc agccccggtc aatgggtcag tctttattgc gctgcttcca 900
agattcgact gcaactttga cctcaaggtc ctaaagtccc agaaggctgg atatggtgcc 960
gctgtagtac acaatgtgaa ttccaatgaa cttctgaaca tgggtgtggaa tagtgaggaa 1020
atccagcaga agatctggat cccgtctgta tttattgggg agagaagctc cgagtacctg 1080
cgtgccctct ttgtctacga gaagggggct cgggtgcttc tggttccaga caataccttc 1140
cccttgggct attacctcat ccccttcaca gggattgtgg gactgctggt tttggccatg 1200
ggagcagtaa tgatagctcg ttgtatccag caccggaaac ggctccagcg gaatcgactt 1260
accaaaagag aactgaaaca gattcctaca catgactatc agaagggaga ccagtatgat 1320
gtctgtgcca tttgcctgga tgaatatgag gatggggaca agctgcgggt actccctgt 1380
gctcatgcct accacagccg ctgcgtggac ccctggctca ctcagaccgg gaagacctgc 1440
cccatttgca agcagcctgt tcatcggggt cctggggacg aagaccaaga ggaagaaact 1500
caagggcaag agggagggtga tgaaggggag ccaagggacc accctgcctc agaaaggacc 1560
ccacttttgg gttctagccc cactcttccc accctcttgg gttccttagc cccagctccc 1620
cttggttttc ctgggccttc aacagatccc ccactgtccc ctccctcttc cctgtttatc 1680
ctgggtcta at aacccccac acatacacct ctggtgacct atttgacag accgtcgtct 1740
tccctccagt cttctgaggg ataggggaca ttccatccca agcttctccc ttaccacac 1800
ctatcctttt gaggggcttt ggggtggggc agagggactg ggtcttcaact 1860
tcttgggcta ataaaattgt ttctttgtgg actaaaaaaa aaaa 1904

```

<210> 86

<211> 1249

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 1432557CB1

<400> 86

```

cgtttatcct gttgggggag gaagtggag agcccttatt cgtattggct tagatttgca 60
agcggcagtt gtctatcaaa tctatcaagt cgccttagc gctcaggaag tacgacaccg 120

```



```

gaaggggtgg gctttgcgaa gatggcgccg ctgggggtgc tggagtccga cctgccaaagt 180
gccgtgacac ttctgaaaaa tctccaggag caagtgatgg ctgtaactgc acaagtgaaa 240
tcactgacac aaaaagttca agctgggtgcc tatcctacag aaaagggtct cagcttcttg 300
gaagtgaaag accagctgct gctcatgtac cttatggatt tgaccacact cattctggac 360
aaagcctcag gaggatctct tcagggacat gatgcagttt tgagactggg agagattcga 420
acggtttttg aaaagcttcg tcccttgagc caaaagctga agtatcaa atgacaagctg 480
atcaagactg cagtgcacag cagccttagt gagaatgacc cacttcgttt taagcctcat 540
cccagcaata tgatgagcaa gttgagctct gaggatgagg aggaagatga agcagaagat 600
gaccagctct aggccttcagg gaagaaatct gtgaaggag tgtctaagaa atatgttcct 660
ccacgcttgg ttccagtaca ttatgatgaa acagaagctg agcgggagaa gaagcgtcta 720
gaacgagcca agagacgggc attgagcagc tctgtcattc gtgaacttaa ggagcagtac 780
tcagatgctc cagaggaaat ccgtgatgct cggcatcccc atgttaccgg ccagagtcag 840
gaggaccaac acaggattaa ctatgaggag agcatgatgg tgcgtttgag cgtcagtaag 900
cgagagaaaag gacggcgaaa acgagcaaat gtcatgagct cacaacttca ttcccttaca 960
cacttcagtg acatcagtg tttgacaggg ggaactgttc atcttgatga ggatcagaat 1020
cctattaaga agcggaagaa gatacctcag aaaggctcga agaaaaaagg ttttcggagg 1080
cggcggtgat tatgggtgta catatttgta tttttttgt catcctgaga tacttcta at 1140
ttcattgtat ataggtggtt ttccctggaa ttcattaatt gtttgctttg gacatgtgga 1200
aagagcctta ctaataaaat tgattttact tatgaaaaaa aaaaaaaaaa 1249

```

<210> 87
 <211> 1064
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1441770CB1

```

<400> 87
cggctctcca gagcgtctgt aaacaccag agactgtcat ggagggggag gaggaggcgg 60
cggcgccgaa gggaggcgtt tggggccgcc tccagggtcc gctctgccat tcctgaactg 120
gtccctcgtc cccgtgactc tggcatcagg gaagcgaact gttaggcgag aggaggaggc 180
agccagaaac atatccctt cttcctcggg gcggggggcc ggccaggccg gctgagcgg 240
gggaggggctc cgggaggggag tgcttgccca ggccggcctg tctgccgga tggatgacag 300
taaggtggtt ggaggcaaaag taaagaagcc cggtaaacgt ggtcgggaagc cagccaaaat 360
tgacttgaaa gcaaaacttg agaggagccg gcagagtga agagaatgcc gagcccgaaa 420
aaagctgaga tatcagtatt tggaaagatt ggtatccagt cgagaaagag ctatatgtgc 480
cctcagagag gaactggaaa tgtacaagca gtggtgcatg gcaatggacc aaggaaaaat 540
cccttctgaa ataaaggccc tactcactgg agaagagcag aacaaatctc agcagaactc 600
aagcaggcat accaaggctg ggaagacaga tgctaatagc aattcctggt gaagattata 660
taaagatgag tcagtgattg aagccaatat tctgattccc atggaagatg gatgggcaag 720
agtgtacttc ttggctccat ttactaccta ctgctcagta gtcactcttg taaatctgca 780
atttctacca aaatgtgtga tcgtagatct caaaggatct tgctttaact ttcaacactt 840
agaaaatcta caaacattca gacctgtctg ggttggtatt gccacccatg acatttaaca 900
tgttgtgatg cttgaaaaca caggagtaga gaaaatcgat gaagattgta tttttgcacc 960
ttaactccac attgctttat tgggttaatt atattctttc catgtaattc atgtaattgt 1020
atgtctgtgt gtgttttatg tgtcaccacc tttcatgttt ttga 1064

```

<210> 88
 <211> 1398
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 1456684CB1

```

<400> 88
agacccaaaa gcattaagga gaaaaagaaa actacatcac ataccagggg agaaataccg 60

```

```

gaggagtcaa actatgttgc tgatcctgga ggatcactga gcaaaaccac aaatattgct 120
gaagaaacca gcaaaattga aacctacatt gcaaaacctg ctctgccggg aacctccaca 180
aatagtaaatg ttgcacccct ttgccaaata acagtgaataa ttggaaacga agccattgtg 240
aaaaggcaca ttctaggatc taaattgttt tataaaagag ggagaagacc caagtatcag 300
atgcaggagg agcctttgcc acaggggaat gaccagaac ccagtggaga cagcccactc 360
gggctttgcc aatccgagtg catggagatg agtgaagtgt tcgatgacgc aagtgaccag 420
gattccactg acaaaccgtg gcgcccttac tacaactaca aacccaaaaa gaaatccaga 480
cagttgaaaa aaatgaggaa agtcaactgg aggaaggagc acggaacag gagcccagc 540
cataaatgta aatacccagc agaactggat tgcgcgtgg ggaaggctcc tcaggataaa 600
ccctttgagg aagaagaaac taaagagatg ccaagctgc agtgtgaact ctgtgatgga 660
gacaaagcag tgggggctgg aaaccaagga agggccacc gacatcttac ttctggcca 720
tatgcctcgc agctctgcgc caagcagttc cagagccctt ccacactcaa aatgcacatg 780
agatgtcaca ccggggagaa gccataccag tgcaagacct gcggacgggtg cttttcgggtg 840
caaggaaact tacagaaaca tgaacgcac cactggggt tgaaggagt cgtctgtcag 900
tattgcaaca aggcattcac cttgaatgag accctcaaaa tccatgaaag aatccatact 960
ggagaaaagc gttaccactg tcagtctctgc ttccagagat tttgtatct ctccaccaa 1020
aggaatcacg agcagaggca tattcgggag cataatggga agggctatgc ctgcttccag 1080
tgcccaaaaa ttgcaaaaac agctgctgcc cttggaatgc accaaaagaa acacttattc 1140
aaaagcccaa gtcagcagga gaaaataggt gacgtgtgcc acgaaaactc aaatcccttg 1200
gagaatcaac atttcattgg ttcagaagac aatgacaaa aggataacat acaaaccggt 1260
gtggaaaatg ttgtcctttg agtggcaaga attagaaaaa tcttcaaaaa tatagttggt 1320
ggttttttta gttatgattt aagtttagtt tcattttgtc catgtgacag tcatgaagga 1380
gtgaaaaaaa aaaaaaaa

```

<210> 89

<211> 746

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 1602916CB1

<400> 89

```

cggctcgagg cgtctttatg ggcccccttt aaggccggcg gaggcattctc gggccggggcg 60
cggcgctccg tccgtcgggc gtagcgactg aactgcgcgc ggatccctcc ggggggctcc 120
tcgtcccctg cacgctgact ttccgtgcag tgccgtggtg cgaaaatgac tcgccggtgc 180
gcaccggaga cagccgattt ttacgacca gcaagaggcc gagctggtac aatatcctga 240
ctgtaaatcg tccagtggta atattggcga ggaccagac cacttaaatc agagctcgtc 300
tccttctcaa atgtttccgt ggatgagacc acaagcagct cctggtagac gaagaggaag 360
acaaacctac agtcgcttcc aaactctaga gttggaaaag gaatttcttt ttaaccctta 420
tctgaccagg aaaagaagaa tcgaggttcc ccacgcccta gccctcacgg agagacaggt 480
aaaaatctgg ttccagaaca ggagaatgaa atggaaaaag gaaaacaaca aggacaaatt 540
tcccgtttcc cggcaggagg tgaaggacgg ggaacgaaa aaggaaagccc aagagctgga 600
ggaagacaga gccgaacgct tgacaaatta actttacct ttaaaattta ccacagacta 660
ttaaaactaa taatcaccat atgctgtgga caccacctat tttctttggt ggaagagacc 720
ttactgtgtt tcaagctacc ttcagt

```

<210> 90

<211> 1270

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 1692816CB1

<400> 90

```

gttggtcacg tgggtgttcg gagcgggcga gcggagttag cagggtttta ctgcagagcg 60
cgccgggcac tccagcgacc gtggggatca gcgtaggtga gctgtggcct tttgcgaggt 120

```

```

gctgcagcca tagctacgtg cgttcgctac gaggattgag cgtctccacc cagtaagtgg 180
gcaagaggcg gcaggaagtg ggtacgcagg ggcgcaaggc gcacagcctc tagacgactc 240
gctttccctc cggccaacct ctgaagccgc gtccctacttt gacagctgca gggccgcggc 300
ctgggtcttct gtgcttcacc atctacataa tgaatcccag tatgaagcag aaacaagaag 360
aaatcaaaga gaatatataa aatagttctg tcccaagaag aactctgaag atgattcagc 420
cttctgcctc tggatctctt gttggaagag aaaatgagct gtccgcaggc ttgtccaaaa 480
ggaaacatcg gaatgaccac ttaacatcta caacttocag ccctgggggtt attgtcccag 540
aatctagtga aaataaaaaa cttggaggag tcaccagga gtcatttgat cttatgatta 600
aagaaaaatc atcctctcag tattggaagg aagtggcaga aaaacggaga aaggcgctgt 660
atgaagcact taaggaaaaa gagaaacttc ataaagaaat tgaacaaaag gacaatgaaa 720
ttgcccgctc gaaaaaggag aataaagaac tggcagaagt agcagaacat gtacagtata 780
tggcagagct aatagagaga ctgaatgggtg aacctctgga taattttgaa tcaactggata 840
atcaggaatt tgattctgaa gaagaaactg ttgaggattc tctagtggaa gactcagaaa 900
ttggcacgtg tctgtaaggc actgtatctt cctctacgga tgcaaagcca tgtatatgaa 960
atgcattaat atttgactgt tgagaatctt actgccgaag ttacctcca ctagttcttt 1020
gtagcagagt acataactac ataatgccaa ctctgggaatc aaatttcctt gtttgaatcc 1080
tgggacccta ttgcattaaa gtacaaatac tatgtatctt taactatgaa tggtttatgt 1140
gaataggatt ttctcagttg tcagccatga cttatgttta ttactaaata aacttcaaac 1200
tcctgttgaa cattgtgtat aacttagaat aatgaaatat aaggagtatg tgtagaaaaa 1260
aaaaaaaaaa 1270

```

<210> 91

<211> 943

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 1968191CB1

<400> 91

```

gtcccatttc cagaaatcac aaggatgtta ggcaatgaat ggagtaaact gcctcctgag 60
gaaaaacagc gctaccttga tgaagcagac agagataagg agcgttacat gaaggaaactg 120
gaacagtatc agaaaacaga ggcctacaag gtcttcagta ggaaaaccca ggaccgtcag 180
aaaggcaaat ctcataggca agatgcagcc cggcaggcca ctcatgatca tgagaaagaa 240
acagaggtaa aggaacgggtc tggttttgac atccctatat ttacagagga attcttgaac 300
catagcaaaag ctcggaagag agagctccgc cagcttcgca aatccaacat ggagtttgag 360
gagaggaatg cagccctgca aaagcacgtg gagagcatgc gcacagcagt ggagaagctg 420
gaggtggatg tgatccagga gcggagccgc aacacagtct tacagcagca cctggagacc 480
ctcgggcagg tgctgaccag cagctttgcc agcatgccct tgccctggaag tggagagaca 540
cctacagtgg acaccattga ctcatatatg aacagactgc acagtattat tttagctaat 600
cccaagaca atgaaaactt catagctaca gtctgagaa ttgtgaacag actcgatcgt 660
tagggaatgg tgagtgtctc ctgataaata tttatatgcc agcacatcat caaaaataag 720
atgtcatcag actttatcaa tactactaaa accctgggat tacattggat gaacaagtgt 780
gagacttggg taagattcct gttgcatggg tgttaaatgt agtaataat attagaaaag 840
agaatcactg tagtcccagc tacttgggag gctgaggtgg gaggattgct tgagcccagc 900
agttcaagtc cagagagatc ctgtctctaa aaataaaaga aaa 943

```

<210> 92

<211> 1997

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2052061CB1

<400> 92

```

cgaccacgcg tccgctcgcg tggggcgcg ggtaccccg ggggcagcta ggtgcctcca 60
agaaaacccc cccagagga cccgcacgag ttgttgccat ttttcgctga agccctgcc 120

```

```

ctgggggtgg tgttcttctc tatgattcta ccaatcgctg gaattttttt cctcccttct 180
ttcttttttag gaaggcgctg ggggtgtgcg cggaacggc ggcgggtgaag ggatcctctt 240
gtgggtatctc ctccccggag aaatagggga gtggggggccc .aagaacgaga agacgagaac 300
gcgctcgccct gcgctatgtc agaatggggc ggggtgtgagg ggaacagctc tcttgcgac 360
agatcaggag tatgagcctc cgggaggacg gcatgagttc tggacacttc aggagtccct 420
agctagtgtc atggctcgata tggataaaact cataaacaac ttggaggtcc aacttaattc 480
agaaggtggc tcaatgcagg tattcaagca ggtcactgct tctgttcgga acagagatcc 540
ccctgagata gaatacacaa gtaatatgac ttctccaaca ctctggatg ccaaccccat 600
ggagaaccca gcaactgttta atgacatcaa gattgagccc ccagaagaac ttttggttag 660
tgatttcagc ctgcccgaag tgggaaccagt tgacctctcc ttccacaagc ccaaggtctc 720
tctccagcct gctagtcagc tacaagctcc aatacgtccc cccaagccac agtcttctcc 780
ccagaccctt gtgggtgtcca cgtcaacatc tgacatgagc acttcagcaa acattcctac 840
tgttctgacc ccaggctctg tccctgacct ctctcagagc actggtagcc agcagatctt 900
acatgtcatt cacactatcc cctcagtcag tctgccaagt aagatgggtg ggctgaagac 960
catcccaatg gtagtgcagt ctctgcccac ggtgtatact actttgctg cagatggggg 1020
gcctgcagcc attacagtc cactcattgg gggagatggg aaaaatgctg gatcagtgaa 1080
agttgacccc acctccatgt ctccactgga aattccaagt gacagtgagg agagtacaat 1140
tgagagtgga tctcagcct tgacaggtct gcagggacta cagcaagaac cagcagcaat 1200
ggcccaaatg cagggagaag agtcgcttga cttgaagaga agacggattc accaatgtga 1260
ctttgcagga tgcagcaag tgtaacacaa aagctctcac ctgaaagctc accgcagaat 1320
ccatacagga gagaagcctt ataaatgcac ctgggatggc tgctcctgga aatttgctcg 1380
ctcagatgag ctactcgcc atttccgcaa gcacacaggg atcaagcctt ttgggtgcac 1440
agactgcaac cgcagctttt ctggttctga ccacctgtcc ctgcatcgcc gtcgcatga 1500
caccatgtga gccgcacagg tcacactaga gaagctgcgc tgggtatctt cctggctctg 1560
tgctgaggtt gggacaattt tttctctt gactcagct tgcatatggg gttgaagcag 1620
cccactgagc caagttgagg agactggagg aaaagagagc tgggtctccc tggggctctt 1680
catattctac ctccacttct ccaactgtcc gacctgtttt ttcaacccc cacatgggtt 1740
gacttccagc gtggcaccac tgggtgcctt cccatcccc cctgttctga aatagggaat 1800
ttttccccc gcaacaccac aggaatcaaa ctcaaggctg gcaaccacat ccgctgttct 1860
ttctccccc ttccctcttg tctctagaac tcttatccaa tgtcttaaca tctaccaaa 1920
agggctcgat cgtttccaca gatttttcc tctttattgc agggatacat tegtggctgc 1980
cacataaggg ttgcttc

```

<210> 93

<211> 4334

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2056207CB1

<400> 93

```

cgggtgggtgg ggcggggcgg ggcagtgagc ggaggaggat taccgctacg aggtgctcac 60
ggccgagcag attctacaac acatgggtgga atgtatccgg gaggtcaacg aggtcatcca 120
gaatccagca actatcacaa gaatactcct tagccacttc aattgggata aagagaagct 180
aatggaaaagg tactttgatg gaaacctgga gaagctcttt gctgagtgtc atgtaattaa 240
tccaagtaaa aagtctcgaa cagccagat gaatacaagg tcatcagcac aggatatgcc 300
ttgtcagatc tgctacttga actaccctaa ctgctatttc actggccttg aatgtggaca 360
taagttttgt atgcagtgtc ggagtgaata tttaactacc aaaataatgg aagaaggcat 420
gggtcagact atttcgtgtc ctgctcatgg ttgtgatc ttagtggatg acaacacagt 480
tatgcgctg atcacagatt caaaagttaa attaaagtat cagcatttaa taacaaatag 540
ctttgtagag tgcaatcgac tgttaaagtg gtgtcctgcc ccagattgcc accatgttgt 600
taaagtccaa tctcctgatg ctaaacctgt tcgctgcaaa tggggcgccc aattttgctt 660
taactgtgga gaaaattggc atgatcctgt taaatgtaag tgggttaaga aatggattaa 720
aaagtgtgat gatgacagtg aaacctccaa ttggattgca gccaacacaa aggaatgtcc 780
caaatgccat gtcacaattg agaaggatgg tgggtgtaat cacatggctc gtcgtaacca 840
gaattgtaaa gcagagtttt gctgggtgtg tcttgcccca tgggaaccac atggatctgc 900
ctgggtacaac tgtaaccgct ataatagaag tgatgcaaag gcagcaagag atgcacagga 960
gcgatctagg gcagccctgc agaggtacct gttctactgt aatcgctata tgaaccacat 1020
gcagagcctg cgctttgagc acaactata tgctcaggtg aaacagaaaa tggaggagat 1080
gcagcagcac aacatgtcct ggattgaggt gcagtctctg aagaaggcag ttgatgtcct 1140

```

```

ctgccagtgt cgtgccacac tcatgtacac ttatgtcttc gctttctacc tcaaaaagaa 1200
taaccagtcct attatctttg agaataacca agcagatcta gagaatgcc aagaggtgct 1260
ctcggggtac cttgaacgag atatttccca agattctctg caggatataa agcagaaagt 1320
acaagacaag tacagatact gtgagagtcg acgaagggtt ttgttacagc atgtgcatga 1380
aggctatgaa aaagatctgt gggagtacat tgaggactga gaatggccct gcataaaatg 1440
aactctgaaa actttaccat cttagtggtc catgcaatta aaacaaaaca aacacaaaaca 1500
aggaggcact aagcctattc tgacaccact ggtctgtagt accagaattg ttttgtaaat 1560
ggaaagttta agtaaattat attgtaataa aaaggtagat aaaccattgt acaacagtat 1620
tctaggccgc caacaaaagt gtgacagaca cactaaaagc cctccaactt taacttgtaa 1680
cgtagcttca ttctcaaagc tgactccttt tttttctttt tccttttctt gagtgtagta 1740
cagttaaaat ttcaaacagc tccttgacac tgcttttcat gttcaaacca gccattttgt 1800
tgtacttttg taaaggacct cttccccttc ctcccctaca catacagata cccccacaca 1860
cagactgact ctctttctct cataccctaa ggtcatgagt gaatgatgct tagttccttg 1920
taaaagaaaat cttgggatgg ggaagggggt aggcagcaag aggattcaac aaacgaaaaa 1980
cataaaaact ttgtatatga cttttaaaac aagaggacaa cacagtattt ttcaaaattg 2040
tatatagcgc atatgcatgg acaagcaag cgtggcacgt gtttgcataa tgtttaatta 2100
caaaaaataa tttattcttt aaaaatcttc aagattatgt ctatttgctg tgcattttct 2160
ttcactttgc ttctctttcc cgggttgggg ttgggataaa ggtgtgtcgg tttagcacct 2220
ctggaagacc tatctagagc ttcttcaact tcctgaggtt attttgccct ttctgggtgt 2280
ggtatgtctg ttgcccgcga tgggcctcat gccttgaatt cctgctcttg atcagggaca 2340
agggaggtca agctctgact aatgccatga cctgattaag ggttacagca gggagtttg 2400
ttcctacagc ttctggaatta acctgtccca acctaattcc cctccatggc atcatgcctc 2460
taccacaagc tttgtgtgcc catgttatgc acacagctgc aggcattctt aagtccctg 2520
tcgcattccg tggagcatt ttaaaatttc ttttactttt tggttttccc ttaattgctg 2580
cttttcagat tttagttatg gctcgtctgc tcaccccttc tctacattag ggtgtcaaa 2640
agaatgtttt gctttaaata taaatagcca ttcatttagt ctcagattgt gaatttaaaa 2700
tgggtgtatc cgaattgtct tgtgtgtgtt gctgtgggtt tggtttgaag gcaaacaccc 2760
ctagaacatg atattcccat ctagtgcatt taaatagaaa tcaactgagt tgctgctttt 2820
ttattgtcag cagataggag aattaataat gcatttttagc tgtgatgtcc atttttatga 2880
aattcctact aagagctatg ttaaaagtaa aggatgggtg tggttgtatt aactatata 2940
ctgtttaggc cattctggct gtggtatttt agcatctgta aatctgtcag 3000
ttttatagag gagtgcagag tgaactagcc aactagatta agaggtctaa atatgaaat 3060
ccagttgagg ctgaggacct ctctgtcttc ctttagatgt cttttgccta gggagtgttt 3120
accatttgtg aggcagcttt gtctgtcttc acactgtaca tcctattact ccattgggaa 3180
gtagggtcac tttcctctgg ctttttgctt aagttaggct ttgctgaatc aaccctactt 3240
ttccttttag aaaaggttgt tacaggagat ttactggcaa ctgttctttt cccatcaaaa 3300
atcagtgaat gtttgctgag tataaatgct gcttccttaa accactgtgc gctttaggat 3360
caactttacc tgtacctttt ctcttttctt cccttgccac ctcaggtgca aatctgaact 3420
cagtgtctgc ttcttccatt ttctcgtctc tctcccctct tccccatta tccatatgac 3480
attattttac ttcaaatgac agcatcaatc ttaaaaagat atacattaaa actaaggagt 3540
ttttttaaag aaagcctgaa taagttcctt tccctggtaa ctttgaaaag cagtcagagt 3600
tgctatatag atatatgtgg ctcttttaaa atgctttgtg tatgtgtggt gtttaaaaaa 3660
aaaaatctta ccagttaact ttgcagtgtc taggtttgag tgtcataaat ccacgtgttc 3720
ctgttgcaac aaatacccaa aaattgtgtg tgcacttctt aataccagtc ttcacccatg 3780
gaggaacagt gctttttaga gatgctttct atttcaatgt tggcatactg cctgagggtg 3840
ttgcagttgt ggtgcatc cctaatttgt atgatcaaga tgaactggcc cttttctact 3900
tccaagcttt taacagatac caccatattt gacagaattc ccagagtga ttgcttgtgt 3960
tattagtaga ttcagtgcct ccagctggga taggcaagcc atgacagctt ccctgtttca 4020
cctacagaag tcttatctga gggatctatt cacagtaagc accaaggctt ccatgtcttg 4080
aggtcagttt cattgtcttt tgaaaagtgc atgcttcatt tgaacaatc attcagcagc 4140
agatggactt tcagtgtatt aaaataaaat tttgatccaa agctcaggac acaaaaccac 4200
gtggtaaaat tgagtgcatt ataatacag actaaattat ctgtaatttt ccacaaccca 4260
gattgtatgt gttttatgtg tgtttaaata aatatgttag atacacgtgt atacatacac 4320
ccatatacag caga 4334

```

<210> 94
<211> 1706
<212> DNA
<213> Homo sapiens

<220>
<221> misc-feature

<223> Incyte ID No.: 2101803CB1

<400> 94

```

ccgcgcceccg cccgccaccat gaggggcccag ggcctcggcg gcctggagcg cttctgcagc 60
ccgggcaaaag gccgggggct gcgggctctg cagcccttcc aggtggggga cttgctgttc 120
tcctgcccgg cctatgccta cgtgctcacg gtcaacgagc ggggcaacca ctgcgagtac 180
tgcttcacca ggaaagaagg attgtccaaa tgtggaagat gcaagcaggc attttactgc 240
aatgtggagt gtcagaaaaga agattggccc atgcacaagc tggaaatgttc tcccatggtt 300
gtttttgggg aaaactggaa tccctcggag actgtaagac taacagcaag gattctggcc 360
aaacagaaaaa tccaccaga gagaacacct tcggaaaaat tgttagctgt gaaggagttt 420
gaatcacatc tggataagtt agacaatgag aagaaggatt tgattcagag tgacatagct 480
gctctccatc acttttactc caagcatctc gaattccctg acaatgatag cctcgtagta 540
ctctttgcac aggttaactg taatggcttc acaattgaag atgaagaact ttctcatttg 600
ggtacgcaga tatttcctga tgttgcatgt atgaatcata gctgttgccc caatgtcatt 660
gtgacctaca aagggaccct ggcagaagtc agagctgtac aggaaatcaa gccgggagag 720
gaggttttta ccagctatat tgatctcctg taccacaacg aagatagaaa tgaccgggta 780
agagattctt atttctttac ctgtgagtgc caggagtgtt ccaccaagga caaggataag 840
gccaagtggt aaatccggaa gctcagcgat ccccaagg cagaagccat ccgagacatg 900
gtcagatatg cacgcaacgt cattgaagag ttccggaggg ccaagcacta taaatccct 960
agtgaagctg tggagatctg cgagctcagc caggagaaga tgagctctgt gtttgaggac 1020
agtaacgtgt acatgttgca catgatgtac caggccatgg gtgtctgctt gtacatgcag 1080
gactgggaag gagccctgca atatggacag aaaatcatta agccctacag taagcactat 1140
cctttgtact cctccaacgt ggcctccatg tgggtgaagc tagggagact ctacatgggc 1200
ctggaacaca aagccgcagg ggagaaagcc ctgaagaagg ccattgcaat catggaagta 1260
gctcacggca aagatcatcc atatatcttct gagatcaaac aggaaattga aagccactga 1320
aactatgcag catttcagtt ttcatttaaa cacttagttc agaaacctta aaggatttga 1380
atatttcaaa ttgcacagct cactocagca tctctgtaaa ataattgga tgaaaatact 1440
tcttgcaact aaacactgca catgccgtac tttgagggta gtctgaatct tgaactttaa 1500
taccaaaatta attttgaatg cttttgtttc ctaagagata atggcatggt ttcatatggt 1560
atactttgga cagacagagt tttaaaaatg gaattatttt ttctttcatg cctcttgtta 1620
tgttctgaac aaacttgaat gatgaaagta ttaaagagat atcagtaaaa agaacaaaaa 1680
ataaagatcc agaaagaaaa aaaggg 1706

```

<210> 95

<211> 1602

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2112362CB1

<400> 95

```

ccgtggggca gtgcaggatg tcgggtgaatt acgcggcggg gctgtcgccg tacgaggaca 60
agggcaagtg cggcctcccg gagatcttcg acccccggga ggagctggag cggaagggtg 120
gggaactggc gaggtgggtc tggcagttct ccaatgtggg gttccacacg ggtgccggca 180
tcagcactgc ctctggcatc cccgacttca ggggtcccca cggagtctgg accatggagg 240
agcgagggtc ggcccccaag ttgcacacca cttttgagag cgcgcggccc acgcagacc 300
acatggcgct ggtgcagctg gagecgcgtg gectcctccg cttcctgggtc agccagaacg 360
tggacgggct ccagtgcgc tcaggcttcc ccagggacaa actggcagag ctccacggga 420
acatgtttgt ggaagaatgt gccaaagtga agacgcagta cgtccgagac acagtctgtg 480
gcaccatggg cctgaaggcc acgggccggc tctgcaccgt ggctaaggca agggggctgc 540
gagcctgcag gggagagctg agggacacca tcctagactg ggaggactcc ctgcccagac 600
gggacctggc actcggcatg gagggcagca ggaacgcgc cctgtccatc acgtgggta 660
catcgctgca gatccggccc agcgggaacc tggcgtggc taccaagcgc cggggaggcc 720
gcctggctcat cgtcaacctg cagcccacca agcacgaccg ccagtctgac ctccgcatcc 780
atggctacgt tgacgaggtc atgaccggc tcatgaagca cctggggctg gagatccccg 840
cctgggacgg ccccgtgtg ctggagaggg cgtgccacc cctgccccgc ccgccaccc 900
ccaagctgga gcccaaggga gaatctccca cccggatcaa cggctctatc ccgcgcggcc 960
ccaagcagga gccctgcgcc cagcacaacg gctcagagcc cgcagcccc aaacgggagc 1020
ggcccaccag ccctgcccc cagagacccc ccaaagggt gaaggccaag gcggtcccca 1080
gctgaccagg gtgcttgggg aggggtgggg tttttgtaga aactgtggat tctttttctc 1140

```

```

tcgtgggtctc actttgttac ttgtttctgt ccccgaggagc ctcaggggctc tgagagctgt 1200
gctccagggcc aggggttaca cctgccctcc gtggtccctc cctggggctcc aggggcctct 1260
gggtcggttc cgggaagaag ccacacccca gaggtgacag ctgagcccct gccacacccc 1320
agcctctgac ttgctgtgtt gtccagaggt gaggtctgggc cctccctggt ctccagctta 1380
aacaggagtg aactccctct gtccccaggg cctcccttct gggcccccta cagcccaccc 1440
taccctcct ccatgggccc tgcaggaggg gagaccacac ttgaagtggg ggatcagtag 1500
aggcttgac tgcttttggg gctggaggga gacgtgggtc caccaggctt ctggaaaagt 1560
cctcaatgca ataaaaacaa tttctttctt gcaaaaaaaaa aa 1602

```

<210> 96
 <211> 1951
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 2117346CB1

```

<400> 96
gaggcgcgcg cgaccggcgg ctcttttggcg cggattaggg ggtctcggcg agggagtcac 60
caagcttttg tgtatgtgtt ggccgggttct gaagtcttga agaagctctg ctgaggaaga 120
ccaaagcagc actcgttgcc aattagggaa tggaccgttt gggttccctt agcaatgac 180
cctctgataa gccaccttgc cgaggctgct cctcctacct catggagcct tatatcaagt 240
gtgctgaatg tgggccacct ccttttttcc tctgcttgca gtgtttcact cgaggctttg 300
agtacaagaa acatcaaagc gatcatactt atgaaataat gacttcagat tttcctgtcc 360
ttgatcccag ctggactgct caagaagaaa tggccctttt agaagctgtg atggactgtg 420
gcttttgaaa ttggcaggat gttagccaatc aaatgtgcac caagaccaag gaggagtgtg 480
agaagcacta tatgaagcat ttcataata accctctgtt tgcactacc ctgctgaacc 540
tgaaacaagc agaggaagca aaaactgctg acacagccat tccatttcac tctacagatg 600
accctccccg acctaccttt gactccttgc tttctcgga catggccggg tacatgccag 660
ctcgagcaga tttcattgag gaatttgaca attatgcaga atgggacttg agagacattg 720
attttgttga agtgactcg gacattttac atgctctgaa gatggctgtg gtagatatct 780
atcattccag gttaaaggag agacaaagac gaaaaaaaaat tataagagac catggattaa 840
tcaaccttag aaagtttcaa ttaatggaac ggcgttatcc caaggaggct caggacctgt 900
atgaaacaat gaggcgattt gcaagaattg tggggccagt ggaacatgac aaattcattg 960
aaagccatgc attggaattt gaactccgaa gggaaatcaa gaggctccaa gaatacagga 1020
cagcaggcat taccaatttt tgtagtgcga gaacctacga tcacctcaag aagacacggg 1080
aggaagagcg ccttaaagcg actatgctct cagaagttct ccagtatatc caggacagta 1140
gtgcttgcca gcagtggctc cgccggcaag ctgacattga ttccggcctg agtccttcca 1200
ttccaatggc ttcgaaattca ggtagacgga gtgcaccacc cttgaacctc actggcctcc 1260
ctggcacaga gaagctgaat gaaaaagaaa aggagctctg tcagatgggt aggttgggtcc 1320
ctggagccta tttagaatac aaatctgctc tattgaacga atgtaacaag caaggaggct 1380
taagactggc gcaggcaaga gactcatca agatagatgt gaacaaaacc cggaaaatct 1440
atgatttcct catcagagaa ggatacatca ctaaaggcta aggtctccaa agcttgggat 1500
cagaagtcag aagtttggaa tgtgtgggt caaaggacaa tatgggtggg cattctggag 1560
agttgttttt cagctgaatt ctcatggtga aaacagggga aaggacaaag gaaaccttaa 1620
gttgatttgt ctactttctt ctccatcctg ctttaaaaca ctctgtttgt tgggtattatg 1680
ctgcagagtt gtgtgctaca taagctatta ttaaagtgtg gtgggcattc attcctaaca 1740
cctctgttaa ctaaaagacc catagtacc tcacatcaca gtgcctggga gaaattggac 1800
ctaaaccac agtctgtaaa tcccaggagt tcaagcctag aatccttaat attgtccggg 1860
gaaagggttt tctgaataaa acacggcttt ggcctttaa aaaactttgg ggaaacgaat 1920
tttaaaaatt aaaagggaaa aggtttttt a 1951

```

<210> 97
 <211> 854
 <212> DNA
 <213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2119917CB1

<400> 97

```

cccacgcgtc cgcggacggt gggcgcgcgcc tatgacggcc agcgcacagc cgcgcgggcg 60
gcggccagga gtcggagtcg gagtcgtggt gaccagctgc aagcatccgc gttgcgtcct 120
cctgggggaag aggaaaggct cgggttgagc tggcagtttc caactccctg gaggtcatct 180
ggagttcgggt gaaacctggg aagaatgtgc tcaaagggaa acctgggaag aagcagctct 240
tcacctgaaa aatgttcact ttgcctcagt tgtgaattct ttcattgaga aggagaatta 300
ccattatgtt actatattaa tgaaaggaga agtggatgtg actcatgatt cagaaccaa 360
gaatgtagag cctgaaaaaa atgaaagtgt ggagtggtt ccttgggaag aactacctcc 420
cctggaccag cttttctggg gactgcgttg tttaaaagaa caaggctatg atccatttaa 480
agaagatctg aacctctgg tgggatacaa aggaatcat ctctagtggt ccgagaagat 540
ttgattttct ttaaaaagac aagaataagg tctggttagg gaatgaaaaa tgtatacatt 600
tcggaacaac tccattttat ctaaaaaagt tcttgtgatt gccagtttat ttgcagtctc 660
ttaatgtatc cccactctt tcagccagta cttgagaaaa tttttctgaa atatgtcatt 720
gaattgtatt ccagacacag aatacatgat aaatactgat attatgggta atctgcttct 780
catatttacc tatggatatg tacgtgcaat gtgccataac tagttgagag ggtgggtgag 840
gcagattttc cttt                                     854

```

<210> 98

<211> 1581

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2123456CB1

<400> 98

```

tccgggaaaag tttctttgga ggtccggccc ggagcggcca tgtcccacgg cccaagcag 60
cccggcgcgg ccgcctgtcc ggcggggcggc aaggctccgg gccagcatgg gggcttcgtg 120
gtgactgtca agcaagagcg cggcgagggg ccacgcgcgg gcgagaaggg gtcccacgag 180
gaggagcccg gttctgcagc cgggtgaagaa acgcggctgg cccaagggca agaagcggaa 240
gaagattctg ccgaatgggc ccaaggcacc ggtcacgggc tacgtgcgct tcctgaacga 300
gcggcgcgag cagatccgca cgcgccaccc ggatctgccc tttcccgaga tcaccaagat 360
gctgggcgcc gagtggagca agctgcagcc aacggaaaag cagcggtagc tggatgaggc 420
cgagagagag aagcagcagt acatgaagga gctgcggggc taccagcagt ctgaagccta 480
taagatgtgc acggagaaga tccaggagaa gaagatcaag aaagaagact cgagctctgg 540
gctcatgaac actctcctga atggacacaa ggggtggggac tgcgatggct tctccacctt 600
cgatgtttcc atcttctact aagagttctt ggacaaaac aaagcgcgtg aggcggagct 660
tcggcgcttg cggaagatga atgtggcctt cgaggagcag aacgcggtac tgcagaggca 720
cacgcagagc atgagcagcg cgcgcgagcg tctggagcag gagctggcgc tggaggagcg 780
gaggacgctg gcgctgcagc agcagctcca ggccgtgcgc caggcgctca ccgccagctt 840
cgcctcactg ccggtgcggg gcacggggcga aacgcccacg ctgggcactc tggacttcta 900
catggcccgg cttcacggag ccaccccgcc cagcacgaga agctcatcgt 960
ccgcatcaag gaaatcctgg ccaggtcgc cagcgagcac ctgtgaggag tgggcgggce 1020
cacgatgcag aggagaagct gtgggcgcgg ccctgccaca cccaccccg tggacgagag 1080
gctgggggtc caccctttgg ggcctgggtc catcctgcac cttgggggct ccagccccc 1140
taaaattaaa tttctgcagc atccctttag ctttcaattc cccagcccc ctgaaccgg 1200
aaaaagcact cgctgcgcga tacaccaga agaacctcac agccgagggt gcccctctc 1260
ggaggacagc cacgcgctac actggctctc cgggccaccc ccaggacaca gggcagacga 1320
aaccaccccc cagcacacgg caggaccccc caaattactc actacggggg gctgtgccat 1380
aggccacaca ggaagctgcc ttgtggggac ttacctggg tgcctccgc atgctgtac 1440
cccagatggg tggggggcgg ctttggccat cctgctctcc tccagccgag ggaccctggt 1500
gggggtggct ccttctcact gctggatcgg gactttttta ataaaaacaa gtaaaatttg 1560
tgttttaaaa aaaaaaaaaa a                                     1581

```

<210> 99

<211> 2150

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2148792CB1

<400> 99

```

gtctcgatct cctgacctcg ccatccaccc gcctcagcca cccgaagtgt tgggatttca 60
ggggccaaat agttgcatca catgtatcta atccgagagt ctcatgcttc tggtagctcc 120
tcagtgaacca gctcctgtct actgccctca gaaagcccaa cctcaggca atggcggcct 180
tgttcctgtc tgccccaccc caggccgagg tgaccttcga ggacgtggct gtgtacctct 240
cccgaggagga atggggccgc ctgggccttg ctgagagggg cctctacagg gacgtgatgc 300
tggagacctc cgggaacctc gtctcactgg gagtaggacc tgcaggcccc aagcctggag 360
tgatctcgca gttggagcga ggggatgagc cctgggtcct ggatgttcag ggcacctctg 420
ggaaagagca cctgagagtc aacagcccag ctcttgggac cagaactgag tacaaggagt 480
tgacttcaca ggagacattt ggtgaggaag atccccaggg atctgagcca gtagaagcct 540
gtgaccacat cagtgaagtc gaggggagcc tggaaaagct agtggagcag agaggcccca 600
gggcagtcac actgaccaac ggggagagca gcagggagtc tgggggaaac ctgaggttgc 660
tgtcaagacc tgttcctgat cagagacctc acaaatgtga tatatgtgag caaagttttg 720
aacagagatc atatctcaac aaccataagc gtgtacacag gtcaaaaaaa acaaatcacg 780
ttcgtaactc tggggaaatc tttagtgcaa acttagttgt taaagaagat cagaaaattc 840
ctactgggaa aaaattgcat tattgcagtt actgtgggaa aacattcagg tacagtgcga 900
accttgtaa gcatcagcgg cttcacactg aagagaagcc ctacaaatgt gatgagtgtg 960
ggaaagcctt cagccagagc tgcgagttca tcaatcaccc aaggatgcac tcaggagaga 1020
ttccctaccg gtgtgacgag tgtgggaaga cattcacccg gagggccaac ctcatgaagc 1080
accagaggat tcacactggg gagaaccctt acaagtgtgg ggagtgtggg aagcacttta 1140
gcgccacttc ttccctgatt tatcaccaga gaatccacac cggagagaaa ccctataaat 1200
gtaatgactg cgggaaagcc tttagtgatg gctcaatcct tatccgacat cgtcggactc 1260
acaccggaga gaagccattt gagtgcgaagg aatgtggcaa aggctttaca caaagtctta 1320
accttatcca aactcagaga attcacactg gagagaaacc ctataaatgt aatgaatgtg 1380
agaaagcttt cattcaaaaa accaaactcg tggaaacatc gagaagccac actggagaga 1440
agccctatga atgcaatgac tgtggcaaaag ttttcagcca aagcacacac ctcatccagc 1500
accagagaat ccacacagga gagaagccct acaagtgcag cgagtgtggg aaagccttcc 1560
acaacagttc cagactcact caccaccaga ggtgcacca cggagagaaa cctacagat 1620
gcagcgattg caagaaagcc tttagccaga gcacgtactt gattcagcac cggaggatcc 1680
acaccgggga gaagccctac aagtgcagcg agtgtgggaa ggccttcccg cacagtccca 1740
acatgtgtca gcatcagcgg attcacctcc gggaggactt ctccatgtaa cagtggcgcg 1800
gtgtccgagg ctagactcca gctgagcact tctgcatgc gcccccggca cctgactctg 1860
ccctttatgt attatccaca cgatgttttc acagagtgaagg aggcgtttc tcattaaaca 1920
aacctctttt cttaaatcaa aaaaaaaaaa agggggggcc gctttagggg tcccaggggt 1980
tcccaacccc taatcctaaa caatgtaata gctgttcccg gtgtaaaatt gttagcggcc 2040
caaaattccc tggcaaatca tcgaaccggg aatccttaaa tgtttaaaac ccccgggggtg 2100
gccccaaaaa aggttgact cgaaacttcc acattaaaat tcgggggggg 2150

```

<210> 100

<211> 691

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 2751943CB1

<400> 100

```

cggcgccgga gcggcgctca tggcgcggt cctctggttg ctccggggcc tgaccctcgg 60
aactgcgcct cggcgggcgg tgcggggcca agcggggcgg ggcgggcccg gcaccgggcc 120
gggactgggg gaggcagggt ctcttgcaac gtgtgagctg cctcttgcca agagtgaagt 180
gcaaaagaaa ctaaccccgg agcagttcta cgtcacaaga gaaaagggaa cggaaccgcc 240
tttcagtggt atctacctga ataacaagga agcaggaatg tatcattgag tgtgctgcga 300
cagtccactc tttagttctg agaaaaagta ctgctctggc actgggtggc cttcgttttc 360
cgaggctcat ggtacgtctg gctctgatga aagccacaca gggatcctga gacgtctgga 420

```

```

tacctcgtta ggatcagctc gcacagaggt tgtctgcaag cagtgtgaag ctcattctagg 480
tcacgtgttt cctgatggac ctggggcccaa tggctcagagg ttttgcatac acagtgtggc 540
tttgaagttc aaaccaagga aacctgacc atcttcaaga gtcccgttcc cttgccaccc 600
cttcacgtgc accctcaatt tccacaattc acttgaatga cttgttttat ttgcaataaa 660
actgggctga atttgcaaaa aaaaaaaaaa a 691

```

<210> 101

<211> 2101

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 3128913CB1

<400> 101

```

gtggccttga gctcgggggtg ggtggctcat ttcctggccg ctctcgggct tcgcggaaaag 60
aagagattac tcacactcct tctcaagcac agaaccagtt gtactgagct ttttgctaag 120
ctgttttcagc caagaatggc tgtggaatct ggagtgattt caaccctgat acctcaggat 180
cctccgggaaac aagaactaat actagtgaata gtagaagata acttttctctg ggatgagaaa 240
tttaagcaga atgggagtac tcaatcctgc caagaattgt ttcgtcagca attcagaaaa 300
ttttgctacc aggagacacc tggggcccgagg gaggctctga gccgactcca ggaactttgc 360
tatcagtggc taatgccaga gttgcacaca aaggagcaga tcttagaact gctggtactg 420
gagcagttcc tgagcattct gcctgaggag ctgcagatct gggttcagca acataatcca 480
gaaagcggcg aggaagctgt gacctgttg gaggatttag agaggaggtt tgatgacca 540
gggcagcagg tcccagctag tccacaggga ccagcagtg catggaagga tttaacatgt 600
ctcagagcat cccaagagtc aacagacatc cacctccagc ccttaaagac acagctgaaa 660
tcctggaaac catgcctttc ccctaaaagt gattgtgaga acagtgaac agcaacaaaa 720
gagggcatct cagaagaaaa atcacaggga ctccctcagg aaccttcatt tcgaggaatt 780
agtgcattcg aaagcaattt agtgtggaag caaggagtg ctacagggga gaaactaaga 840
tctccttccc aagggggcag ttttagtcaa gtgatcttca caaacaatc tctaggaaaag 900
agagaccttt atgatgaggc tgaaagatgc ttgattctaa ctacagactc tataatgtgt 960
cagaaagttc ctccagaaga gagaccttat agatgtgatg tatgtgggca cagcttcaag 1020
cagcattcct ctctaacaca acatcagaga atccatactg gagaaaagcc ctataaatgt 1080
aaccagtgtg ggaaggcctt tagtttgagg tcttatctta ttattcatca gagaattcat 1140
agtgggtgaga aagcatatga atgtagtga tgtgggaaaag ctttcaatca gagctcagcc 1200
ctcattagac atcgggaaat ccatactggt gagaaagctt gtaaatgtaa tgagtgtggc 1260
aaagcattca gtcaaaagttc atatctcatt atacatcaaa gaattcacac tggtgagaaa 1320
ccttatgagt gtaatgaatg tgggaaaacc ttttagccaga gctcaaaact cattagacat 1380
cagcgaattc acacaggaga gagaccctat gaatgtaatg aatgtggaaa agctttcagg 1440
cagagctcag agctgattac tcatcagaga atacatagtg gagagaaacc ctatgaatgt 1500
agtgaatgtg gaaaagcttt cagtttgagc tcaaaccctta tcagacatca gagaattcat 1560
agtggggagg aaccttatca gtgtaatgaa tgtggcaaaa ctttcaaaag gagctcagcc 1620
cttgttcagc atcagagaat tcattctggg gatgaagctt atatatgtaa tgaatgtggg 1680
aaggctttca ggcacagatc ggtccttatg cgccatcaaa gagtccacac tataaagtaa 1740
tttgtaata ctgtgaatag tgtaaatact tcagtcagat ttttaagttt gtagtcaaaa 1800
agagtttact ttggagcaaa actccataaa ggttataaaa tactaggctt tgagtctagc 1860
ttgctttgtg cagcatttcc cagtgcataa gtaaaagtgc ccttgaaagc ttttctctgt 1920
actaatcaga acagaatata gaagaatcat tacttccagc tcttctctta ttaggaatac 1980
tcaggaaata cgaagagtggt gaatgtaaca ttgaaacctc attttgtatg aaagtgtcat 2040
gaatatagca acccaggctc tgtcactgca tctgattgtt agtgtgccag gttatggggg 2100
g 2101

```

<210> 102

<211> 2196

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 3282941CB1

<400> 102
 cctacttttct ctagaccaga aaattcccaa aattgtgcac attagaatca cgtgggtagc 60
 ccacaaaatt cccagtgcc aattataacc tgcctttcct gtggcccagt ggtttccaac 120
 gacattgggc cctcgggtctg agaagtgcct ataatactgg cttacatggc cctcaggctg 180
 aaccttggac gtggcaggta ttgcaaatat ttcccgcggt gcgcgcggat tctggacttg 240
 ggcgcgaact cgtagctcac gctcccggg gtcagcagag gggcgtcacg ctctcgccac 300
 ccacctcgct ttctcaccct gcgcttcccg gcctgggttt ttagtcttcc ttggagcgct 360
 ctctggcctc cgcctccgcc agggagcggg aggcggagac agcgagactg gccagggggg 420
 aggaaagagg acgcgtgtgg gcaaggggga caacgggatg tccacgggct cgggtagtga 480
 tccgggtggc atggagcttc gggggctgca gcgggagtag ccggtcccg cctccaagag 540
 gccgcccctc cgcggcgtag agcgcagcta cgcctcgccc agtgacaact cgtcggcaga 600
 ggaggaggac cccgacggcg agggaggagc ctgcgctctg ggcacagcgg gcagcggga 660
 aggctgcaag aggaagcggc cccgtgtggc tggggcgggc ggcgcagggt gtagcgggg 720
 cgggtggggc aagaagcccc tcccggccaa gggtcagcc gcagagtga agcagtcgca 780
 gcggaacgcg gccaacgccc gtgagcgtgc cgggatgcgc gtgctgagca aagccttctc 840
 caggctcaag accagcctgc cctgggtgcc ccccgacact aagctctcca agctggacac 900
 gctccggctg gcttccagtt acatcgctca cctgcggcag ctgttgcaag aggaccgcta 960
 tgagaacggc tagtgacac cagtgaacct gacatggcca ttcgtggtct cgggaagacc 1020
 ggactctgac accaaagaag tttccgcagc caacagacta tgtggaacca ccgcttaaat 1080
 cggactggaa ctcaactgat gggattatc gttaaatgcg agtggttggg ggccacggag 1140
 agaagggaga gctcgtgaga tgggaagaag tttccgctgg attctccttg acccttcccc 1200
 ttccctgga actgtgatcg tgacaggtgg cgggtgtggc tgtcactgca cagcggccac 1260
 ggctacagct gcgcgggatc tgggcgacca gcttttgct ctccaaaaag agcttcttt 1320
 cgtgacgaga cgcggacgca ggtccaccct cgggccctag ctctgtagac taactctcgg 1380
 ctgctgcccc agcccgcgcc agacagccca cggatccgtt ctacagcgac gcagattcat 1440
 cgcacacgtg cgggacgggt ccacacagcc cggcctttc gcggtgacac aatggttagg 1500
 gaacggttag acgcgctct acatccgct taaagacaga ggtctagacg tgagatccgc 1560
 gtcgggacag ggttttaagt gacaaagaag ggcgagtggc ttctctgggc cgggttcgta 1620
 ctccagcaca gcgcccttct aacgggcggg aggaaggccg ctgctcgag ggctaggtgg 1680
 agacacactt cccagatcac cgcaggcggg ttttaccgg agagctctgg gccgttcggc 1740
 ctccctgccc ggtggcttct tcaatccgt ctccctccca agctccggc tttttctaat 1800
 caggcaggcg tctgtcaacc ctctccact ctgggctgaa gcctcccaa gccccgctgc 1860
 gccaacctgt gtgggttctt ctccggcct cctctccgc cccgctcctg ctctacctg 1920
 cagcaccctc agctccgact ccagactctc tgcacaggt cteccactc cagctccgg 1980
 gcgccccaac tccaacacca cgtcctgccc cgcaggttct tcccgcgcg gaggagcgcg 2040
 cagggtgggc ggcttaccat agcaagtgat cctgcgatag ggaacgcgc cttgccccga 2100
 ggctgcacta ccacaggaaa taacatatgt aaataaattt atttttttat gaataataaa 2160
 acgcgctgta aaaaccgtga aaaaaaaaaa aaaagg 2196

<210> 103
 <211> 749
 <212> DNA
 <213> Homo sapiens

<220>
 <221> unsure
 <222> 735, 736, 741
 <223> a or g or c or t, unknown, or other
 <223> Incyte ID No.: 3286656CB1

<400> 103
 gctgagctgt gctgcggcgc gcggcgcggc gcggtgcggc acggcacggg gggagtgtct 60
 ccggctggct tgcaggggaga acaccgactg agacctcaaa ccctggctcc agtgtcatgg 120
 aatccgtcac ctttgaggat gtggccgtgg agttcatcca ggagtgggca ttgctggaca 180
 gcgcacggag ggcctgtgac aaatacagga gtgcaggacc ctggcctcca 240
 ggggaactcc accatgcaaa cccagtgtgt tctccagct ggggcaaaga gcagagccaa 300
 aggcaacaga acgagggatt ctccgtgcca cagggtgttg ctgggaatct caacttaaac 360
 ccgaagagtt gccttctatg caggatcttt tggaagaagc atcctccagg gacatgcaaa 420
 tggggccggg cgtgttctg aggatgcagc ttggtgccct catagaagag agggagacac 480
 cattgactcg agaggaccgg ccagctctcc agggagcggc ttggtctctg ggatgcacgg 540
 gactgaaggc cgctatgcag attcagaggg tgggtgatac agtgctact ctggggccac 600
 gcaacccatg ggtggccagg gattctgcca tgtagcacgt gcctgcttcc cctttgcctt 660

ccgccatgat tgtaagtttc tcgagtcctc cccagccatg cgtcctgtac aacctgtgga 720
accaggcagc caaanncagt natagttgt 749

<210> 104
<211> 1311
<212> DNA
<213> Homo sapiens

<220>
<221> unsure
<222> 1294
<223> a or g or c or t, unknown, or other
<223> Incyte ID No.: 3490802CB1

<400> 104
gggcctttgt ttctcgtctg agcgaggagct ccagggtttat cctctgtgtt ctgtgtcctg 60
tgcttataga ggcccgctct ctgtggccgt gtgacctgca agtattggga gagccacagc 120
taaaccccgagg gacccctgga agcctagaaa tgggaccatt gcaatttaga gatgtggcca 180
tagaattctc tctggaggag tggcattgcc tggacactgc acagcggaat ttatataggg 240
atgtgatgtt agagaactac agaaacttgg tcttccttgg tattgttgtc tctaagccag 300
acctgggttac ctgtctggag caaggaaaaa aacctttaac tatggaaaaga catgagatga 360
ttgccaaacc ccaggttatg agttctcatt ttgcccaaga cctttggcca gagaacatac 420
aaaattcttt ccaaataagg atgctgagaa gatatgaaga atgcagacat gacaatttac 480
agttaaaaaa aggctgtaaa agcgtgggtg agcataaggt gcacaaaagg ggttataatg 540
gacttaacca atgtttgaca actaccaga aagaaatatt tcaatgtgat aaatatggaa 600
aagtctttca taagtcttca aattcaaca catataagac aagacatact ggaataaatc 660
ttttcaaatg tataatatgt ggcaaagctt ttaaaccgtc ctcaaccctt actacacata 720
agaaaattca tactggagag aaaccttaca aatgtgaaga atgtggcaaa gcttttaacc 780
aatcctcaaa ccttactaca cataagagaa ttcatactgg agagaaacct tacaatgtg 840
aagactgtgg caagctttt aactggtcct cagaccttaa taaacataag aaaattcata 900
ttgaacgaaa accctacata gtgaagaatg tgacagatct tttaaatgtt cctccacttt 960
taattagcat aagataattc atactggaga gaaaccttat gaatgtgatg aatgtgggaa 1020
agcctttaac cagccctcga ctcttagtaa atttgagagt ttatatggaa cacaaacctt 1080
acaaataaca agaatgtgac aaagcttttt aaggaaagttc tcaaccotta ttacacataa 1140
ttcatacca acagaagccc tacaagtgtg aagaatgtgg caaaacctat aaacctataa 1200
caagttctca attccttttt ttttgagatg gagtttctact cttgtcaccc aggcagaggt 1260
tgcagtgagc actccagctc aggcgacaga gtangccttg tcgcgatccc a 1311

<210> 105
<211> 990
<212> DNA
<213> Homo sapiens

<220>
<221>
<223> Incyte ID No.: 3507366CB1

<400> 105
caaaaaggaa agaataaggc aaattcttag aatgtatgca caacttaata tccttgctcg 60
tttgatgtgc attgatctca tttaataggt ttttgaatct gaaaattcaa gaagggtgaag 120
ctcacacat tttttgccct gcatatgatt gcttccaact tgtacctgtg gatcatatag 180
aaagtgtagt ttcaaaggag atggacaaac gatacctaca gtttgatatt aaggcctttg 240
ttgaaaataa tcctgccatt aaatgggtgc ctactccagg ctgtgacaga gcagtaagac 300
taacgaataa agagtcattt acatctggat ctgatacact cagcttccca ttgctgagag 360
ctcctgctgt tgattgtgga aaaggacacc tcttctgctg ggagtgcctt ggtgaagcac 420
atgagccttg tgactgcaa acatggaaga attggctgca aaaaataacc gaaatgaaac 480
cagaagaact tgtgggagtt agtgaagcct acgaggatgc cgccaattgt ctctgggtat 540
taactaactc caagctttag gccaactgta agtctccaat acagaagaat gaaggctgca 600
atcacatgca gtgtgctaag tgcaagtatg acttttgctg gatttgcctt gaagagtgga 660
aaaaacatag ttcgtccact ggagggtatt acagatgtac tcgctatgaa gtcattcaac 720
acgtggagga gcaatccaag gaaatgactg tggaggctga gaaaaaacac aaacgatttc 780

```

aggaacttga cagatttatg cactattata caagatttaa aaaccatgag catagtattc 840
agctagaaca acgccttctt aaaacagcca aagaaaagat ggagcaaatg agcagagtct 900
caaaagaactg aaggaggctg tccagatacc actttcattg gagatgagtt catgtgctct 960
aaaaatcggc gcatctcaag tgtcttatca 990

```

<210> 106

<211> 1048

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 3573060CB1

<400> 106

```

ccgctgcagc tctccgcggg acatctcacc gttctggaga cagggctcgc tcgctctcac 60
ggcttcttag gccgggggtg gacagccgcc ttccggccag aggggatgag gttgcgctgc 120
gctccggggag cgcgatggc gtgactggcc ccgcgcggag cagcgacact gcccggccag 180
cccgttcttc tgcccgagc catgaatctc agtagcgcca gtagcacgga ggaaaaggca 240
gtgacgaccg tgctctgggg ctgcgagctc agtcaggaga ggcggacttg gaccttcaga 300
cccagctgg aggggaagca gagctgcagg ctgttgcttc atacgatttg cttgggggag 360
aaagccaaag agggagagga agagggaagag gaagatgatg aggatgagga tgcagatata 420
aagatgcagc cggtcaccat tgcctcactc caggcctcag tctctcccat ggtctccatg 480
gtaggagtgc agctttctcc cccagttact ttccagctcc gggctggctc aggaccctg 540
ttctcagtg gccaggaacg ttatgaagca tcagacctaa cctgggagga ggaggaggaa 600
gaagaagggg agggagagga agagggaagag gaagatgatg aggatgagga tgcagatata 660
tctctggagg agcaaagccc tgtcaaacia gtcaaaaggc tgggtgcccc gaagcaggcg 720
agcgtggcta agaaaaaaaa gctggaaaaa gaagaagagg aaataagagc cagcgttaga 780
gacaagagcc ctgtgaaaaa ggccaaagcc acagccagag ccaagaagcc aggattcaag 840
aaatgaggag ccagcccttg gggggcacgg tgcaaaagtgg gccttccctg ggctgtgctg 900
caggcacagg gtgcccctgt ccagcccctc cacctgtgtc tgaatgcaac aggggtgttg 960
cgggggcaac atgagagccc ctcaccccca actctccact ttcaggaggc cccagtgaa 1020
gagccccacc tcgggtcaca ataagtgt 1048

```

<210> 107

<211> 1349

<212> DNA

<213> Homo sapiens

<220>

<221> misc-feature

<223> Incyte ID No.: 3573661CB1

<400> 107

```

gagaaagcag agctatttca agagtgaagg acagaaggga atccaggagg catctaagcg 60
aggaagggtc tacaggcagt gagtgaaggc caggagcagg gcccaggcca ggcacgacca 120
ccgaggggat gaacttcaca gtgggtttca agccgctgct aggggatgca cacagcatgg 180
acaacctgga gaagcagctc atctgcccga tctgcctgga gatgttctcc aaaccagtgg 240
tgatcttgcc ctgccaacac aacctgtgcc gcaaatgtgc caacgacgtc ttccaggcct 300
cgaatcctct atggcagtc cggggctcca ccactgtgtc ttcaggaggc cgtttccgct 360
gcccatcgtg caggcatgag gttgtcctgg acagacacgg tgtctacggc ctgcagcgaa 420
acgtgctagt ggagaacatt atcgacattt acaagcagga gtcattccaag ccgctgcact 480
cgaaggctga gcagcacctc atgtgcgagg agcatgaaga agagaagatc aatatttact 540
gcctgagctg tgaggtgccc acctgctctc tctgcaaggc cttcgggtgc cacaaggact 600
gtgaggtggc cccactgccc accatttaca aacgccagaa gaggtagctc agcgatggca 660
tcgcgatgct ggtggcaggc aatgaccgcg tgcaagcagt gatcacacag atggaggagg 720
tgtgccagac tatcgaggac aatagccgga ggcagaagca gttgttaaac cagaggtttg 780
agagcctgtg cgcagtgtc gaggagcgca agggtagct gctgcaggcg ctggcccggy 840
agcaagagga gaagctgcag cgcgtccgcg gcctcatccg tcagtatggc gaccacctgg 900
aggcctcctc taagctgggt gagtctgcca tccagtccat ggaagagcca caaatggcgc 960

```

```

tgtatctcca gcaggccaag gagctgatca ataagggtcgg ggccatgtcg aagggtggagc 1020
tggcaggggcg gccggagcca ggctatgaga gcatggagca attcaccgta aggggtggagc 1080
acgtggccga aatgctgcgg accatcgact tccagccagg cgcttccggg ggaggaagag 1140
gagggtggccc cagacggaag aagagggcaa cccggggggc ggaagaaaaa acggcccgga 1200
tggggcctta taggcctttg cgcccgaacc ccgacccctt gcttcgaaaa agccccgggc 1260
gcttaagaat ttccggggga aggaattctt gccgcaaaaa aaccccggca agcttttaac 1320
ccccaaaaat tccggggcgc ccggggccc 1349

```

<210> 108
 <211> 1642
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3633422CB1

```

<400> 108
ctagttctag atcgcgagcg cccgcccagg caccaccag cggcgccgac ctcagcgcg 60
acctatgggc tcgctaccag gacatgcgga gactggtgca cgacctcttg ccccccagg 120
tctgcagtct cctgaaccca gcagccatct acgccaacaa cgagatcagc ctgctgagc 180
ttgaggtcta cggctttgac tacgactaca ccctggccca gtatgcagac gcaactgacc 240
ccgagatctt cagtaccgcc cgtgacatcc tgcagagca ctacaagtac ccagaaggga 300
ttcggaagta tgactacaac cccagctttg ccatcctggt cctccactat gacattcaga 360
agagccttct gatgaagatt gacgccttcc actacgtgca gctggggaca gcctacaggg 420
gcctccagcc tgtgccagac gaggagggtga ttgagctgta tgggggtacc cagcacatcc 480
cactatacca gatgagtggc ttctatggca aggtccctc cattaagcag ttcattggaca 540
tcttctcgct accggagatg gctctgctgt cctgtgtggt ggactacttt ctgggccaca 600
gcctggagtt tgaccaagca catctctaca aggacgtgac ggacgccatc cgagacgtgc 660
atgtgaaggg cctcatgtac cagtggatcg agcaggacat ggagaagtac atcctgagag 720
gggatgagac gtttgctgtc ctgagccgcc tgggtggcca tgggaaacag ctgttctca 780
tcaccaacag tcttttcagc ttctagaca aggggatgcg gcacatggtg ggtcccatt 840
ggcgccagct cttcgatgtg gtcattgtcc aggcagacaa gccagcttc ttcactgacc 900
ggcgcaagcc tttcagaaaa ctcgatgaga agggctcact tcagtgggac cggatcacc 960
gcttgaaaaa gggcaagatg tatcggcagg gaaacctgtt tgacttctta cgcttgacgg 1020
aatggcggtg ccccgcggtg ctctacttcg gggaccacct ctatagtgat ctggcggtac 1080
tcatgctgcg gcacggctgg cgcacaggcg ccatcatccc cgagctggag cgtgagatcc 1140
gcatcatcaa cacggagcag tacatgcact cgctgacgtg gcagcaggcg ctcacggggc 1200
tgctggagcg catgcagacc tatcaggacg cggagtgcag gcaggtgctg gctgcctgga 1260
tgaaagagcg gcaggagctg aggtgcatca ccaaggccct gttcaatgcg cagttcggca 1320
gcatcttcg caccttccac aaccccacct acttctcaag gcgcctctg cgcttctctg 1380
acctctacat ggcctccctc agctgcctgc tcaactaccg cgtggacttc acctctacc 1440
cacgccgtac gccgctgcag cacgaggcac cctctggat ggaccagctc tgcacgggct 1500
gcatgaagac ccccttccct ggtgacatgg ccacatccg ctgagggcac ctttattgtc 1560
tgggacaggc cctcagcccc tctgccccca tccaccaga caagcaataa aagtgtgtc 1620
ctcctgaaa aaaaaaaaaa aa 1642

```

<210> 109
 <211> 1818
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc-feature
 <223> Incyte ID No.: 3993377CB1

```

<400> 109
gaagactctc tggagccctt caactctctg gcaccagagc cagtgagtgg aggactatat 60
ggtattgatg acacggagct gatgggtgca gaggacaagc tgctcttgga ggacagccct 120
gtgattgctg cccttgattg cccttccctc aataatgcta ctgccttcag tctcctggca 180
gatgatagtc aaacatcaac ctctatcttt gccagtccca cctctccacc tgtcctaggg 240

```

```

gagtctgtcc tgcaagataa cagctttgac ctgaataatg gtagtgacgc tgaacaggaa 300
gaaatggaaa ctcaatcttc agacttccca ccatccctga cccagccagc tcctgatcag 360
tcatccacta ttcagctaca tccagcaacc tcaccagcag tctcgccaac aacctcccca 420
gcagctctcc tagtggtttc tccagcagcc tccccagaaa tctctccaga agtttgtccc 480
gcagcttcta cagttgtctc tccagcagtc ttctcagtg tctctccagc ttctcagca 540
gtcctccagc cagttctctt agaagtcocg ttgacggctt cagtgacatc cccaaaagcc 600
tctcccgtaa cttcccccagc agctgccttt ccaacagcct cccagcaaa taaggatgtc 660
agcagctttc tagaaaccac tgctgacgtg gaagagatca ctggagaagg actcactgct 720
tctggtagtg gtgatgtcat gaggagacgt attgctaccc cagaagaagt tcgtcttccc 780
ctccaacatg ggtggcggag agaggtgcgc atcaagaaca gcagccaccg atggcagggg 840
gagacctggt attatggccc ctgtgggaag aggatgaagc aatttccaga agtgatcaag 900
tacctgagcc gcaacgtggt acacagtgtc cgccgagagc acttcagctt cagtccccgt 960
atgcctgttg gagatttctt tgaagaaaga gacacgccag agggcttgca gtgggtgcag 1020
ctctcagcag aggagatccc gtctgaggatt caggcaatta ctggcaaacg ggttcgacct 1080
cgaaacactg agaaggctaa gactaaggaa gtccccagg tgaaacgggg tcgaggtcgg 1140
ccacctaaag tcaaaatcac tgagctattg aacaagacag agattgctaa aagcaagaag 1200
ctggaggccc aagaacatt gaatgaggag gataaagcaa agattgctaa aagcaagaag 1260
aagatgagc agaaggttca acggggagag tgctcagacta ctatccaagg gcaggccaga 1320
aataagcgga aacaagagac caagagctta aagcagaagg aagctaagaa gaaatccaag 1380
gctgagaaag aaaaaggaaa gacaaaagcag gaaaaactga aggaaaaagt caagagggaa 1440
aagaaggaga aggtaaaaat gaaggaaaag gaggaggtga ccaaagccaa gccagcctgt 1500
aaagcagata agaccctggc cacacagagg cgcttggagg aacggcagag gcagcagatg 1560
atcttggagg acatgaagaa gccgacagag gatattgtgtc tgactgacca ccagcccctg 1620
cctgacttct cagcagtcoc tggctctgaca ttgcccagtg gacgcttctc agactgcttg 1680
accattgttg agttcctgca tagctttggc aaggtgctgg gccttgatcc tgcccaaggt 1740
tgtgcctagc ctgggggggtc ctgcaggaag gggctcctgt gttcaagggt accactctgg 1800
gtgaagggtgc aaaacccc 1818

```

<210> 110

<211> 785

<212> DNA

<213> Homo sapiens

<220>

<221> unsure

<222> 738

<223> a or g or c or t, unknown, or other

<223> Incyte ID No.: 4717936CB1

<400> 110

```

gtctctgtag tcatgaggct gaaggggggtg gggacagtgt tgataaaagg cactagaggc 60
agctcccaca ccttctctcg gaactgttgc ccacatgcag ccccgacac agcccctagc 120
ccaaacctta ccttctctcc tcggaggggc cctcagagac actgggctgc ggtgacctgt 180
cattaagatg ggcacagggt gggagggctt ccagcggacc ctgaaggaaag tcgctacat 240
cctcctctgc tgctggtgta tcaaggact gctggcatcg ctcaggtggg cagggtagag taaacaggag 360
ctctcccccac cagcaccatg gctggcatcg ccttagcttc tcattctctt ctttagcccc 420
gcatagctgc agcttctgtg gcagagcttg ccttagcttc tcattctctt ctttagcccc 480
cagcccaatt gccatcaaga ctcctgaagc cagctgtgct tgaccaagga tgggccataa 540
acaatgagta aacagtaaag tgtggatcct gctttgagct gtgtcatcta gcagacctgc 600
ctcatctctg agcctccttc attcccaacc cctgcctctg gtggacctct ggtgggcagg 660
agcttggact gctctggcta ggacccagc aagattgtgg caagacctgg cactcctcca 720
agcttggcac agtgagccca ccagctcaga tggttgatct taccataccc tcatagtacc 780
aagaatggac tgccccnta agaaacctgt ttgagaatca ctgaattata aatataaacc 840
cagat 875

```